

# Global, regional and national burden of dietary iron deficiency from 1990 to 2021: a Global Burden of Disease study

Received: 23 September 2024

Accepted: 28 February 2025

Published online: 22 April 2025

 Check for updates


Sooji Lee<sup>1,918</sup>, Yejun Son<sup>2,918</sup>, Jiyoung Hwang<sup>3,918</sup>, Min Seo Kim<sup>4,5,918</sup>, GBD 2021 Dietary Iron Deficiency Collaborators\*, Jae Il Shin<sup>6,919</sup> , Dong Keon Yon<sup>7,919</sup>  & Nicholas. J. Kassebaum<sup>8,9,10</sup> 


Although iron deficiency is well documented, less is known about dietary involvement in symptomatic iron deficiency manifesting in medical conditions. In this study, we quantified the global burden of dietary iron deficiency, focusing on where inadequate dietary iron intake leads to clinical manifestations such as anemia. We analyzed data from the Global Burden of Diseases, Injuries, and Risk Factors Study (GBD) 2021 to estimate dietary iron deficiency prevalence and disability-adjusted life years (DALYs), stratified by age, sex, geography and socio-demographic index (SDI) across 204 countries. In 2021, global age-standardized prevalence and DALY rates were 16,434.4 (95% uncertainty interval (UI), 16,186.2–16,689.0) and 423.7 (285.3–610.8) per 100,000 population, with rates decreasing by 9.8% (8.1–11.3) and 18.2% (15.4–21.1) from 1990 to 2021. A higher burden was observed in female individual (age-standardized prevalence, 21,334.8 (95% UI, 20,984.8–21,697.4); DALYs, 598.0 (402.6–854.4)) than in male individual ((age-standardized prevalence, 11,684.7 (11,374.6–12,008.8); DALYs, 253.0 (167.3–371.0)). High-SDI countries presented greater improvement, with a 25.7% reduction compared to 11.5% in low-SDI countries. Despite global improvements, dietary iron deficiency remains a major health concern with a global prevalence of 16.7%, particularly affecting female individuals, children and residents in low-SDI countries. Urgent interventions through supplementation, food security measures and fortification initiatives are essential.

Iron deficiency is one of the most common micronutrient deficiencies, leading to iron deficiency anemia and causing a substantial disease burden worldwide<sup>1</sup>. Although it is considered relatively preventable with iron supplementation<sup>1</sup>, dietary iron deficiency is ranked as the eighth highest modifiable cause of years lived with disability (YLDs) in 2021 across all age groups, underscoring its global importance<sup>2</sup>. Previous studies highlighted infants, young children and pregnant people as particularly susceptible to dietary iron deficiency due to their high demand for supplementary iron<sup>3</sup>. Iron deficiency in

infants and children can impair brain development, metabolism and immune system development, and maternal iron deficiency is associated with an increased risk of preterm birth, low birth weight and adverse neonatal outcomes, leading to substantial associated disease burden and health consequences<sup>4,5</sup>.

Earlier research focused primarily on investigating the global burden of anemia<sup>4</sup>, with limited data on dietary iron deficiency at a global scale. Although one study estimated the global burden of micronutrient deficiency, it only provided prevalence rates of dietary

A full list of affiliations appears at the end of the paper.  e-mail: [shinji@yuhs.ac](mailto:shinji@yuhs.ac); [yonkkang@gmail.com](mailto:yonkkang@gmail.com)



**Fig. 1 | Global map of age-standardized prevalence and DALY rates of dietary iron deficiency and their percentage changes over time, both sexes combined, 2021. a,b.** Age-standardized prevalence and DALY rates for dietary iron deficiency (a) and percentage changes of age-standardized prevalence and DALY rates for dietary iron deficiency (b).

iron deficiency at the country level, with limited subgroup analysis for each sex and age group<sup>6</sup>. Given the known disparities in dietary iron deficiency across demographics, subgroup analysis is crucial to prioritize high-risk populations requiring urgent resource allocation and tailored strategies for effective prevention and management<sup>1</sup>. Moreover, previous studies on dietary iron deficiency often assessed disease burden based solely on iron levels, which may capture cases of iron deficiency without clinical symptoms or those of lesser clinical relevance, potentially overestimating the overall burden<sup>7,8</sup>.

This Global Burden of Disease (GBD) Study 2021 analysis focused on complicated dietary iron deficiency—specifically individuals presenting with anemia—to better capture the impact of such cases and identify high-priority populations for dietary interventions. We estimated the prevalence and burden (that is, DALYs) associated with dietary iron deficiency across 204 countries and territories from 1990 to 2021.

## Results


### Global burden of dietary iron deficiency

In 2021, the global age-standardized prevalence and DALY rates of dietary iron deficiency were 16,434.4 (95% UI, 16,186.2–16,689.0) and 423.7 (285.3–610.8) per 100,000 population, respectively (Supplementary Table 6). The all-age global prevalence of dietary iron deficiency was 16.7% (95% UI, 16.4–16.9). These country-level age-standardized rates and their percentage changes are presented in Fig. 1. The number of prevalent cases worldwide increased by 29.0%, from 984.6 (95% UI, 970.8–997.8) million in 1990 to 1,270.6 (1,252.4–1,290.7) million in 2021, as illustrated in Fig. 2. Detailed numbers of cases and age-standardized rates of prevalence and DALYs for each SDI region are presented in Supplementary Tables 7–11. In Tables 1 and 2, the percentage changes of age-standardized rates of prevalence and

DALYs for each decade are presented. The global age-standardized prevalence rate and DALY rate decreased by 9.8% (95% UI, 8.1–11.3) and 18.2% (15.4–21.1) from 1990 to 2021, respectively. From 2000 to 2010, the prevalence rate decreased by 5.0% (95% UI, 4.1–5.9), the most considerable reduction observed, whereas the lowest change was observed between 2019 and 2021, with a decline of 0.1% (–0.6 to 0.4). Similarly, the DALY rate was the most substantially reduced between 2000 and 2010 (8.1% (95% UI, 6.5–9.9)) and between 2010 and 2019 (8.6% (6.3–11.4)), with the smallest reduction between 2019 and 2021, at 1.0% (0.3–1.8).

### Burden of dietary iron deficiency according to age and sex

Global ranks and distributions of dietary iron deficiency according to each age group are depicted in Extended Data Figs. 1–7. Dietary iron deficiency ranked particularly high among those aged 5–14 years in low and lower-middle SDI countries. Subgroup analysis presented in Fig. 3 revealed that female individuals had higher prevalence rates than male individuals in age groups under 65 years. DALY rates were higher in female than in male individuals across all ages, particularly during reproductive years. In 2021, the age-standardized prevalence rates per 100,000 population were 21,334.8 (20,984.8–21,697.3) in female and 11,684.7 (11,374.6–12,008.8) in male individuals. Similarly, the age-standardized DALY rates per 100,000 population were higher in female individuals (598.0 (402.6–854.4)) than in male individuals (253.0 (167.2–370.9)) (Supplementary Tables 12 and 13). However, from 1990 to 2021, a greater decrease was observed among male individuals (20.6% (18.3–23.0)) compared to female individuals (2.6% (0.6–4.5)). Considerable differences were observed among age groups, with children aged 6–11 months and adults over 95 years showing substantial burdens compared to other age groups (Supplementary Table 14). Similar patterns were observed for each sex (Supplementary Tables 15 and 16).



**Fig. 2 | Numbers and rates of dietary iron deficiency, 1990–2021. a, b.** Number of prevalent cases and DALYs of dietary iron deficiency at the global level (a) and rates (per 100,000 population) of age-standardized and all-age prevalence and DALYs of dietary iron deficiency at the global level (b). The lines represent the

central estimate (median) of the number and rate. Uncertainty was incorporated at every stage of the estimation process by random sampling from possible value ranges for each estimate. The final UI for each result was determined using the 2.5th and 97.5th percentiles of the draws.

### Burden of dietary iron deficiency by SDI and region

There were substantial disparities in age-standardized dietary iron deficiency prevalence rates among SDI quintiles (Supplementary Tables 7–11). In 2021, lower SDI countries exhibited higher age-standardized prevalence rates per 100,000 population compared to higher SDI countries (low SDI, 27,301.1 (26,624.0–27,996.5); lower-middle SDI, 25,686.6 (25,141.4–26,256.7); middle SDI, 13,569.0 (13,297.7–13,840.1); higher-middle SDI, 8,010.2 (7,718.0–8,300.7); high SDI, 4,572.6 (4,338.6–4,812.6)). A similar pattern was observed in DALY rates (low SDI, 756.7 (507.9–1,079.0); lower-middle SDI, 701.7 (473.8–995.6); middle SDI, 325.7 (217.3–471.7); higher-middle SDI, 158.5 (105.2–230.7); high SDI, 71.1 (46.7–107.1)).

Figure 4 presents the male-to-female (M/F) ratio of dietary iron deficiency, comparing both prevalence and DALY rates across age groups and SDI levels. Female participants exhibited a markedly higher prevalence of dietary iron deficiency during childhood and

reproductive ages compared to male participants, with this disparity being most pronounced in high-SDI countries. Specifically, in high-SDI countries, the M/F prevalence ratio was 0.1 among the age group between 25 years and 29 years, whereas low-SDI countries exhibited a ratio of 0.3 for the same age group. This pattern shifted in older populations, where high-SDI countries showed a higher burden in male adults compared to female adults (M/F ratio of 1.7 for ages 80–84 years), whereas low-SDI countries (ratio of 1.1 for the same age group) maintained a relatively similar burden distribution between sexes. Similar patterns were also observed for DALY rates (Supplementary Tables 17 and 18).

Disparities in both age-standardized prevalence and DALY rates were observed across world regions (Supplementary Tables 19–21). The highest prevalence and DALY rates were observed in South Asia (prevalence per 100,000 population, 31,696.8 (95% UI, 31,046.8–32,415.5); DALYs, 885.5 (95% UI, 600.3–1,271.4)) and Western Sub-Saharan Africa

**Table 1 | Prevalence of dietary iron deficiency, 1990–2021**

	Percentage change in prevalence (95% UI)				
	1990–2021	1990–2000	2000–2010	2010–2019	2019–2021
Global (overall)					
Absolute number	29,050 (26,883 to 31,362)	8,998 (7,872 to 10,247)	7,624 (6,519 to 8,623)	11,934 (10,085 to 13,673)	1,350 (0.897 to 1.802)
Age-standardized rate	-9.762 (-11.256 to -8.109)	-3.318 (-4.287 to -2.287)	-4.982 (-5.919 to -4.116)	-2.145 (-3.832 to -0.468)	-0.092 (-0.550 to 0.376)
Sex					
Male					
Absolute number	12,081 (8,835 to 15,494)	5,993 (4,146 to 7,809)	2,040 (0,309 to 3,780)	4,433 (1,100 to 7,523)	0.013 (-0.894 to 0.903)
Age-standardized rate	-20.605 (-22.951 to -18.246)	-5.263 (-6.798 to -3.709)	-9.500 (-10,978 to -7,957)	-8.786 (-11,733 to -6,056)	-1.151 (-2.047 to -0.279)
Female					
Absolute number	40,503 (37,698 to 43,346)	11,026 (9,627 to 12,482)	11,222 (9,958 to 12,384)	16,381 (14,399 to 18,406)	2,084 (1,665 to 2,548)
Age-standardized rate	-2.577 (-4.503 to -0.610)	-2.153 (-3.312 to -0.947)	-2.128 (-3.203 to -1.120)	1.999 (0.167 to 3.878)	0.530 (0.110 to 1.006)
SDI					
Low SDI					
Absolute number	95,343 (88,846 to 102,106)	25,432 (22,873 to 28,172)	22,035 (19,202 to 24,961)	32,844 (28,398 to 37,279)	4,094 (2,923 to 5,190)
Age-standardized rate	-11.523 (-13.987 to -8.836)	-3.694 (-5.348 to -1.933)	-6.253 (-7.999 to -4.367)	-2.338 (-5.231 to 0.554)	-0.485 (-1.420 to 0.403)
Lower-middle SDI					
Absolute number	38,782 (34,797 to 43,389)	14,585 (12,495 to 16,932)	10,510 (8,602 to 12,494)	11,501 (8,297 to 14,431)	1,035 (0.307 to 1.803)
Age-standardized rate	-14.993 (-17,178 to -12,483)	-4.016 (-5.567 to -2.340)	-6.008 (-7,444 to -4,446)	-6.814 (-9,250 to -4,519)	-1.105 (-1,771 to -0.385)
Middle SDI					
Absolute number	9,748 (7,324 to 12,229)	3,357 (1,797 to 4,896)	2,265 (0,857 to 3,636)	4,615 (2,600 to 6,848)	0.302 (-0.273 to 0.864)
Age-standardized rate	-22.017 (-23,733 to -20,340)	-7.522 (-8,906 to -6,237)	-9.366 (-10,608 to -8,129)	-8.294 (-10,124 to -6,215)	-0.951 (-1,558 to -0.384)
Higher-middle SDI					
Absolute number	-19,825 (-22,632 to -16,572)	-3,535 (-5,599 to -1,286)	-10,763 (-12,739 to -8,711)	-8,089 (-11,053 to -4,583)	-1,341 (-2,265 to -0.414)
Age-standardized rate	-34.429 (-36,910 to -31,689)	-9.773 (-11,781 to -7,620)	-17.901 (-19,999 to -15,816)	-13.763 (-16,757 to -9,972)	-1.215 (-2,223 to -0.193)
High SDI					
Absolute number	-3,863 (-10,933 to 3,880)	-14,041 (-18,852 to -9,637)	6,276 (2,254 to 11,008)	6,246 (-0,102 to 12,844)	0.693 (-0,848 to 2,284)
Age-standardized rate	-25.749 (-31,078 to -19,700)	-20.191 (-24,397 to -16,301)	-3.845 (-7,750 to 0.316)	-3.845 (-9,819 to 2,948)	-0.538 (-2,293 to 1,231)

(prevalence, 26,171.5 (95% UI, 25,195.0–27,200.2); DALYs, 708.8 (95% UI, 461.7–1,016.5)), respectively. The greatest reduction in prevalence rate was observed in East Asia with a decline of 53.9% (52.7–55.1). Western Sub-Saharan Africa was the only region with a possible increasing trend in prevalence rate (4.9% (95% UI, -1.4 to 10.9)) compared to other regions. The percentage change for each region is presented in Supplementary Tables 22 and 23.

## Discussion

We investigated the global, regional and national burden of dietary iron deficiency using estimates from GBD 2021. The total number of people with dietary iron deficiency increased from 1990 to 2021, likely due to population growth, as age-standardized prevalence rates


decreased during the same period. A decrease in age-standardized DALY rates for dietary iron deficiency over time might suggest improvements in nutritional deficiency management. Female participants generally showed higher prevalence and DALY rates than male participants, with the reduction in prevalence rates for female participants over time being much smaller than male participants. Those aged 6–11 months particularly had a higher burden compared to other age groups for both sexes. Moreover, individuals from low-SDI countries exhibited a higher burden of dietary iron deficiency. These findings underscore the urgent need for targeted interventions to address the persistent burden of dietary iron deficiency, particularly among female individuals, young children and populations in low-SDI countries.

**Table 2 | DALYs associated with dietary iron deficiency, 1990–2021**

	Percentage change in DALYs (95% UI)				
	1990–2021	1990–2000	2000–2010	2010–2019	2019–2021
Global (overall)					
Absolute number	13,763 (9,907 to 17,608)	7,133 (5,321 to 9,034)	2,900 (0,937 to 4,720)	3,870 (1,001 to 6,432)	0,164 (–0,571 to 0,846)
Age-standardized rate	–18,194 (–21,089 to –15,413)	–4,029 (–5,563 to –2,404)	–8,096 (–9,862 to –6,479)	–8,630 (–11,357 to –6,269)	–1,040 (–1,817 to –0,333)
Sex					
Male					
Absolute number	–9,687 (–15,248 to –3,261)	1,925 (–1,041 to 5,096)	–6,084 (–9,877 to –2,632)	–6,539 (–12,581 to –1,029)	–1,668 (–3,361 to –0,227)
Age-standardized rate	–32,001 (–36,127 to –27,471)	–6,777 (–9,364 to –4,220)	–14,949 (–18,313 to –11,962)	–16,918 (–22,450 to –11,985)	–2,177 (–3,841 to –0,767)
Female					
Absolute number	27,285 (23,164 to 31,640)	10,136 (7,979 to 12,149)	7,695 (5,803 to 9,531)	8,732 (5,906 to 11,627)	0,933 (0,171 to 1,663)
Age-standardized rate	–10,518 (–13,610 to –7,454)	–2,589 (–4,464 to –0,791)	–4,489 (–6,189 to –2,777)	–4,555 (–7,224 to –1,860)	–0,530 (–1,276 to 0,209)
SDI					
Low SDI					
Absolute number	69,446 (58,228 to 80,568)	19,918 (15,034 to 24,525)	17,063 (11,752 to 22,024)	24,677 (18,051 to 31,333)	2,837 (1,346 to 4,303)
Age-standardized rate	–21,962 (–26,053 to –17,656)	–8,327 (–11,468 to –5,575)	–9,355 (–12,566 to –6,082)	–7,133 (–11,464 to –2,749)	–1,383 (–2,671 to –0,216)
Lower-middle SDI					
Absolute number	14,567 (8,001 to 20,954)	10,118 (7,005 to 13,632)	3,768 (0,599 to 6,968)	0,491 (–4,116 to 5,024)	–0,764 (–2,030 to 0,524)
Age-standardized rate	–28,029 (–31,919 to –24,335)	–7,182 (–9,671 to –4,545)	–10,927 (–13,365 to –8,510)	–15,236 (–18,955 to –11,721)	–2,648 (–3,778 to –1,478)
Middle SDI					
Absolute number	–4,461 (–8,018 to –0,694)	1,364 (–0,714 to 3,293)	–2,006 (–3,977 to 0,010)	–4,475 (–7,550 to –1,539)	–0,861 (–1,633 to –0,103)
Age-standardized rate	–29,420 (–32,204 to –26,600)	–8,211 (–10,038 to –6,663)	–11,654 (–13,541 to –9,922)	–15,439 (–18,267 to –12,761)	–1,821 (–2,610 to –1,055)
Higher-middle SDI					
Absolute number	–31,821 (–35,802 to –27,884)	–5,916 (–8,925 to –3,070)	–18,533 (–21,383 to –15,896)	–13,183 (–16,734 to –9,320)	–1,446 (–2,482 to –0,355)
Age-standardized rate	–43,702 (–46,865 to –40,316)	–11,520 (–14,315 to –8,814)	–24,340 (–26,931 to –22,036)	–19,129 (–22,902 to –15,491)	–1,387 (–2,492 to –0,112)
High SDI					
Absolute number	–6,470 (–12,819 to 0,244)	–12,024 (–15,597 to –8,706)	3,456 (–0,784 to 7,464)	3,102 (–3,619 to 9,555)	1,231 (–0,146 to 2,639)
Age-standardized rate	–27,549 (–32,203 to –22,470)	–18,422 (–21,708 to –15,359)	–6,168 (–10,028 to –2,414)	–6,531 (–12,225 to –0,610)	–0,038 (–1,522 to 1,280)

Previous studies predominantly focused on all-cause anemia or all-cause iron deficiency, with limited studies specifically addressing dietary iron deficiency. From a public health perspective, understanding dietary iron deficiency is crucial as it represents a preventable burden that can be addressed through feasible interventions. A recent pooled iron level analysis revealed that iron deficiency rates among preschool-aged children were higher than 20% in 13 countries and 10–19% in 8 other countries<sup>7</sup>. Similarly, for non-pregnant reproductive-age people, 10 datasets indicated prevalence rates exceeding 20%, with six datasets showing between 10% and 19%<sup>7</sup>. However, this analysis was limited by its exclusion of men, school-aged children and older adults and coverage of only 22 countries with median data collection in 2013. Another study by Passarelli et al.<sup>8</sup> estimated the

global prevalence of dietary iron deficiency to be 65%—substantially higher than the 16.7% found in our analysis. This large variation likely reflects differences in scope and case definitions; Passarelli et al.<sup>8</sup> accounted for both complicated and uncomplicated iron deficiency cases by analyzing iron level data, whereas this GBD centered on dietary iron deficiency related to anemia, representing symptomatic or complicated cases. We aimed to estimate the global burden of complicated dietary iron deficiency to better capture its impact and identify high-priority populations for dietary interventions. Despite these differences, both our analysis and that of Passarelli et al.<sup>8</sup> consistently showed higher prevalence among female individuals and in lower SDI regions, particularly in South Asian and sub-Saharan African countries. The convergence of findings across different methodologies



**Fig. 3 | Numbers and age-specific rates (per 100,000 population) of prevalence and DALYs from dietary iron deficiency at the global level by age group and sex, 2021.** The bars and lines represent the central estimate (median) of the number and rate, respectively. The error bars represent uncertainty, which was

incorporated at every stage of the estimation process by random sampling from possible value ranges for each estimate. The final uncertainty interval for each result was determined using the 2.5th and 97.5th percentiles of the draws.

underscores the importance of urgent global action to reduce dietary iron deficiency, especially in at-risk populations.


From 1990 to 2021, the age-standardized prevalence rate of dietary iron deficiency declined worldwide, a trend attributable to several factors. The inclusion of anemia in the United Nations' Sustainable Development Goals (SDGs) prompted substantial and widespread policy actions<sup>9</sup>. Nutrition programs, such as iron supplementation for pregnant people and children, as well as food fortification with iron-containing multiple micronutrient powders, were recommended by the World Health Organization (WHO)<sup>10</sup>. These policies likely contributed to the reduction in the prevalence of dietary iron deficiency, particularly in low- and middle-income countries (LMICs) with high rates of severe anemia<sup>11</sup>. Furthermore, studies indicated that countries experiencing economic growth have seen increased access to animal-source foods and implementation of iron fortification programs, resulting in improvements in nutritional status<sup>12</sup>. The correlation between diversifying food supplies and lower rates of anemia, malnutrition and stunting was suggested<sup>12</sup>. This could account for the higher prevalence of anemia from dietary iron deficiency in lower SDI countries, where food supply and diversity are often limited<sup>13</sup>.

Substantial differences between sexes and age groups were also observed in our study. Female individuals had higher prevalence rates than male individuals in age groups under 65 years, and DALY rates were higher in female individuals across all ages, particularly during reproductive years. One possible reason for the sex differences in the prevalence of dietary iron deficiency is that female individuals, particularly those of reproductive age, face higher demands for iron due to menstrual blood loss and childbirth<sup>6</sup>; previous studies indicated that iron stores in pre-menopausal women are substantially influenced by menstrual blood loss, requiring more iron intake to sustain essential body function<sup>14</sup>. In addition to menstrual blood loss, sex-hormonal influences also play a critical role in iron metabolism. Estrogen, which rises during the reproductive years, is known to affect iron absorption and distribution, further emphasizing why women of reproductive age require higher dietary iron intake<sup>3</sup>. Similarly, pregnancy imposes an additional burden on iron stores as fetal development significantly increases iron demands<sup>14</sup>. Such life stages create fluctuations in body iron demand throughout a woman's life, potentially increasing vulnerability to even marginal changes in dietary iron intake. This is distinct from cases where pathological conditions, such as heavy menstrual bleeding or other menstrual disorders, are the primary cause of iron deficiency, which are categorized separately in our analysis. In addition to the increased

physiological iron requirements of female individuals of reproductive age, a range of complex and interrelated factors may contribute. Studies suggest that variations in dietary access and food choices, influenced by socioeconomic contexts—such as unequal food distribution within households and limited dietary diversity—disproportionately affect female individuals, particularly those from low-income households<sup>15</sup>.

Conversely, post-menopausal women typically experience a decrease in iron loss due to the cessation of menstruation, which can lead to relatively higher iron stores<sup>1</sup>. However, older adults, regardless of sex, may still face iron deficiency due to factors such as reduced dietary intake, chronic diseases or decreased gastrointestinal absorption. Therefore, although reproductive-age women are particularly vulnerable, age-related physiological changes also substantially impact iron status across the lifespan. In older age groups, we observed no significant sex differences in dietary iron deficiency, with a slightly higher prevalence rate in male adults. This pattern may be explained by age-related testosterone decline in male adults, which reduces hemoglobin and hematocrit levels and may increase iron demands<sup>16</sup>.

SDG 2, which primarily focuses on achieving zero hunger, also includes efforts to improve nutrition and address malnutrition, such as reducing the prevalence of anemia among women of reproductive age by 2030. This aligns with its broader objectives to enhance health outcomes for vulnerable populations<sup>9</sup>. However, our data suggest that progress toward this goal has been minimal, particularly in LMICs. The recent impacts of the COVID-19 pandemic have further exacerbated existing challenges. We think that this limited advancement can be attributed to several complex and interrelated factors. Global food supply chain disruptions, intensified by the pandemic, have impacted the availability and affordability of nutrient-rich foods in low-SDI countries. This impact has been particularly severe in LMICs, where existing food distribution infrastructure is often unstable<sup>17</sup>. Compared to high-income countries, healthy diets with sufficient nutrients are more difficult to afford in LMICs<sup>18</sup>. Furthermore, economic crises at national and household levels have led to increased food insecurity, making it challenging for people to afford a nutrient-adequate diet<sup>19</sup>. Fruits, vegetables and animal-source foods, which are rich in essential nutrients, including iron, remain comparatively costly sources of calories in contrast to starchy staples and cereals in LMICs<sup>20</sup>. These cost disparities often force families to rely heavily on less nutritious but more affordable food options, a scenario exacerbated during the pandemic<sup>17</sup>, which altogether may partly explain the minimal reduction in global dietary iron deficiency burden observed during this period



**Fig. 4 | The M/F ratio of prevalence and DALY rates (per 100,000 population) of dietary iron deficiency by SDI and age group, 2021.** M/F ratio of dietary iron deficiency burden in 2021 is shown across age groups and SDI quintiles. The left panel represents the M/F ratio of age-specific prevalence rates (per 100,000 population), and the right panel displays the M/F ratio of DALYs per 100,000

population. Rows indicate age groups, ranging from 6–11 months to more than 95 years, and columns correspond to SDI quintiles, from low to high. In both panels, color gradients reflect the M/F ratio, with darker red shades indicating higher M/F ratios and blue-green shades representing lower ratios.

(Tables 1 and 2). The situation is further complicated by armed conflicts in various regions, which have intensified food insecurity, trade routes, agriculture and disrupted health systems<sup>21</sup>. Climate change also influences food availability and nutritional quality as it impacts crop yields and food production patterns<sup>22</sup>.

Despite global efforts to reduce dietary iron deficiency, its prevalence has only slightly decreased, falling short of the SDG targets. This persistent challenge highlights ongoing disparities in global food security and resource allocation, reflecting the complex interplay of economic, environmental and political factors influencing nutrition

worldwide. Given the particular susceptibility of populations in LMICs to malnutrition and related health burdens, it is crucial to bolster global efforts to stabilize global food supply chains and enact targeted policies.

Recognizing the relationship between dietary iron deficiency and nutritious diet access highlights the need for social services to support those lacking such diets. Ensuring continued delivery of nourishing school meals through home delivery, take-home rations and vouchers when schools are closed might be crucial in some regions<sup>23</sup>. Universal free school meals are one of the options to support nutrition and academic equity for all children and adolescents<sup>24</sup>. Additionally, income support for low-income households can reduce food insecurity, underscoring the importance of social protection<sup>25</sup>. The provision of iron supplements to at-risk populations could be considered a straightforward public health approach. This can be accomplished through various strategies, including supplementation, fortification of staple foods and the use of multiple micronutrient powders<sup>26</sup>. Targeted promotions aimed at populations with inadequate iron intake can effectively alleviate the burden of iron deficiency. A previous study reported that prenatal iron supplementation can reduce maternal anemia by 70%<sup>27</sup>, suggesting the potential for implementation. However, challenges remain due to limited access to prenatal care or iron supplementation, particularly among individuals at low socioeconomic status<sup>28</sup>.

With 2030 approaching, it is clear that considerable efforts are still required to meet the broader SDG 2 target, within which reducing anemia is one of the sub-goals. The WHO has set a target to achieve a 50% reduction in the prevalence of anemia among women of reproductive age (15–49 years) by 2025, as part of its Global Nutrition Targets 2025 (ref. 9). Our analysis of the population-level burden of dietary iron deficiency can provide the insights required to appropriately tailor interventions at the national and regional levels in an effort to reduce the prevalence across all age and sex groups. Our estimates can have implications for determining funding allocations, shaping programmatic priorities and supporting advocacy initiatives, all of which are imperative to achieving the targets set by the SDGs.

This study has several limitations. It relies on estimates derived from GBD methodology, where there are potential biases that need to be considered. Because complete data on dietary iron deficiency were not available for all 204 countries, missing data were supplemented through the GBD modeling process. Consequently, the actual level of dietary iron deficiency in certain regions or countries may have been either overestimated or underestimated, and results may be presented with broad and overlapping uncertainty intervals. Therefore, when interpreting the estimates, it is crucial to carefully consider each country's or region's data collection environment and the potential for missing information, acknowledging that the true prevalence of dietary iron deficiency may be higher or lower than the reported estimates<sup>2</sup>. Our methodology did not capture non-anemic iron deficiency with normal hemoglobin concentration, which could potentially lead to an underestimation of the total disease burden. However, current evidence indicates that the most substantial adverse health outcomes of iron deficiency manifest when hemoglobin levels decrease to the point of anemia<sup>4,26,29</sup>, justifying our focus on symptomatic/complicated dietary iron deficiency. Moreover, when compared to other GBD disability weights, mild anemia has one of the lowest disability weights across all categories: mild anemia (0.004), moderate anemia (0.052) and severe anemia (0.149). This suggests that non-anemic iron deficiency, with presumably even lower disability weights, could minimally contribute to burden estimates. However, it would also substantially increase case counts, which could overestimate iron deficiency prevalence due to the potential inclusion of individuals without anemia. By limiting to anemic iron deficiency attributable to inadequate diet, our approach estimated clinically significant burden associated with a dietary component. Third, the current modeling framework falls short in fully distinguishing dietary deficiencies from

other causes due to the lack of richer and more integrated data sources that provide direct, simultaneous measurement of dietary intake and menstrual health factors as well as other contributing conditions for iron deficiency. Finally, this study did not account for the role of ethnicity in the burden of dietary iron deficiency. Previous research indicates significant variations in hemoglobin concentrations among different ethnic groups, with East Asians exhibiting higher concentrations than African Americans<sup>30</sup>.

This study also has several strengths. It estimated the burden of iron deficiency due to inadequate nutrition, facilitating a direct estimation of the iron deficiency burden that can potentially be prevented by diet interventions. Given that dietary iron deficiency is the largest contributor to anemia (Extended Data Fig. 8) and can be modified through relatively simple dietary measures, our targeted focus allows us to identify population groups that could benefit from accessible interventions, such as iron supplements. In contrast to previous studies that broadly encompassed non-anemic iron deficiency, we focused on iron deficiency manifesting as anemia to better capture the clinical impact of iron deficiency and prioritize food resources. Lastly, the data highlighted demographics most susceptible to dietary iron deficiency by examining disease burden across age groups and sexes, with potential insights for policymakers looking to tailor public health actions to specific populations.

In conclusion, dietary iron deficiency remained a prevalent and persistent global burden, with a global prevalence of 16.7%, despite a decreasing trend globally from 1990 to 2021. Dietary iron deficiency disproportionately affected women, children and populations residing in the African continent and in low-SDI countries. Given that dietary iron deficiency can be effectively managed through interventions such as iron supplements, balanced diets and fortified foods, urgent and universal actions are warranted. Further research should focus on integrating data from diverse sources, including food systems, dietary intake patterns and population health metrics, to gain a more comprehensive understanding of dietary iron deficiency.

## Online content

Any methods, additional references, Nature Portfolio reporting summaries, source data, extended data, supplementary information, acknowledgements, peer review information; details of author contributions and competing interests; and statements of data and code availability are available at <https://doi.org/10.1038/s41591-025-03624-8>.

## References

1. Pasricha, S.-R., Tye-Din, J., Muckenthaler, M. U. & Swinkels, D. W. Iron deficiency. *Lancet* **397**, 233–248 (2021).
2. Ferrari, A. J. et al. Global incidence, prevalence, years lived with disability (YLDs), disability-adjusted life-years (DALYs), and healthy life expectancy (HALE) for 371 diseases and injuries in 204 countries and territories and 811 subnational locations, 1990–2021: a systematic analysis for the Global Burden of Disease Study 2021. *Lancet* **403**, 2133–2161 (2024).
3. Stevens, G. A. et al. Global, regional, and national trends in haemoglobin concentration and prevalence of total and severe anaemia in children and pregnant and non-pregnant women for 1995–2011: a systematic analysis of population-representative data. *Lancet Glob. Health* **1**, e16–e25 (2013).
4. Kinyoki, D., Osgood-Zimmerman, A. E., Bhattacharjee, N. V., Kassebaum, N. J. & Hay, S. I. Anemia prevalence in women of reproductive age in low- and middle-income countries between 2000 and 2018. *Nat. Med.* **27**, 1761–1782 (2021).
5. Ohuma, E. O. et al. Association between maternal haemoglobin concentrations and maternal and neonatal outcomes: the prospective, observational, multinational, INTERBIO-21st fetal study. *Lancet Haematol.* **10**, e756–e766 (2023).

6. Han, X., Ding, S., Lu, J. & Li, Y. Global, regional, and national burdens of common micronutrient deficiencies from 1990 to 2019: a secondary trend analysis based on the Global Burden of Disease 2019 study. *EClinicalMedicine* **44**, 101299 (2022).
7. Stevens, G. A. et al. Micronutrient deficiencies among preschool-aged children and women of reproductive age worldwide: a pooled analysis of individual-level data from population-representative surveys. *Lancet Glob. Health* **10**, e1590–e1599 (2022).
8. Passarelli, S. et al. Global estimation of dietary micronutrient inadequacies: a modelling analysis. *Lancet Glob. Health* **12**, e1590–e1599 (2024).
9. Daru, J. Sustainable Development Goals for anaemia: 20 years later, where are we now? *Lancet Glob. Health* **10**, e586–e587 (2022).
10. Pasricha, S.-R. et al. Net benefit and cost-effectiveness of universal iron-containing multiple micronutrient powders for young children in 78 countries: a microsimulation study. *Lancet Glob. Health* **8**, e1071–e1080 (2020).
11. Csölle, I. et al. Health outcomes associated with micronutrient-fortified complementary foods in infants and young children aged 6–23 months: a systematic review and meta-analysis. *Lancet Child Adolesc. Health* **6**, 533–544 (2022).
12. Bentham, J. et al. Multi-dimensional characterisation of global food supply from 1961–2013. *Nat. Food* **1**, 70–75 (2020).
13. Willett, W. et al. Food in the Anthropocene: the EAT–Lancet Commission on healthy diets from sustainable food systems. *Lancet* **393**, 447–492 (2019).
14. Munro, M. G. et al. The relationship between heavy menstrual bleeding, iron deficiency, and iron deficiency anemia. *Am. J. Obstet. Gynecol.* **229**, 1–9 (2023).
15. Waage, J. et al. Changing food systems and infectious disease risks in low-income and middle-income countries. *Lancet Planet. Health* **6**, e760–e768 (2022).
16. Ferrucci, L. et al. Low testosterone levels and the risk of anemia in older men and women. *Arch. Intern. Med.* **166**, 1380–1388 (2006).
17. Bai, Y. et al. Retail prices of nutritious food rose more in countries with higher COVID-19 case counts. *Nat. Food* **3**, 325–330 (2022).
18. Hirvonen, K., Bai, Y., Headey, D. & Masters, W. A. Affordability of the EAT–Lancet reference diet: a global analysis. *Lancet Glob. Health* **8**, e59–e66 (2020).
19. Osendarp, S. et al. The COVID-19 crisis will exacerbate maternal and child undernutrition and child mortality in low- and middle-income countries. *Nat. Food* **2**, 476–484 (2021).
20. Laborde, D., Herforth, A., Headey, D. & de Pee, S. COVID-19 pandemic leads to greater depth of unaffordability of healthy and nutrient-adequate diets in low- and middle-income countries. *Nat. Food* **2**, 473–475 (2021).
21. The Lancet Global Health. Food insecurity will be the sting in the tail of COVID-19. *Lancet Glob. Health* **8**, e737 (2020).
22. Läderach, P., Pacillo, G., Thornton, P., Osorio, D. & Smith, D. Food systems for peace and security in a climate crisis. *Lancet Planet. Health* **5**, e249–e250 (2021).
23. Fore, H. H., Dongyu, Q., Beasley, D. M. & Ghebreyesus, T. A. Child malnutrition and COVID-19: the time to act is now. *Lancet* **396**, 517–518 (2020).
24. Cohen, J. F. W., Verguet, S., Giyose, B. B. & Bundy, D. Universal free school meals: the future of school meal programmes? *Lancet* **402**, 831–833 (2023).
25. Reeves, A., Loopstra, R. & Tarasuk, V. Family policy and food insecurity: an observational analysis in 142 countries. *Lancet Planet. Health* **5**, e506–e513 (2021).
26. Stevens, G. A. et al. National, regional, and global estimates of anaemia by severity in women and children for 2000–19: a pooled analysis of population-representative data. *Lancet Glob. Health* **10**, e627–e639 (2022).
27. Peña-Rosas, J. P., De-Regil, L. M., Garcia-Casal, M. N. & Dowswell, T. Daily oral iron supplementation during pregnancy. *Cochrane Database Syst. Rev.* **2015**, CD004736 (2015).
28. Thiele, L. R. et al. Direct dispensation of prenatal supplements with iron and anemia among pregnant people. *JAMA Netw. Open* **6**, e2332100 (2023).
29. Gardner, W. M. et al. Prevalence, years lived with disability, and trends in anaemia burden by severity and cause, 1990–2021: findings from the Global Burden of Disease Study 2021. *Lancet Haematol.* **10**, e713–e734 (2023).
30. Kang, W. et al. Ethnic differences in iron status. *Adv. Nutr.* **12**, 1838–1853 (2021).

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

© The Author(s), under exclusive licence to Springer Nature America, Inc. 2025

<sup>1</sup>Department of Medicine, Kyung Hee University College of Medicine, Seoul, South Korea. <sup>2</sup>Department of Precision Medicine, Kyung Hee University College of Medicine, Seoul, South Korea. <sup>3</sup>Center for Digital Health, Medical Science Research Institute, Kyung Hee University College of Medicine, Seoul, South Korea. <sup>4</sup>Cardiovascular Disease Initiative, Broad Institute of MIT and Harvard, Cambridge, MA, USA. <sup>5</sup>Cardiovascular Research Center, Massachusetts General Hospital, Boston, MA, USA. <sup>6</sup>Department of Pediatrics, Yonsei University College of Medicine, Seoul, South Korea. <sup>7</sup>Department of Pediatrics, Kyung Hee University, Seoul, South Korea. <sup>8</sup>Institute for Health Metrics and Evaluation, University of Washington, Seattle, WA, USA. <sup>9</sup>Department of Health Metrics Sciences, School of Medicine, University of Washington, Seattle, WA, USA. <sup>10</sup>Department of Anesthesiology & Pain Medicine, University of Washington, Seattle, WA, USA. <sup>918</sup>These authors contributed equally: Sooji Lee, Yejun Son, Jiyoung Hwang, Min Seo Kim. <sup>919</sup>These authors jointly supervised this work: Jae Il Shin, Dong Keon Yon. \*A list of authors and their affiliations appears at the end of the paper.

✉ e-mail: [shinji@yuhs.ac](mailto:shinji@yuhs.ac); [yonkkang@gmail.com](mailto:yonkkang@gmail.com)

### GBD 2021 Dietary Iron Deficiency Collaborators

**Sooji Lee<sup>1,918</sup>, Yejun Son<sup>2,918</sup>, Jiyoung Hwang<sup>3,918</sup>, Min Seo Kim<sup>4,5,918</sup>, Jae Il Shin<sup>6</sup>, Dong Keon Yon<sup>7,919</sup>, Nicholas J. Kassebaum<sup>8,9,10</sup>, Semagn Mekonnen Abate<sup>11</sup>, Yohannes Habtegiorgis Abate<sup>12</sup>, Samar Abd ElHafeez<sup>13</sup>, Sherief Abd-Elsalam<sup>14</sup>, Meriem Abdoun<sup>15,16</sup>, Rizwan Suliankatchi Abdulkader<sup>17</sup>, Auwal Abdullahi<sup>18,19</sup>, Mesfin Abebe<sup>20</sup>, Armita Abedi<sup>21</sup>, Alemwork Abie<sup>22</sup>, Olumide Abiodun<sup>23</sup>, Richard Gyan Aboagye<sup>24</sup>, Hassan Abolhassani<sup>25,26</sup>,**

Mohamed Abouzid<sup>27</sup>, Lucas Guimarães Abreu<sup>28</sup>, Hasan Abualruz<sup>29</sup>, Hana J. Abukhadajah<sup>30</sup>, Salahdein Aburuz<sup>31,32</sup>, Ahmed Abu-Zaid<sup>33,34</sup>, Lawan Hassan Adamu<sup>35,36</sup>, Mesafint Molla Adane<sup>37</sup>, Isaac Yeboah Addo<sup>38,39</sup>, Oyelola A. Adegboye<sup>40</sup>, Victor Adekanmbi<sup>41</sup>, Charles Oluwaseun Adetunji<sup>42</sup>, Temitayo Esther Adeyeoluwa<sup>43,44</sup>, Qorinah Estiningtyas Sakilah Adnani<sup>45</sup>, Leticia Akua Adzigbli<sup>46</sup>, Saira Afzal<sup>47,48</sup>, Suneth Buddhika Agampodi<sup>49</sup>, Williams Agyemang-Duah<sup>50</sup>, Aqeel Ahmad<sup>51</sup>, Muayyad M. Ahmad<sup>52</sup>, Shahzaib Ahmad<sup>53,54</sup>, Tauseef Ahmad<sup>55</sup>, Ali Ahmed<sup>56,57</sup>, Ayman Ahmed<sup>58,59</sup>, Haroon Ahmed<sup>60</sup>, Mehrunnisha Sharif Ahmed<sup>61</sup>, Syed Anees Ahmed<sup>62</sup>, Marjan Ajami<sup>63</sup>, Karolina Akinosoglou<sup>64,65</sup>, Ema Akter<sup>66</sup>, Salah Al Awaidey<sup>67,68</sup>, Syed Mahfuz Al Hasan<sup>69</sup>, Omar Al Omari<sup>70</sup>, Muaaz M. Alajlani<sup>71</sup>, Ziyad Al-Aly<sup>72,73</sup>, Rasmieh Mustafa Al-amer<sup>74,75</sup>, Mohammed Albashtawy<sup>76</sup>, Wafa A. Aldhaleei<sup>77</sup>, Abdelazeem M. Algammal<sup>78</sup>, Fadwa Naji Alhalaiqa<sup>79</sup>, Abid Ali<sup>80</sup>, Akhtar Ali<sup>81</sup>, Liaqat Ali<sup>82</sup>, Mohammed Usman Ali<sup>83,84</sup>, Syed Shujait Ali<sup>85</sup>, Waad Ali<sup>86</sup>, Shohreh Alian Samakkhah<sup>87</sup>, Sheikh Mohammad Alif<sup>88,89</sup>, Sabah Al-Marwani<sup>90</sup>, Hesham M. Al-Mekhlafi<sup>91,92</sup>, Sami Almustanyir<sup>93,94</sup>, Jaber S. Alqahtani<sup>95</sup>, Rajaa M. Mohammad Al-Raddadi<sup>96</sup>, Mohammed A. Alsabri<sup>97,98</sup>, Zaid Altaany<sup>99</sup>, Awais Altaf<sup>100</sup>, Alaa B. Al-Tammemi<sup>101,102</sup>, Khalid A. Altirkawi<sup>103</sup>, Hany Aly<sup>104</sup>, Karem H. Alzoubi<sup>105,106</sup>, Reza Aman<sup>107,108</sup>, Sohrab Amiri<sup>109</sup>, Hubert Amu<sup>110</sup>, Ganiyu Adeniyi Amusa<sup>111,112</sup>, Robert Ancuceanu<sup>113</sup>, Tudorel Andrei<sup>114</sup>, Boluwatife Stephen Anuoluwa<sup>115</sup>, Iyadunni Adesola Anuoluwa<sup>116</sup>, Saeid Anvari<sup>117</sup>, Sumadi Lukman Anwar<sup>118</sup>, Anayochukwu Edward Anyasodor<sup>119</sup>, Jalal Arabloo<sup>120</sup>, Mosab Arafat<sup>121</sup>, Aleksandr Y. Aravkin<sup>8,9,122</sup>, Demelash Areda<sup>123,124</sup>, Brhane Berhe Aregawi<sup>125</sup>, Abdulfatai Aremu<sup>126</sup>, Hany Ariffin<sup>127,128</sup>, Mesay Arkew<sup>129</sup>, Ni Ketut Aryastami<sup>130,131</sup>, Mubarek Yesse Ashemo<sup>132,133</sup>, Jamila Abdulhamid Atata<sup>134</sup>, Seyyed Shamsadin Athari<sup>135</sup>, Maha Moh'd Wahbi Atout<sup>136</sup>, Alok Atreya<sup>137</sup>, Amlaku Mulat Aweke<sup>138</sup>, Mamaru Ayenew Awoke<sup>139</sup>, Beatriz Paulina Ayala Quintanilla<sup>140,141</sup>, Gulrez Shah Azhar<sup>142</sup>, Ahmed Y. Azzam<sup>143,144</sup>, Giridhara Rathnaiah Babu<sup>145</sup>, Nayereh Baghcheghi<sup>146</sup>, Nasser Bagheri<sup>147</sup>, Saeed Bahramian<sup>148</sup>, Atif Amin Baig<sup>149</sup>, Shankar M. Bakkannavar<sup>150</sup>, Senthilkumar Balakrishnan<sup>151</sup>, Wondu Feyisa Balcha<sup>138</sup>, Palash Chandra Banik<sup>152</sup>, Hansi Bansal<sup>153</sup>, Simachew Animen Bante<sup>138</sup>, Afisu Basiru<sup>154</sup>, Mohammad-Mahdi Bastan<sup>155,156</sup>, Kavita Batra<sup>157</sup>, Tahmina Begum<sup>158,159</sup>, Muhammad Bashir Bello<sup>160,161</sup>, Luis Belo<sup>162,163</sup>, Isabela M. Bensenor<sup>164</sup>, Girma Beressa<sup>165,166</sup>, Alemshet Yirga Berhie<sup>167</sup>, Devidas S. Bhagat<sup>168</sup>, Neeraj Bhala<sup>169,170</sup>, Nikha Bhardwaj<sup>171</sup>, Pankaj Bhardwaj<sup>172,173</sup>, Prarthna V. Bhardwaj<sup>174</sup>, Sonu Bhaskar<sup>175,176</sup>, Ajay Nagesh Bhat<sup>177</sup>, Jasvinder Singh Bhatti<sup>178</sup>, Bijit Biswas<sup>179</sup>, Trupti Bodhare<sup>180</sup>, Aadam Olalekan Bodunrin<sup>181</sup>, Archith Bloor<sup>182</sup>, Kaustubh Bora<sup>183</sup>, Hamed Borhany<sup>184</sup>, Souad Bouaoud<sup>185,186</sup>, Yasser Bustanji<sup>187,188</sup>, Mehtap Çakmak Barsbay<sup>189</sup>, Daniela Calina<sup>190</sup>, Luis Alberto Cámara<sup>191,192</sup>, Fan Cao<sup>193</sup>, Monica Cattafesta<sup>194</sup>, Luca Cegolon<sup>195,196</sup>, Francieli Cembranel<sup>197</sup>, Muthia Cenderadewi<sup>198,199</sup>, Kelly M. Cercy<sup>8</sup>, Chiranjib Chakraborty<sup>200,201</sup>, Sandip Chakraborty<sup>202</sup>, Rama Mohan Chandika<sup>203</sup>, Vijay Kumar Chattu<sup>204,205</sup>, Akhilanand Chaurasia<sup>206</sup>, Haowei Chen<sup>207</sup>, Gerald Chi<sup>208</sup>, Patrick R. Ching<sup>209</sup>, Jesus Lorenzo Chirinos-Caceres<sup>210</sup>, William C. S. Cho<sup>211</sup>, Bryan Chong<sup>212</sup>, Hitesh Chopra<sup>213</sup>, Sonali Gajanan Choudhari<sup>214</sup>, Dinh-Toi Chu<sup>215,216</sup>, Isaac Sunday Chukwu<sup>217</sup>, Federica Concina<sup>218</sup>, Michael H. Criqui<sup>219</sup>, Natalia Cruz-Martins<sup>220,221</sup>, Alanna Gomes da Silva<sup>222</sup>, Omid Dadras<sup>223,224</sup>, Emanuele D'Amico<sup>225</sup>, Samuel Demissie Darcho<sup>226</sup>, Saswati Das<sup>227</sup>, Nihar Ranjan Dash<sup>228</sup>, Shayom Debopadhaya<sup>229</sup>, Cristian Del Bo<sup>230</sup>, Ivan Delgado-Enciso<sup>231,232</sup>, Hardik Dineshbhai Desai<sup>233</sup>, Fikreab Desta<sup>234</sup>, Devananda Devegowda<sup>235</sup>, Amol S. Dhane<sup>236</sup>, Vishal R. Dhulipala<sup>237</sup>, Thanh Chi Do<sup>238</sup>, Thao Huynh Phuong Do<sup>239</sup>, Saeid Doaei<sup>240,241</sup>, Sushil Dohare<sup>242</sup>, Mario D'Oria<sup>195,243</sup>, Ojas Prakashbhai Doshi<sup>244</sup>, Robert Kokou Dowou<sup>46</sup>, Ashel Chelsea Dsouza<sup>245</sup>, Haneil Larson Dsouza<sup>246,247</sup>, Senbagam Duraisamy<sup>248</sup>, Sulagna Dutta<sup>249</sup>, Hisham Atan Edinur<sup>250</sup>, Ferry Efendi<sup>251</sup>, Aziz Eftekharmehrabad<sup>252,253</sup>, Foolad Eghbal<sup>254</sup>, Michael Ekholuenetale<sup>255</sup>, Ibrahim Farahat El Bayoumy<sup>256,257</sup>, Iffat Elbarazi<sup>258</sup>, Ghada Metwally Tawfik ElGohary<sup>259,260</sup>, Muhammed Elhadi<sup>261,262</sup>, Waseem El-Huneidi<sup>263</sup>, Mohammed Elshaer<sup>264</sup>, Ibrahim Elsohaby<sup>265,266</sup>, Theophilus I. Emeto<sup>267</sup>, Adeniyi Francis Fagbamigbe<sup>268,269</sup>, Ayesha Fahim<sup>270</sup>, Ildar Ravisovich Fakhradiyev<sup>271,272</sup>, MoezAlIslam Ezzat Mahmoud Faris<sup>273</sup>, Nelsensius Klau Fauk<sup>274,275</sup>, Patrick Fazeli<sup>276</sup>, Alireza Feizkhah<sup>277</sup>, Ginenus Fekadu<sup>265,278</sup>, Nuno Ferreira<sup>279</sup>, Florian Fischer<sup>280</sup>, Bobirca Teodor Florin<sup>281</sup>, Morenike Oluwatoyin Oluwátóyìn Folayan<sup>282,283</sup>, Artem Alekseevich Fomenkov<sup>284</sup>, Takeshi Fukumoto<sup>285</sup>, Blima Fux<sup>286</sup>, G. Sridevi<sup>287</sup>, Muktar A. Gadanya<sup>288,289</sup>, Abhay Motiramji Gaidhane<sup>290</sup>, Yaseen Galali<sup>291,292</sup>, Aravind P. Gandhi<sup>293</sup>, William M. Gardner<sup>8</sup>, Rupesh K. Gautam<sup>294</sup>, Miglas Welay Gebregergis<sup>295</sup>, Mesfin Gebrehiwot<sup>296</sup>, Tesfay B. B. Gebremariam<sup>297</sup>, Habtamu Geremew<sup>298</sup>, Lemma Getacher<sup>299</sup>, Genanew K. Getahun<sup>300</sup>, Molla Getie<sup>301</sup>, Simegnew Asmer Getie<sup>138</sup>, Delaram J. Ghadimi<sup>302</sup>, Khalid Yaser Ghailan<sup>242,303</sup>, Ramy Mohamed Ghazy<sup>304,305</sup>, Maryam Gholamalizadeh<sup>306</sup>, Ali Gholamrezanezhad<sup>307</sup>, Artyom Urievich Gil<sup>308</sup>, Bikash Ranjan Giri<sup>309</sup>, Alem Abera Girmay<sup>310</sup>, Alessandra C. Goulart<sup>311</sup>, Mohammed Ibrahim Mohialdeen Gubari<sup>312</sup>, Damitha Asanga Gunawardane<sup>313</sup>, Anish Kumar Gupta<sup>314,315</sup>, Ishita Gupta<sup>316,317</sup>, Rajat Das Gupta<sup>318,319</sup>, Sapna Gupta<sup>320</sup>, Vivek Kumar Gupta<sup>321</sup>, Farrokh Habibzadeh<sup>322</sup>, Parham Habibzadeh<sup>323</sup>, Najah R. Hadi<sup>324</sup>, Hailey Hagins<sup>8</sup>, Sobia Ahsan Halim<sup>325</sup>, Nadia M. Hamdy<sup>326</sup>, Samer Hamidi<sup>327</sup>, Alexis J. Handal<sup>328</sup>, Demelash Woldeyohannes Handiso<sup>133</sup>, Ahmed I. Hasaballah<sup>329</sup>, Md. Kamrul Hasan<sup>330,331</sup>, Hamidreza Hasani<sup>332</sup>, Md Saquib Hasnain<sup>333</sup>

Ikrama Ibrahim Hassan<sup>334,335</sup>, Hadi Hassankhani<sup>336,337</sup>, Simon I. Hay<sup>8,9</sup>, Khezay Hayat<sup>338,339</sup>, Golnaz Heidari<sup>340</sup>, Mehdi Hemmati<sup>341,342</sup>, Kamal Hezam<sup>343,344</sup>, Yuta Hiraike<sup>345</sup>, Nguyen Quoc Hoan<sup>346</sup>, Ramesh Holla<sup>347</sup>, Md Mahbub Hossain<sup>348,349</sup>, Mehdi Hosseinzadeh<sup>350,351</sup>, Mihaela Hostiuc<sup>352</sup>, Sorin Hostiuc<sup>353,354</sup>, Junjie Huang<sup>355</sup>, Tsegaye Gebreyes Hundie<sup>356</sup>, Javid Hussain<sup>357</sup>, Hong-Han Huynh<sup>358</sup>, Bing-Fang Hwang<sup>359,360</sup>, Segun Emmanuel Ibitoye<sup>361</sup>, Nayu Ikeda<sup>362</sup>, Olayinka Stephen Ilesanmi<sup>363,364</sup>, Irena M. Ilic<sup>365</sup>, Milena D. Ilic<sup>366</sup>, Mustapha Immurana<sup>367</sup>, Leeberk Raja Inbaraj<sup>368</sup>, Arit Inok<sup>369</sup>, Muhammad Iqhrammullah<sup>370</sup>, Lalu Muhammad Irham<sup>371</sup>, Md. Rabiul Islam<sup>372</sup>, Sheikh Mohammed Shariful Islam<sup>373</sup>, Nahlah Elkudssiah Ismail<sup>374,375</sup>, Gaetano Isola<sup>376</sup>, Chidozie Declan Iwu<sup>377</sup>, Assefa N. Iyasu<sup>310</sup>, Mahalaxmi Iyer<sup>378</sup>, J. Vinothini<sup>287</sup>, Abdollah Jafarzadeh<sup>379,380</sup>, Kasra Jahankhani<sup>381</sup>, Akhil Jain<sup>382</sup>, Ammar Abdulrahman Jairoun<sup>383</sup>, Mihajlo Jakovljevic<sup>384,385</sup>, Reza Jalilzadeh Yengejeh<sup>386</sup>, Safayet Jamil<sup>159,387</sup>, Talha Jawaid<sup>388</sup>, Krishnamurthy Jayanna<sup>389,390</sup>, Shubha Jayaram<sup>391</sup>, Belayneh Hamdela Jena<sup>392</sup>, Bijay Mukesh Jeswani<sup>393</sup>, Mohammad Jokar<sup>394,395</sup>, Jost B. Jonas<sup>396,397</sup>, Jobinse Jose<sup>398</sup>, Nitin Joseph<sup>399</sup>, Charity Ehimwenma Joshua<sup>400</sup>, Farahnaz Joukar<sup>401,402</sup>, Jacek Jerzy Jozwiak<sup>403</sup>, Ali Kabir<sup>404</sup>, Zubair Kabir<sup>405</sup>, Dler H. Hussein Kadir<sup>406,407</sup>, Farima Kahe<sup>408</sup>, Leila R. Kalankesh<sup>409</sup>, Arun Kamireddy<sup>410</sup>, Kehinde Kazeem Kanmodi<sup>411,412</sup>, Rami S. Kantar<sup>413,414</sup>, Jafar Karami<sup>415,416</sup>, Arman Karimi Behnagh<sup>417,418</sup>, Joonas H. Kauppila<sup>419,420</sup>, Navjot Kaur<sup>421</sup>, Gbenga A. Kayode<sup>422,423</sup>, Kefita Kashala Kayola<sup>424</sup>, Adera Debella Kebede<sup>425</sup>, Leila Keikavoosi-Arani<sup>426</sup>, Peter Njenga Keiyoro<sup>427</sup>, Himanshu Khajuria<sup>428</sup>, Amirmohammad Khalaji<sup>429,430</sup>, Nauman Khalid<sup>431,432</sup>, Alireza Khalilian<sup>433</sup>, Maseer Khan<sup>434</sup>, Vishnu Khanal<sup>435,436</sup>, Haitham Khatatbeh<sup>437</sup>, Moawiah Mohammad Khatatbeh<sup>438</sup>, Khalid A. Kheirallah<sup>439</sup>, Feriha Fatima Khidri<sup>440</sup>, Manoj Khokhar<sup>441</sup>, Atulya Aman Khosla<sup>53,442</sup>, Majid Khosravi<sup>443,444</sup>, Helda Khusun<sup>445,446</sup>, Yun Jin Kim<sup>447</sup>, Ruth W. Kimokoti<sup>448</sup>, Adnan Kisa<sup>449,450</sup>, Sezer Kisa<sup>451</sup>, Shivakumar KM Shivakumar<sup>452</sup>, Ali-Asghar Kolahi<sup>453</sup>, Farzad Kompani<sup>454</sup>, Hamid Reza Koohestani<sup>455</sup>, Soewarta Kosen<sup>456</sup>, Kewal Krishan<sup>457</sup>, Mohammed Kuddus<sup>458</sup>, Mukhtar Kulimbet<sup>459,460</sup>, Dewesh Kumar<sup>461</sup>, Nithin Kumar<sup>399</sup>, Vijay Kumar<sup>462,463</sup>, Satyajit Kundu<sup>464</sup>, Om P. Kurmi<sup>465,466</sup>, Frank Kyei-Arthur<sup>467</sup>, Chandrakant Lahariya<sup>468,469</sup>, Dharmesh Kumar Lal<sup>470</sup>, Iván Landires<sup>471,472</sup>, Kamaluddin Latief<sup>473,474</sup>, Nhi Huu Hanh Le<sup>475,476</sup>, Thao Thi Thu Le<sup>477</sup>, Munjae Lee<sup>478</sup>, Seung Won Lee<sup>479</sup>, Wei-Chen Lee<sup>480</sup>, An Li<sup>481,482</sup>, Ming-Chieh Li<sup>483</sup>, Stephen S. Lim<sup>8,9</sup>, Jialing Lin<sup>484</sup>, Xuefeng Liu<sup>485,486</sup>, Ashwini Lonimath<sup>487</sup>, José Francisco López-Gil<sup>488</sup>, Platon D. Lopukhov<sup>489</sup>, László Lorenzovici<sup>490,491</sup>, Paulo A. Lotufo<sup>492</sup>, Jailos Lubinda<sup>493,494</sup>, Hawraz Ibrahim M. Amin<sup>495,496</sup>, Zheng Feei Ma<sup>497</sup>, Elaheh Malakan Rad<sup>498</sup>, Ahmad Azam Malik<sup>499</sup>, Deborah Carvalho Malta<sup>500</sup>, Yosef Manla<sup>501</sup>, Emmanuel Manu<sup>110</sup>, Mirko Marino<sup>230</sup>, Miquel Martorell<sup>502,503</sup>, Roy Rillera Marzo<sup>504,505</sup>, Yasith Mathangasinghe<sup>506,507</sup>, Elezabeth Mathews<sup>508</sup>, Medha Mathur<sup>509</sup>, Andrea Maugeri<sup>225</sup>, Mohsen Mazidi<sup>510</sup>, Theresa A. McHugh<sup>8</sup>, Asim Mehmood<sup>242</sup>, Rahul Mehra<sup>511</sup>, Kala M. Mehta<sup>512</sup>, Tesfahun Mekene Meto<sup>513</sup>, Birye Dessalegn Mekonnen<sup>514</sup>, Hadush Negash Meles<sup>515</sup>, Endalkachew Belayneh Melese<sup>516,517</sup>, Max Alberto Mendez-Lopez<sup>518</sup>, Walter Mendoza<sup>519</sup>, Tomislav Mestrovic<sup>8,520</sup>, Chamila Dinushi Kukulege Mettananda<sup>521,522</sup>, Sachith Mettananda<sup>523,524</sup>, Tomasz Miazgowski<sup>525</sup>, Irmina Maria Michalek<sup>526,527</sup>, Le Huu Nhat Minh<sup>528,529</sup>, GK Mini<sup>530,531</sup>, Mojgan Mirghafourvand<sup>532</sup>, Awoke Misganaw<sup>8,9,533</sup>, Madhukar Mittal<sup>534</sup>, Ahmed Ismail Mohamed<sup>535,536</sup>, Jama Mohamed<sup>537</sup>, Nouh Saad Mohamed<sup>538,539</sup>, Ameen Mosa Mohammad<sup>540</sup>, Sakineh Mohammad-Alizadeh-Charandabi<sup>541,542</sup>, Saeed Mohammadi<sup>325,543</sup>, Shafiu Mohammed<sup>544,545</sup>, Ali H. Mokdad<sup>8,9</sup>, Hossein Molavi Vardanjani<sup>546</sup>, Lorenzo Monasta<sup>218</sup>, Mohammad Ali Moni<sup>547,548</sup>, AmirAli Moodi Ghalibaf<sup>549</sup>, Maryam Moradi<sup>254</sup>, Abbas Mosapour<sup>550,551</sup>, Rohith Motappa<sup>399</sup>, Amin Mousavi Khaneghah<sup>552,553</sup>, Sumaira Mubarik<sup>554,555</sup>, Sumoni Mukherjee<sup>556,557</sup>, Francesc Mulita<sup>558,559</sup>, Kavita Munjal<sup>560</sup>, Yanjinlkhani Munkhsaikhan<sup>561</sup>, Christopher J. L. Murray<sup>8,9</sup>, Ana-Maria Musina<sup>562,563</sup>, Ghulam Mustafa<sup>564,565</sup>, Sathish Muthu<sup>566,567</sup>, Muhammad Muzaffar<sup>568,569</sup>, Ahamarshan Jayaraman Nagarajan<sup>570,571</sup>, Pirouz Naghavi<sup>572</sup>, Gurudatta Naik<sup>573</sup>, Soroush Najdaghi<sup>574,575</sup>, Hastyar Hama Rashid Najmuldeen<sup>576</sup>, Kannothu Thazha Kuni Nandu<sup>577</sup>, Sreenivas Narasimha Swamy<sup>578</sup>, Delaram Narimani Davani<sup>579</sup>, Abdulqadir J. Nashwan<sup>579</sup>, Zuhair S. Natto<sup>580,581</sup>, Javid Nauman<sup>582,583</sup>, Muhammad Naveed<sup>584</sup>, Biswa Prakash Nayak<sup>428</sup>, Athare Nazri-Panjaki<sup>585</sup>, G. Takop Nchanji<sup>586,587</sup>, Amanuel Tebabal Nega<sup>22</sup>, Mulata H. Nega<sup>588</sup>, Ionut Negoii<sup>281,589</sup>, Ruxandra Irina Negoii<sup>590,591</sup>, Seyed Aria Nejadghaderi<sup>592,593</sup>, Henok Biresaw Netsere<sup>37,594</sup>, Georges Nguefack-Tsague<sup>595</sup>, Josephine W. Ngunjiri<sup>596</sup>, Duc Hoang Nguyen<sup>597,598</sup>, Hau Thi Hien Nguyen<sup>599,600</sup>, Nhan Nguyen<sup>601</sup>, Nhien Ngoc Y. Nguyen<sup>602,603</sup>, Phat Tuan Nguyen<sup>604</sup>, Van Thanh Nguyen<sup>605,606</sup>, Robina Khan Niazi<sup>607</sup>, Ali Nikoobar<sup>453</sup>, Efaq Ali Noman<sup>608,609</sup>, Shuhei Nomura<sup>610,611</sup>, Syed Toukir Ahmed Noor<sup>66,612</sup>, N. A. Nawsherwan<sup>613</sup>, Mehran Nouri<sup>614,615</sup>, Majid Nozari<sup>616</sup>, Chisom Adaobi Nri-Ezedi<sup>617</sup>, Dieta Nurrika<sup>618,619</sup>, Chimezie Igwegbe Nzopotam<sup>620</sup>, Ogochukwu Janet Nzopotam<sup>621,622</sup>, James Odhiambo Oguta<sup>623</sup>, Tolulope R. Ojo-Akosile<sup>624</sup>, Sylvester Reuben Okeke<sup>39,625</sup>, Akinkunmi Paul Okekunle<sup>626,627</sup>, Osaretin Christabel Okonji<sup>628</sup>, Andrew T. Olagunju<sup>629,630</sup>, Matthew Idowu Olatubi<sup>631</sup>, Ahmed Omar Bali<sup>632</sup>, Michal Ordak<sup>633</sup>, Alberto Ortiz<sup>634,635</sup>, Esteban Ortiz-Prado<sup>636</sup>, Uchechukwu Levi Osuagwu<sup>637,638</sup>, Amel Ouyahia<sup>639,640</sup>, Mayowa O. Owolabi<sup>641,642</sup>, P. A. Mahesh Padukudru<sup>643</sup>, Alicia Padron-Monedero<sup>644</sup>, Jagadish Rao Padubidri<sup>645</sup>, Songhomitra Panda-Jonas<sup>646</sup>, Paola Pani<sup>218</sup>, Paraskevi Papadopoulou<sup>647,648</sup>, Shahina Pardhan<sup>649</sup>,

Pragyan Paramita Parija<sup>650</sup>, Romil R. Parikh<sup>651</sup>, Seoyeon Park<sup>652</sup>, Roberto Passera<sup>653,654</sup>, Jay Patel<sup>655,656</sup>, Siddhartha Pati<sup>657</sup>, Shankargouda Patil<sup>658</sup>, Dimitrios Patoulis<sup>659</sup>, Shrikant Pawar<sup>660</sup>, Minjin Peng<sup>661</sup>, Prince Peprah<sup>662</sup>, Gavin Pereira<sup>663,664</sup>, Arokiasamy Perianayagam<sup>665</sup>, Norberto Perico<sup>666</sup>, Anil K. Philip<sup>667</sup>, Zahra Zahid Piracha<sup>668,669</sup>, Dimitri Poddighe<sup>670,671</sup>, Ramesh Poluru<sup>672</sup>, Naeimeh Pourtaheri<sup>673</sup>, Pranil Man Singh Pradhan<sup>674,675</sup>, Jalandhar Pradhan<sup>676</sup>, Manya Prasad<sup>677</sup>, Bharathi M. Purohit<sup>678</sup>, Jagadeesh Puvvula<sup>679</sup>, Nameer Hashim Qasim<sup>680</sup>, Ibrahim Qattea<sup>681</sup>, Asma Saleem Qazi<sup>82</sup>, R. Deepthi<sup>682</sup>, Venkatraman Radhakrishnan<sup>683</sup>, Pankaja Raghav<sup>172</sup>, Fakher Rahim<sup>684,685</sup>, Vafa Rahimi-Movaghar<sup>686</sup>, Md. Mosfequr Rahman<sup>687</sup>, Mohammad Hifz Ur Rahman<sup>688</sup>, Mosiur Rahman<sup>687</sup>, Muhammad Aziz Rahman<sup>689,690</sup>, Amir Masoud Rahmani<sup>691</sup>, Mohammad Rahmanian<sup>692</sup>, Wahid Rahmanian<sup>693</sup>, Setyaningrum Rahmawaty<sup>694</sup>, Adarsh Raja<sup>695</sup>, Sandesh Raja<sup>696</sup>, Vinoth Rajendran<sup>287</sup>, Prashant Rajput<sup>697</sup>, Koushik Ramachandra<sup>698</sup>, Mahmoud Mohammed Ramadan<sup>699,700</sup>, Shakthi Kumaran Ramasamy<sup>701</sup>, Kritika Rana<sup>702</sup>, Rishabh Kumar Rana<sup>703</sup>, Amey Rane<sup>704,705</sup>, Chyitra R. Rao<sup>706</sup>, Mithun Rao<sup>347</sup>, Sowmya J. Rao<sup>707</sup>, Mohammad-Mahdi Rashidi<sup>155,453</sup>, Ashkan Rasouli-Saravani<sup>381</sup>, Devarajan Rathish<sup>708</sup>, Salman Rawaf<sup>709,710</sup>, Rabail Zehra Raza<sup>82</sup>, Christian Razo<sup>8</sup>, Elrashdy M. Moustafa Mohamed Redwan<sup>711,712</sup>, Giuseppe Remuzzi<sup>666</sup>, Kannan RR Rengasamy<sup>713,714</sup>, Nazila Rezaei<sup>155</sup>, Mohsen Rezaeian<sup>715</sup>, Muhammad Riaz<sup>716</sup>, Jennifer Rickard<sup>717,718</sup>, Hermano Alexandre Lima Rocha<sup>719</sup>, Leonardo Roever<sup>720,721</sup>, Ravi Rohilla<sup>722</sup>, Luca Ronfani<sup>218</sup>, Moustaq Karim Khan Rony<sup>723</sup>, Himanshu Sekhar Rout<sup>724,725</sup>, Bedanta Roy<sup>726</sup>, Guilherme de Andrade Ruela<sup>727</sup>, Godfrey M. Rwegerera<sup>728</sup>, Aly M. A. Saad<sup>729</sup>, Cameron John Sabel<sup>342</sup>, Siamak Sabour<sup>730</sup>, Mamta Sachdeva Dhingra<sup>731</sup>, Basema Ahmad Saddik<sup>732,733</sup>, Umar Saeed<sup>734,735</sup>, Fatemeh Saheb Sharif-Askari<sup>736</sup>, Narjes Saheb Sharif-Askari<sup>228</sup>, Amirhossein Sahebkar<sup>737,738</sup>, Pragyan Monalisa Sahoo<sup>739</sup>, Soumya Swaroop Sahoo<sup>740</sup>, S. Mohammad Sajadi<sup>741</sup>, Afeez Abolarinwa Salami<sup>411,742</sup>, Luciane B. Salaroli<sup>743</sup>, Samreen Saleem<sup>744</sup>, Marwa Rashad Salem<sup>745</sup>, Yoseph Leonardo Samodra<sup>746,747</sup>, Vijaya Paul Samuel<sup>748</sup>, Abdallah M. Samy<sup>749,750</sup>, Rama Krishna Sanjeev<sup>751</sup>, Itamar S. Santos<sup>164,752</sup>, Sivan Yegnanarayana Iyer Saraswathy<sup>753</sup>, Aswini Saravanan<sup>754,755</sup>, Tanmay Sarkar<sup>756</sup>, Gargi Sachin Sarode<sup>757</sup>, Sachin C. Sarode<sup>757</sup>, Brijesh Sathian<sup>758,759</sup>, Anudeep Sathyanarayan<sup>245</sup>, Maheswar Satpathy<sup>760,761</sup>, Monika Sawhney<sup>762</sup>, Abu Sayeed<sup>763,764</sup>, Art Schuermans<sup>765,766</sup>, Siddharthan Selvaraj<sup>767,768</sup>, Mohammad H. Semreen<sup>769,770</sup>, Pallav Sengupta<sup>771</sup>, Sadaf G. Sepanlou<sup>772,773</sup>, Yashendra Sethi<sup>774</sup>, Christianus Heru Setiawan<sup>775</sup>, Mahan Shafie<sup>776</sup>, Samiah Shahid<sup>100,569</sup>, Wajeelah Shahid<sup>777</sup>, Moyad Jamal Shahwan<sup>778</sup>, Masood Ali Shaikh<sup>779</sup>, Alireza Shakeri<sup>780</sup>, Sunder Sham<sup>781</sup>, Muhammad Aaqib Shamim<sup>754</sup>, Mehran Shams-Beyranvand<sup>782</sup>, Anas Shamsi<sup>778,783</sup>, Alfiya Shamsutdinova<sup>784</sup>, Mohd Shanawaz<sup>785</sup>, Mohammed Shannawaz<sup>786</sup>, Medha Sharath<sup>245</sup>, Amin Sharifan<sup>787,788</sup>, Javad Sharifi Rad<sup>272</sup>, Anupam Sharma<sup>789</sup>, Vishal Sharma<sup>790</sup>, Maryam Shayan<sup>791,792</sup>, Rahim Ali Sheikhi<sup>793</sup>, Rekha Raghuvver Shenoy<sup>794</sup>, Mahabalesh Shetty<sup>795</sup>, Pavanchand H. Shetty<sup>645</sup>, Premalatha K. Shetty<sup>796</sup>, Shiran Shetty<sup>797</sup>, Rahman Shiri<sup>798</sup>, Aminu Shittu<sup>799</sup>, Velizar Shivarov<sup>800,801</sup>, Sunil Shrestha<sup>802</sup>, Emmanuel Edwar Siddig<sup>803,804</sup>, Luis Manuel Lopes Rodrigues Silva<sup>805,806</sup>, Baljinder Singh<sup>807</sup>, Harmanjit Singh<sup>808</sup>, Jasbir Singh<sup>809</sup>, Jasvinder A. Singh<sup>810,811</sup>, Narinder Pal Singh<sup>812</sup>, Paramdeep Singh<sup>813</sup>, Puneetpal Singh<sup>814</sup>, Surjit Singh<sup>754</sup>, Farrukh Sobia<sup>242</sup>, Ranjan Solanki<sup>815,816</sup>, Shipra Solanki<sup>817</sup>, Soroush Soraneh<sup>818,819</sup>, Reed J. D. Sorensen<sup>8,820</sup>, Suresh Kumar Srinivasamurthy<sup>821</sup>, Vetriselvan Subramanian<sup>822</sup>, Hani Susianti<sup>823,824</sup>, Chandan Kumar Swain<sup>724</sup>, Lukasz Szarpak<sup>825,826</sup>, Mindy D. Szeto<sup>827</sup>, T. Y. Sree Sudha<sup>828</sup>, Seyyed Mohammad Tabatabaei<sup>829,830</sup>, Seyed-Amir Tabatabaeizadeh<sup>831,832</sup>, Celine Tabche<sup>709</sup>, Jabeen Taiba<sup>833,834</sup>, Iman M. Talaat<sup>228,835</sup>, Mircea Tampa<sup>836,837</sup>, Jacques Lukenze Tamuzi<sup>838,839</sup>, Ingan Ukur Tarigan<sup>840</sup>, Md. Tariqujjaman<sup>841</sup>, Nathan Y. Tat<sup>842,843</sup>, Birhan Tsegaw Taye<sup>844</sup>, Yibekal Manaye Tefera<sup>845</sup>, Mohamad-Hani Temsah<sup>846</sup>, Rekha Thapar<sup>399</sup>, Samar Tharwat<sup>847</sup>, Sathish Thirunavukkarasu<sup>848</sup>, Nikhil Kenny Thomas<sup>849</sup>, Jansje Henny Vera Ticoalu<sup>850</sup>, Mariya Vladimirovna Titova<sup>851,852</sup>, Krishna Tiwari<sup>754</sup>, Marcos Roberto Tovani-Palone<sup>767</sup>, Nghia Minh Tran<sup>853</sup>, Ngoc Ha Tran<sup>854</sup>, Thang Huu Tran<sup>855,856</sup>, Domenico Trico<sup>857</sup>, Thien Tan Tri Tai Truyen<sup>858</sup>, Abdul Rohim Tualeka<sup>859</sup>, Munkhtuya Tumurkhuu<sup>860</sup>, Aniefiok John Udoakang<sup>861</sup>, Saeed Ullah<sup>862</sup>, Shahid Ullah<sup>863</sup>, Muhammad Umar<sup>864</sup>, Bhaskaran Unnikrishnan<sup>347</sup>, Tolassa Wakayo Ushula<sup>865</sup>, Seyed Mohammad Vahabi<sup>866</sup>, Sanaz Vahdati<sup>867</sup>, Asokan Govindaraj Vaithinathan<sup>868</sup>, Rohollah Valizadeh<sup>869</sup>, Jef Van den Eynde<sup>766</sup>, Joe Varghese<sup>870</sup>, Tommi Juhani Vasankari<sup>871,872</sup>, Balachandar Vellingiri<sup>873,874</sup>, Georgios-Ioannis Verras<sup>875,876</sup>, Dominique Vervoort<sup>877</sup>, Manish Vinayak<sup>878</sup>, Simona Ruxandra Volovat<sup>879,880</sup>, Yasir Waheed<sup>881,882</sup>, Yanzhong Wang<sup>883</sup>, Kosala Gayan Weerakoon<sup>884</sup>, Holly Wild<sup>89</sup>, Peter Willeit<sup>885,886</sup>, Abay Tadesse Woday<sup>887</sup>, Yihun Miskir Wubie<sup>888</sup>, Xiaoyue Xu<sup>733,889</sup>, Vikas Yadav<sup>890</sup>, Amir Yarahmadi<sup>891,892</sup>, Sanni Yaya<sup>893</sup>, Subah Abderehim Yesuf<sup>894,895</sup>, Muluken Yigezu<sup>896</sup>, Dehui Yin<sup>897</sup>, Naohiro Yonemoto<sup>898,899</sup>, Chuanhua Yu<sup>555</sup>, Burhan Abdullah Zaman<sup>900</sup>, Nelson Zamora<sup>901,902</sup>, Iman Zare<sup>903</sup>, Mohammed G. M. Zeariya<sup>329,904</sup>, Naod Gebrekrstos Zeru<sup>905,906</sup>, Haijun Zhang<sup>907,908</sup>, Liqun Zhang<sup>909,910</sup>, Claire Chenwen Zhong<sup>911</sup>, Abzal Zhumagaliuly<sup>912</sup>, Hafsa Zia<sup>913,914</sup>, Ghazal Zoghi<sup>915</sup>, Sa'ed H. Zyoud<sup>916,917</sup>, Nandita Perumal<sup>8,9</sup>, Leo Zoeckler<sup>8,9</sup>, Victor Vilchis-Tella<sup>8,9</sup> & Susan A. McLaughlin<sup>8,9</sup>

<sup>11</sup>Anesthesiology Department, Wollo University, Dessie, Ethiopia. <sup>12</sup>Department of Clinical Governance and Quality Improvement, Aleta Wondo General Hospital, Aleta Wondo, Ethiopia. <sup>13</sup>Department of Epidemiology, Alexandria University, Alexandria, Egypt. <sup>14</sup>Department of Tropical Medicine and Infectious Diseases, Tanta University, Tanta, Egypt. <sup>15</sup>Department of Medicine, University of Setif Algeria, Sétif, Algeria. <sup>16</sup>Department of Health, Sétif,

Algeria. <sup>17</sup>National Institute of Epidemiology, Indian Council of Medical Research, Chennai, India. <sup>18</sup>Department of Physiotherapy, Bayero University Kano, Kano, Nigeria. <sup>19</sup>Department of Physiotherapy, Federal University Wukari, Wukari, Nigeria. <sup>20</sup>Department of Midwifery, Dilla University, Dilla, Ethiopia. <sup>21</sup>Department of Emergency Medicine, Zanjan University of Medical Sciences, Zanjan, Iran. <sup>22</sup>Midwifery Department, Bahir Dar University, Bahir Dar, Ethiopia. <sup>23</sup>Department of Community Medicine, Babcock University, Ilishan-Remo, Nigeria. <sup>24</sup>Department of Family and Community Health, University of Health and Allied Sciences, Ho, Ghana. <sup>25</sup>Research Center for Immunodeficiencies, Tehran University of Medical Sciences, Tehran, Iran. <sup>26</sup>Department of Medical Biochemistry and Biophysics, Karolinska Institute, Stockholm, Sweden. <sup>27</sup>Department of Physical Pharmacy and Pharmacokinetics, Poznan University of Medical Sciences, Poznan, Poland. <sup>28</sup>Department of Pediatric Dentistry, Federal University of Minas Gerais, Belo Horizonte, Brazil. <sup>29</sup>Department of Nursing, Al Zaytoonah University of Jordan, Amman, Jordan. <sup>30</sup>Medical Research Center, Hamad Medical Corporation, Doha, Qatar. <sup>31</sup>Department of Pharmacology and Therapeutics, United Arab Emirates University, Al Ain, United Arab Emirates. <sup>32</sup>College of Pharmacy, University of Jordan, Amman, Jordan. <sup>33</sup>Department of Biochemistry and Molecular Medicine, Alfaisal University, Riyadh, Saudi Arabia. <sup>34</sup>College of Graduate Health Sciences, University of Tennessee, Memphis, TN, USA. <sup>35</sup>Department of Human Anatomy, Federal University Dutse, Dutse, Nigeria. <sup>36</sup>Department of Anatomy, Bayero University Kano, Kano, Nigeria. <sup>37</sup>College of Medicine and Health Sciences, Bahir Dar University, Bahir Dar, Ethiopia. <sup>38</sup>School of Medicine, University of Sydney, Sydney, New South Wales, Australia. <sup>39</sup>Centre for Social Research in Health, University of New South Wales, Sydney, New South Wales, Australia. <sup>40</sup>Menzies School of Health Research, Charles Darwin University, Darwin, Northern Territory, Australia. <sup>41</sup>Department of Obstetrics and Gynecology, The University of Texas Medical Branch, Galveston, TX, USA. <sup>42</sup>Department of Microbiology, Edo State University Uzairue, Iyamho, Nigeria. <sup>43</sup>Department of Pharmacology and Therapeutics, University of Medical Sciences, Ondo, Ondo, Nigeria. <sup>44</sup>Department of Veterinary Medicine, University of Ibadan, Ibadan, Nigeria. <sup>45</sup>Department of Public Health, Universitas Padjadjaran (Padjadjaran University), Bandung, Indonesia. <sup>46</sup>Department of Epidemiology and Biostatistics, University of Health and Allied Sciences, Ho, Ghana. <sup>47</sup>Department of Community Medicine, King Edward Memorial Hospital, Lahore, Pakistan. <sup>48</sup>Department of Public Health, Public Health Institute, Lahore, Pakistan. <sup>49</sup>Department of New Initiatives, International Vaccine Institute, Seoul, South Korea. <sup>50</sup>Department of Public Health Sciences, Queen's University, Kingston, Ontario, Canada. <sup>51</sup>College of Medicine, Shaqra University, Shaqra, Saudi Arabia. <sup>52</sup>School of Nursing, University of Jordan, Amman, Jordan. <sup>53</sup>Department of Medical Oncology, Miami Cancer Institute, Miami, FL, USA. <sup>54</sup>Department of Community Medicine and Preventive Health, King Edward Medical University Lahore, Lahore, Pakistan. <sup>55</sup>School of Public Health, Zhejiang University, Hangzhou, China. <sup>56</sup>Department of Pharmacy Practice, Riphah Institute of Pharmaceutical Sciences, Islamabad, Pakistan. <sup>57</sup>Division of Infectious Diseases and Global Public Health (IDGPH), University of California, San Diego, San Diego, CA, USA. <sup>58</sup>Institute of Endemic Diseases University of Khartoum, Khartoum, Sudan. <sup>59</sup>Swiss Tropical and Public Health Institute, University of Basel, Basel, Switzerland. <sup>60</sup>Department of Biosciences, COMSATS Institute of Information Technology, Islamabad, Pakistan. <sup>61</sup>College of Nursing, Majmaah University, Al Majmaah, Saudi Arabia. <sup>62</sup>Brody School of Medicine, East Carolina University, Greenville, NC, USA. <sup>63</sup>National Nutrition and Food Technology Research Institute, Shahid Beheshti University of Medical Sciences, Tehran, Iran. <sup>64</sup>Department of Internal Medicine, University of Patras, Patras, Greece. <sup>65</sup>Department of Internal Medicine and Infectious Diseases, University General Hospital of Patras, Patras, Greece. <sup>66</sup>Maternal and Child Health Division, International Centre for Diarrhoeal Disease Research, Dhaka, Bangladesh. <sup>67</sup>Department of Communicable Diseases, Ministry of Health, Muscat, Oman. <sup>68</sup>Middle East, Eurasia and Africa Influenza Stakeholders Network, Muscat, Oman. <sup>69</sup>Division of Public Health Sciences, Washington University in St. Louis, St. Louis, MO, USA. <sup>70</sup>Fundamentals and Administration Department, Sultan Qaboos University, Muscat, Oman. <sup>71</sup>Faculty of Pharmacy, Al-Sham Private University, Damascus, Syria. <sup>72</sup>Department of Research and Development, Washington University in St. Louis, St. Louis, MO, USA. <sup>73</sup>Clinical Epidemiology Center, US Department of Veterans Affairs (VA), St. Louis, MO, USA. <sup>74</sup>School of Nursing, Yarmouk University, Irbid, Jordan. <sup>75</sup>School of Nursing and Midwifery, Western Sydney University, Sydney, New South Wales, Australia. <sup>76</sup>Department of Community and Mental Health, Al al-Bayt University, Mafrqa, Jordan. <sup>77</sup>Division of Gastroenterology and Hepatology, Mayo Clinic, Jacksonville, FL, USA. <sup>78</sup>Department of Bacteriology, Immunology, and Mycology, Suez Canal University, Ismailia, Egypt. <sup>79</sup>College of Nursing, Qatar University, Doha, Qatar. <sup>80</sup>Department of Zoology, Abdul Wali Khan University Mardan, Mardan, Pakistan. <sup>81</sup>School of Agriculture, Food and Ecosystem Sciences, University of Melbourne, Parkville, Victoria, Australia. <sup>82</sup>Department of Biological Sciences, National University of Medical Sciences (NUMS), Rawalpindi, Pakistan. <sup>83</sup>Department of Medical Rehabilitation (Physiotherapy), University of Maiduguri, Maiduguri, Nigeria. <sup>84</sup>Department of Rehabilitation Sciences, Hong Kong Polytechnic University, Hong Kong, China. <sup>85</sup>Center for Biotechnology and Microbiology, University of Swat, Swat, Pakistan. <sup>86</sup>Department of Geography, Sultan Qaboos University, Muscat, Oman. <sup>87</sup>Department of Food Hygiene, Amol University of Special Modern Technologies, Amol, Iran. <sup>88</sup>Institute of Health and Wellbeing, Federation University Australia, Melbourne, Victoria, Australia. <sup>89</sup>School of Public Health and Preventive Medicine, Monash University, Melbourne, Victoria, Australia. <sup>90</sup>Independent Consultant, Amman, Jordan. <sup>91</sup>Department of Parasitology, University of Malaya, Kuala Lumpur, Malaysia. <sup>92</sup>Department of Parasitology, Sana'a University, Sana'a, Yemen. <sup>93</sup>College of Medicine, Alfaisal University, Riyadh, Saudi Arabia. <sup>94</sup>Ministry of Health, Riyadh, Saudi Arabia. <sup>95</sup>Department of Respiratory Care, Prince Sultan Military College of Health Sciences, Dammam, Saudi Arabia. <sup>96</sup>Department of Community Medicine, King Abdulaziz University, Jeddah, Saudi Arabia. <sup>97</sup>Department of Emergency Medicine, Sana'a University, Sanaa, Yemen. <sup>98</sup>Pediatric Emergency Medicine Department, Drexel University, Philadelphia, PA, USA. <sup>99</sup>Department of Basic Sciences, Yarmouk University, Irbid, Jordan. <sup>100</sup>Institute of Molecular Biology and Biotechnology, The University of Lahore, Lahore, Pakistan. <sup>101</sup>Research, Policy, and Training Directorate, Jordan Center for Disease Control, Amman, Jordan. <sup>102</sup>Applied Science Research Center, Applied Science Private University, Amman, Jordan. <sup>103</sup>University of Sharjah, Sharjah, United Arab Emirates. <sup>104</sup>Department of Pediatrics, Cleveland Clinic, Cleveland, OH, USA. <sup>105</sup>Department of Pharmacy Practice and Pharmacotherapeutics, University of Sharjah, Sharjah, United Arab Emirates. <sup>106</sup>Department of Clinical Pharmacy, Jordan University of Science and Technology, Irbid, Jordan. <sup>107</sup>Interdisciplinary Graduate Program in Human Toxicology, University of Iowa, Iowa City, IA, USA. <sup>108</sup>Holden Comprehensive Cancer Center, University of Iowa Hospitals and Clinics, Iowa City, IA, USA. <sup>109</sup>Spiritual Health Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran. <sup>110</sup>Department of Population and Behavioural Sciences, University of Health and Allied Sciences, Ho, Ghana. <sup>111</sup>Department of Medicine, University of Jos, Jos, Nigeria. <sup>112</sup>Department of Internal Medicine, Jos University Teaching Hospital, Jos, Nigeria. <sup>113</sup>Faculty of Pharmacy, Carol Davila University of Medicine and Pharmacy, Bucharest, Romania. <sup>114</sup>Department of Statistics and Econometrics, Bucharest University of Economic Studies, Bucharest, Romania. <sup>115</sup>Department of Environmental and Occupational Health, University of Medical Sciences, Ondo, Nigeria. <sup>116</sup>Department of Microbiology, University of Medical Sciences, Ondo, Nigeria. <sup>117</sup>Regenerative Medicine, Organ Procurement and Transplantation Multi-disciplinary Center, Guilan University of Medical Sciences, Rasht, Iran. <sup>118</sup>Department of Surgery, Gadjah Mada University, Yogyakarta, Indonesia. <sup>119</sup>Rural Health Research Institute, Charles Sturt University, Orange, New South Wales, Australia. <sup>120</sup>Health Management and Economics Research Center, Iran University of Medical Sciences, Tehran, Iran. <sup>121</sup>College of Pharmacy, Al Ain University, Abu Dhabi, United Arab Emirates. <sup>122</sup>Department of Applied Mathematics, University of Washington, Seattle, WA, USA. <sup>123</sup>College of Art and Science, Ottawa University, Surprise, AZ, USA. <sup>124</sup>School of Life Sciences, Arizona State University, Tempe, AZ, USA. <sup>125</sup>College of Medicine and Health Sciences, Adigrat University, Adigrat, Ethiopia. <sup>126</sup>Department of Veterinary Pharmacology and Toxicology, University of Ilorin, Ilorin, Nigeria. <sup>127</sup>Department of Paediatrics, University of Malaya, Kuala Lumpur, Malaysia. <sup>128</sup>University of Malaya Medical Centre, University of Malaya, Kuala Lumpur, Malaysia. <sup>129</sup>Department of Medical Laboratory

Sciences, Haramaya University, Harar, Ethiopia. <sup>130</sup>Directorate of Research and Innovation Foresight, National Research and Innovation Agency (BRIN), Jakarta Pusat, Indonesia. <sup>131</sup>Honorary Council of Nutritionist Ethics, Persatuan Ahli Gizi Indonesia (Indonesian Nutrition Association), Jakarta, Indonesia. <sup>132</sup>Department of Public Health, Jimma University, Jimma, Ethiopia. <sup>133</sup>Department of Public Health, Wachemo University, Hossana, Ethiopia. <sup>134</sup>Department of Veterinary Pathology, University of Ilorin, Kwara, Nigeria. <sup>135</sup>Department of Immunology, Zanjan University of Medical Sciences, Zanjan, Iran. <sup>136</sup>Faculty of Nursing, Philadelpia University, Amman, Jordan. <sup>137</sup>Department of Forensic Medicine, Lumbini Medical College, Palpa, Nepal. <sup>138</sup>Department of Midwifery, Bahir Dar University, Bahir Dar, Ethiopia. <sup>139</sup>Department of Epidemiology and Preventive Medicine, Monash University, Melbourne, Victoria, Australia. <sup>140</sup>The Judith Lumley Centre, La Trobe University, Melbourne, Victoria, Australia. <sup>141</sup>Universidad de San Martín de Porres, Lima, Peru. <sup>142</sup>The World Bank, Washington, DC, USA. <sup>143</sup>ASIDE Healthcare, Lewes, DE, USA. <sup>144</sup>Faculty of Medicine, October 6 University, 6th of October City, Egypt. <sup>145</sup>Department of Population Medicine, Qatar University, Doha, Qatar. <sup>146</sup>Department of Nursing, Saveh University of Medical Sciences, Saveh, Iran. <sup>147</sup>Health Research Institute, University of Canberra, Canberra, Australian Capital Territory, Australia. <sup>148</sup>School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran. <sup>149</sup>International Medical School, Management and Science University, Alam, Malaysia. <sup>150</sup>Department of Forensic Medicine and Toxicology, Manipal Academy of Higher Education, Manipal, India. <sup>151</sup>Division of Biological Sciences, Tamil Nadu State Council for Science and Technology, Chennai, India. <sup>152</sup>Department of Non-communicable Diseases, Bangladesh University of Health Sciences, Dhaka, Bangladesh. <sup>153</sup>Department of Forensic Science, Government Institute of Forensic Science, Nagpur, India. <sup>154</sup>Department of Veterinary Physiology and Biochemistry, University of Ilorin, Ilorin, Nigeria. <sup>155</sup>Non-communicable Diseases Research Center, Tehran University of Medical Sciences, Tehran, Iran. <sup>156</sup>School of Medicine, Iran University of Medical Sciences, Tehran, Iran. <sup>157</sup>Department of Medical Education, University of Nevada Las Vegas, Las Vegas, NV, USA. <sup>158</sup>Center of Research Excellence in Stillbirth, The University of Queensland, Brisbane, Queensland, Australia. <sup>159</sup>Health System and Population Studies Division, International Centre for Diarrhoeal Disease Research, Dhaka, Bangladesh. <sup>160</sup>Infectious Disease Research Department, King Abdullah International Medical Research Center, Riyadh, Saudi Arabia. <sup>161</sup>Department of Veterinary Microbiology, Usmanu Danfodiyo University, Sokoto, Nigeria. <sup>162</sup>Department of Biological Sciences, University of Porto, Porto, Portugal. <sup>163</sup>Research Unit on Applied Molecular Biosciences (UCIBIO), University of Porto, Porto, Portugal. <sup>164</sup>Department of Internal Medicine, University of São Paulo, São Paulo, Brazil. <sup>165</sup>Department of Public Health, Madda Walabu University, Addis Ababa, Ethiopia. <sup>166</sup>Department of Nutrition and Dietetics, Jimma University, Addis Ababa, Ethiopia. <sup>167</sup>Department of Nursing, Bahir Dar University, Bahir Dar, Ethiopia. <sup>168</sup>Department of Forensic Chemistry, Government Institute of Forensic Science, Aurangabad, India. <sup>169</sup>Institute of Applied Health Research, University of Nottingham, Nottingham, UK. <sup>170</sup>Institute of Applied Health Research, University of Birmingham, Birmingham, UK. <sup>171</sup>Department of Anatomy, All India Institute of Medical Sciences, Jodhpur, India. <sup>172</sup>Department of Community Medicine and Family Medicine, All India Institute of Medical Sciences, Jodhpur, India. <sup>173</sup>School of Public Health, All India Institute of Medical Sciences, Jodhpur, India. <sup>174</sup>Division of Hematology Oncology, University of Massachusetts Medical School, Springfield, MA, USA. <sup>175</sup>Global Health Neurology Lab, NSW Brain Clot Bank, Sydney, New South Wales, Australia. <sup>176</sup>Division of Cerebrovascular Medicine and Neurology, National Cerebral and Cardiovascular Center, Suita, Japan. <sup>177</sup>Department of General Medicine, Manipal Academy of Higher Education, Mangalore, India. <sup>178</sup>Department of Human Genetics and Molecular Medicine, Central University of Punjab, Bathinda, India. <sup>179</sup>Department of Community Medicine and Family Medicine, All India Institute of Medical Sciences, Deoghar, India. <sup>180</sup>Department of Community & Family Medicine, All India Institute of Medical Sciences, Ramanathapuram, India. <sup>181</sup>Department of Computer Science, East Carolina University, Greenville, NC, USA. <sup>182</sup>Department of Internal Medicine, Manipal Academy of Higher Education, Mangalore, India. <sup>183</sup>ICMR-Regional Medical Research Centre, North East Region, Indian Council of Medical Research, Dibrugarh, India. <sup>184</sup>Internal Medicine Department, Shahid Beheshti University of Medical Sciences, Tehran, Iran. <sup>185</sup>Department of Medicine, University Ferhat Abbas of Setif, Sétif, Algeria. <sup>186</sup>Department of Epidemiology and Preventive Medicine, University Hospital Saadna Abdenour, Sétif, Algeria. <sup>187</sup>School of Pharmacy, The University of Jordan, Amman, Jordan. <sup>188</sup>Department of Basic Biomedical Sciences, University of Sharjah, Sharjah, United Arab Emirates. <sup>189</sup>Faculty of Health Sciences Healthcare Management Department, Ankara University, Ankara, Turkey. <sup>190</sup>Department of Clinical Pharmacy, University of Medicine and Pharmacy of Craiova, Craiova, Romania. <sup>191</sup>Department of Internal and Geriatric Medicine, Hospital Italiano de Buenos Aires (Italian Hospital of Buenos Aires), Buenos Aires, Argentina. <sup>192</sup>Board of Directors, Argentine Society of Medicine, Buenos Aires, Argentina. <sup>193</sup>Department of Ophthalmology, Beijing Institute of Ophthalmology, Beijing, China. <sup>194</sup>Graduate Program in Nutrition and Health, Federal University of Espírito Santo, Vitória, Brazil. <sup>195</sup>Department of Medical, Surgical, and Health Sciences, University of Trieste, Trieste, Italy. <sup>196</sup>Public Health Unit, University Health Agency Giuliano-Isontina (ASUGI), Trieste, Italy. <sup>197</sup>Department of Nutrition, Federal University of Santa Catarina, Florianópolis, Brazil. <sup>198</sup>College of Public Health, Medical, and Veterinary Sciences, James Cook University, Townsville, Queensland, Australia. <sup>199</sup>Department of Public Health, University of Mataram, Mataram, Indonesia. <sup>200</sup>Department of Biotechnology, Adamas University, Kolkata, India. <sup>201</sup>Institute for Skeletal Aging & Orthopedic Surgery, Hallym University, Chuncheon, South Korea. <sup>202</sup>State Disease Investigation Laboratory, Animal Resources Development Department, Agartala, India. <sup>203</sup>Clinical Nutrition Department, Jazan University, Jazan, Saudi Arabia. <sup>204</sup>Temerty Faculty of Medicine, University of Toronto, Toronto, Ontario, Canada. <sup>205</sup>Department of Community Medicine, Datta Meghe Institute of Medical Sciences, Sawangi, India. <sup>206</sup>Department of Oral Medicine and Radiology, King George's Medical University, Lucknow, India. <sup>207</sup>Clinical Research Center, Zhujiang Hospital of Southern Medical University, Guangzhou, China. <sup>208</sup>Division of Cardiovascular Medicine, Harvard University, Boston, MA, USA. <sup>209</sup>Division of Infectious Diseases, Virginia Commonwealth University, Richmond, VA, USA. <sup>210</sup>Department of Public Health, Administration, and Social Sciences, Cayetano Heredia University, Lima, Peru. <sup>211</sup>Department of Clinical Oncology, Queen Elizabeth Hospital, Hong Kong, China. <sup>212</sup>Department of Medicine, National University of Singapore, Singapore, Singapore. <sup>213</sup>Centre for Research Impact & Outcome, Chitkara University, Rajpura, India. <sup>214</sup>Department of Community Medicine, Jawaharlal Nehru Medical College, Wardha, India. <sup>215</sup>The Interdisciplinary Research Group on Biomedicine and Health, VNU International School (VNUIS), Hanoi, Viet Nam. <sup>216</sup>Faculty of Applied Sciences, VNU International School (VNUIS), Hanoi, Viet Nam. <sup>217</sup>Department of Paediatric Surgery, Federal Medical Centre, Umuahia, Nigeria. <sup>218</sup>Clinical Epidemiology and Public Health Research Unit, Burlo Garofolo Institute for Maternal and Child Health, Trieste, Italy. <sup>219</sup>Department of Family Medicine and Public Health, University of California, San Diego, La Jolla, CA, USA. <sup>220</sup>Department of Diagnostic and Therapeutic Technologies, Cooperativa de Ensino Superior Politécnico e Universitário (Polytechnic and University Higher Education Cooperative), Vila Nova de Famalicão, Portugal. <sup>221</sup>Institute for Research and Innovation in Health (i3S), University of Porto, Porto, Portugal. <sup>222</sup>School of Nursing, Federal University of Minas Gerais, Belo Horizonte, Brazil. <sup>223</sup>Research Center for Child Psychiatry, University of Turku, Turku, Finland. <sup>224</sup>Iranian Research Center for HIV/AIDS (IRCHA), Tehran University of Medical Sciences, Tehran, Iran. <sup>225</sup>Department of Medical and Surgical Sciences and Advanced Technologies 'GF Ingrassia', University of Catania, Catania, Italy. <sup>226</sup>Department of Public Health, Haramaya University, Harar, Ethiopia. <sup>227</sup>Department of Biochemistry, Ministry of Health and Welfare, New Delhi, India. <sup>228</sup>Clinical Sciences Department, University of Sharjah, Sharjah, United Arab Emirates. <sup>229</sup>Medical College, Albany Medical College, Albany, NY, USA. <sup>230</sup>Department of Food, Environmental and Nutritional Sciences (DeFENS), Università degli Studi di Milano (University of Milan), Milan, Italy. <sup>231</sup>School of Medicine, University of Colima, Colima, Mexico. <sup>232</sup>Department of Research, State Cancerology Institute of Colima, IMSSBIENESTAR, Colima, Mexico. <sup>233</sup>Department of Research, Gujarat Adani Institute of Medical Sciences, Bhuj, India. <sup>234</sup>Department of Public Health, Madda Walabu University, Goba, Ethiopia. <sup>235</sup>JSS Medical College Department of Biochemistry, Jagadguru Sri Shivarathreeswara University, Mysuru, India.

<sup>236</sup>Research and Development Cell, Dr. D. Y. Patil Vidyapeeth, Pune, India. <sup>237</sup>University of South Carolina, Columbia, SC, USA. <sup>238</sup>Department of Medicine, Pham Ngoc Thach University of Medicine, Ho Chi Minh City, Viet Nam. <sup>239</sup>Department of Medicine, Can Tho University of Medicine and Pharmacy, Can Tho, Viet Nam. <sup>240</sup>School of Health, Guilan University of Medical Sciences, Rasht, Iran. <sup>241</sup>Department of Community Nutrition, Shahid Beheshti University of Medical Sciences, Tehran, Iran. <sup>242</sup>Department of Public Health, Jazan University, Jazan, Saudi Arabia. <sup>243</sup>Cardio-Thoraco-Vascular Department, Azienda Sanitaria Universitaria Giuliano Isontina, Trieste, Italy. <sup>244</sup>Independent Consultant, South Plainfield, NJ, USA. <sup>245</sup>Department of Medicine, Bangalore Medical College and Research Institute, Bangalore, India. <sup>246</sup>Manipal Academy of Higher Education, Manipal, India. <sup>247</sup>Department of Forensic Medicine and Toxicology, Kasturba Medical College, Mangalore, India. <sup>248</sup>Faculty of Science and Humanities, SRM Institute of Science and Technology, Kattankulathur, India. <sup>249</sup>College of Medicine, Ajman University, Ajman, United Arab Emirates. <sup>250</sup>School of Health Sciences, Universiti Sains Malaysia (University of Science Malaysia), Kubang Kerian, Malaysia. <sup>251</sup>Advanced Nursing Department, Universitas Airlangga (Airlangga University), Surabaya, Indonesia. <sup>252</sup>Department of Biochemistry, Ege University, Izmir, Turkey. <sup>253</sup>Azerbaijan State University of Economics (UNEC), Baku, Azerbaijan. <sup>254</sup>Iran University of Medical Sciences, Tehran, Iran. <sup>255</sup>Faculty of Science and Health, University of Portsmouth, Hampshire, UK. <sup>256</sup>Department of Public Health and Community Medicine, Tanta University, Tanta city, Egypt. <sup>257</sup>School of Public Health, Texila American University, Guyana, Guyana. <sup>258</sup>Institute of Public Health, United Arab Emirates University, Al Ain, United Arab Emirates. <sup>259</sup>Department of Internal Medicine, Ain Shams University, Cairo, Egypt. <sup>260</sup>Section of Adult Hematology, King Saud University, Riyadh, Saudi Arabia. <sup>261</sup>Faculty of Medicine, University of Tripoli, Tripoli, Libya. <sup>262</sup>Houston Methodist Hospital, Houston, TX, USA. <sup>263</sup>Department of Basic Medical Sciences, University of Sharjah, Sharjah, United Arab Emirates. <sup>264</sup>Department of Clinical Pathology, Mansoura University, Mansoura, Egypt. <sup>265</sup>Department of Infectious Diseases and Public Health, City University of Hong Kong, Hong Kong, China. <sup>266</sup>Department of Animal Medicine, Zagazig University, Zagazig, Egypt. <sup>267</sup>Department of Public Health and Tropical Medicine, James Cook University, Townsville, Queensland, Australia. <sup>268</sup>Department of Epidemiology and Medical Statistics, University of Ibadan, Ibadan, Nigeria. <sup>269</sup>Research Centre for Healthcare and Community, Coventry University, Coventry, UK. <sup>270</sup>Department of Oral Biology, Riphah International University, Islamabad, Pakistan. <sup>271</sup>Director of the Scientific and Technological Park, Kazakh National Medical University, Almaty, Kazakhstan. <sup>272</sup>Department of Medicine, Korea University, Seoul, South Korea. <sup>273</sup>Department of Clinical Nutrition and Dietetics, Applied Science Private University, Amman, Jordan. <sup>274</sup>Centre for Public Health, Equity and Human Flourishing, Torrens University Australia, Adelaide, South Australia, Australia. <sup>275</sup>Institute of Resource Governance and Social Change, Kupang, Indonesia. <sup>276</sup>Department of Biology and Medicine, Brown University, Providence, RI, USA. <sup>277</sup>Department of Social Medicine and Epidemiology, Guilan University of Medical Sciences, Rasht, Iran. <sup>278</sup>Department of Pharmacy, Wollega University, Nekemte, Ethiopia. <sup>279</sup>Department of Social Sciences, University of Nicosia, Nicosia, Cyprus. <sup>280</sup>Institute of Public Health, Charité Universitätsmedizin Berlin (Charité Medical University Berlin), Berlin, Germany. <sup>281</sup>Department of General Surgery, Carol Davila University of Medicine and Pharmacy, Bucharest, Romania. <sup>282</sup>Department of Child Dental Health, Obafemi Awolowo University, Ife, Nigeria. <sup>283</sup>Clinical Science Department, Nigerian Institute of Medical Research, Lagos, Nigeria. <sup>284</sup>Department of Cell Biology and Biotechnology, K.A. Timiryazev Institute of Plant Physiology, Moscow, Russia. <sup>285</sup>Department of Dermatology, Kobe University, Kobe, Japan. <sup>286</sup>Department of Pathology, Federal University of Espirito Santo, Vitória, Brazil. <sup>287</sup>Department of Community Medicine and Family Medicine, All India Institute of Medical Sciences, Gorakhpur, India. <sup>288</sup>Department of Community Medicine, Bayero University Kano, Kano, Nigeria. <sup>289</sup>Department of Community Medicine, Aminu Kano Teaching Hospital, Kano, Nigeria. <sup>290</sup>Department of Community Medicine, Datta Meghe Institute of Medical Sciences, Wardha, India. <sup>291</sup>Department of Food Technology, Salahaddin University-Erbil, Erbil, Iraq. <sup>292</sup>Department of Nutrition and Dietetics, Cihan University-Erbil, Erbil, Iraq. <sup>293</sup>Department of Community Medicine and Family Medicine, All India Institute of Medical Sciences, Nagpur, India. <sup>294</sup>Department of Pharmacology, Indore Institute of Pharmacy, Indore, India. <sup>295</sup>Department of Midwifery, Adigrat University, Adigrat, Ethiopia. <sup>296</sup>Department of Environmental Health, Wollo University, Dessie, Ethiopia. <sup>297</sup>Department of Public Health Nutrition, Aksum University, Mekelle, Ethiopia. <sup>298</sup>College of Health Science, Oda Bultum University, Chiro, Ethiopia. <sup>299</sup>Department of Public Health, Debre Berhan University, Debre Berhan, Ethiopia. <sup>300</sup>Department of Public Health, Menelik II Medical and Health Science College, Addis Ababa, Ethiopia. <sup>301</sup>Department of Medical Laboratory Science, Addis Ababa University, Addis Ababa, Ethiopia. <sup>302</sup>School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran. <sup>303</sup>Center of Health Management, Aden University, Aden, Yemen. <sup>304</sup>Tropical Health Department, Alexandria University, Alexandria, Egypt. <sup>305</sup>Family and Community Medicine Department, King Khalid University, Abha, Saudi Arabia. <sup>306</sup>Cancer Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran. <sup>307</sup>Department of Radiology, University of Southern California, Los Angeles, CA, USA. <sup>308</sup>Country Office, World Health Organization (WHO), Astana, Kazakhstan. <sup>309</sup>Department of Zoology, KKS Women's College, Balasore, India. <sup>310</sup>Department of Nursing, Aksum University, Aksum, Ethiopia. <sup>311</sup>Department of Epidemiology, Universidade de São Paulo (University of São Paulo), São Paulo, Brazil. <sup>312</sup>Department of Clinical Science, University Of Sulaimani, Sulaimani, Iraq. <sup>313</sup>Department of Community Medicine, University of Peradeniya, Kandy, Sri Lanka. <sup>314</sup>Department of Nephrology, Max Super Speciality Hospital, New Delhi, India. <sup>315</sup>Non-communicable Diseases Division (NCD), Indian Council of Medical Research, New Delhi, India. <sup>316</sup>Department of Internal Medicine, Independent Consultant, Bharatpur, India. <sup>317</sup>Independent Consultant, Delhi, India. <sup>318</sup>Department of Epidemiology and Biostatistics, University of South Carolina, Columbia, SC, USA. <sup>319</sup>Centre for Noncommunicable Diseases and Nutrition, BRAC University, Dhaka, Bangladesh. <sup>320</sup>Department of Toxicology, Shriram Institute for Industrial Research, Delhi, India. <sup>321</sup>Faculty of Medicine Health and Human Sciences, Macquarie University, Sydney, New South Wales, Australia. <sup>322</sup>Global Virus Network, Middle East Region, Shiraz, Iran. <sup>323</sup>Department of Medicine, University of Pittsburgh Medical Center, Pittsburgh, PA, USA. <sup>324</sup>Department of Clinical Pharmacology and Medicine, University of Kufa, Najaf, Iraq. <sup>325</sup>Natural and Medical Sciences Research Center, University of Nizwa, Nizwa, Oman. <sup>326</sup>Biochemistry Department, Ain Shams University, Cairo, Egypt. <sup>327</sup>School of Health and Environmental Studies, Hamdan Bin Mohammed Smart University, Dubai, United Arab Emirates. <sup>328</sup>Department of Epidemiology, University of Michigan School of Public Health, Ann Arbor, MI, USA. <sup>329</sup>Department of Zoology and Entomology, Al-Azhar University, Cairo, Egypt. <sup>330</sup>Department of Health Research Methods, Evidence, and Impact, McMaster University, Hamilton, Ontario, Canada. <sup>331</sup>Department of Biochemistry and Molecular Biology, Tejgaon College, Dhaka, Bangladesh. <sup>332</sup>Department of Ophthalmology, Iran University of Medical Sciences, Tehran, Iran. <sup>333</sup>Department of Pharmacy, Marwadi University, Rajkot, India. <sup>334</sup>Public Health Department, Dalhatu Araf Specialist Hospital, Lafia, Nigeria. <sup>335</sup>Department of Public Health, Federal University of Lafia, Lafia, Nigeria. <sup>336</sup>School of Nursing and Midwifery, Tabriz University of Medical Sciences, Tabriz, Iran. <sup>337</sup>Independent Consultant, Tabriz, Iran. <sup>338</sup>Institute of Pharmaceutical Sciences, University of Veterinary and Animal Sciences, Lahore, Pakistan. <sup>339</sup>Department of Pharmacy Administration and Clinical Pharmacy, Xian Jiaotong University, Xian, China. <sup>340</sup>Independent Consultant, Santa Clara, CA, USA. <sup>341</sup>Department of Medicine, MedStar Health, Washington, DC, USA. <sup>342</sup>Department of Medicine, Georgetown University, Washington, DC, USA. <sup>343</sup>Department of Microbiology, Taiz University, Taiz, Yemen. <sup>344</sup>School of Medicine, Nankai University, Tianjin, China. <sup>345</sup>Graduate School of Medicine, University of Tokyo, Tokyo, Japan. <sup>346</sup>School of Dentistry, Hanoi Medical University, Hanoi, Viet Nam. <sup>347</sup>Kasturba Medical College Mangalore, Manipal Academy of Higher Education, Manipal, India. <sup>348</sup>Department of Decision and Information Sciences, University of Houston, Houston, TX, USA. <sup>349</sup>Public Health Research Group, Nature Study Society of Bangladesh, Khulna, Bangladesh. <sup>350</sup>School of Computer Science, Duy Tan University, Da Nang, Viet Nam. <sup>351</sup>Jadara University Research Center, Jadara University, Irbid, Jordan. <sup>352</sup>Department of Internal Medicine, Carol Davila University of Medicine and Pharmacy, Bucharest, Romania. <sup>353</sup>Department of Legal Medicine and

Bioethics, Carol Davila University of Medicine and Pharmacy, Bucharest, Romania. <sup>354</sup>Department of Clinical Legal Medicine, National Institute of Legal Medicine Mina Minovici, Bucharest, Romania. <sup>355</sup>Faculty of Medicine, The Chinese University of Hong Kong, Hong Kong, China. <sup>356</sup>Department of Epidemiology, Johns Hopkins University, Baltimore, MD, USA. <sup>357</sup>Department of Biological Sciences and Chemistry, University of Nizwa, Nizwa, Oman. <sup>358</sup>International Master Program for Translational Science, Taipei Medical University, Taipei, Taiwan. <sup>359</sup>Department of Occupational Safety and Health, China Medical University, Taichung, Taiwan. <sup>360</sup>Department of Occupational Therapy, Asia University, Taichung, Taiwan. <sup>361</sup>Department of Health Promotion and Education, University of Ibadan, Ibadan, Nigeria. <sup>362</sup>Center for Nutritional Epidemiology and Policy Research, National Institutes of Biomedical Innovation, Health and Nutrition, Osaka, Japan. <sup>363</sup>West Africa RCC, Africa Centre for Disease Control and Prevention, Abuja, Nigeria. <sup>364</sup>Department of Community Medicine, University College Hospital, Ibadan, Nigeria. <sup>365</sup>Faculty of Medicine, University of Belgrade, Belgrade, Serbia. <sup>366</sup>Faculty of Medical Sciences, University of Kragujevac, Kragujevac, Serbia. <sup>367</sup>Institute of Health Research, University of Health and Allied Sciences, Ho, Ghana. <sup>368</sup>Department of Health Research, ICMR National Institute for Research in Tuberculosis, Chennai, India. <sup>369</sup>Faculty of Health and Life Sciences, University of Exeter, Exeter, UK. <sup>370</sup>Faculty of Public Health, Universitas Muhammadiyah Aceh, Banda Aceh, Indonesia. <sup>371</sup>Faculty of Pharmacy, Universitas Ahmad Dahlan, Yogyakarta, Indonesia. <sup>372</sup>School of Pharmacy, BRAC University, Dhaka, Bangladesh. <sup>373</sup>Institute for Physical Activity and Nutrition, Deakin University, Burwood, Victoria, Australia. <sup>374</sup>Department of Clinical Pharmacy & Pharmacy Practice, Asian Institute of Medicine, Science and Technology, Bedong, Malaysia. <sup>375</sup>Malaysian Academy of Pharmacy, Puchong, Malaysia. <sup>376</sup>Department of General Surgery and Medical-Surgical Specialties, University of Catania, Catania, Italy. <sup>377</sup>School of Health Systems and Public Health, University of Washington, Seattle, WA, USA. <sup>378</sup>Department of Microbiology, Central University of Punjab, Bathinda, India. <sup>379</sup>Department of Immunology, Kerman University of Medical Sciences, Kerman, Iran. <sup>380</sup>Department of Immunology, Rafsanjan University of Medical Sciences, Rafsanjan, Iran. <sup>381</sup>Department of Immunology, Shahid Beheshti University of Medical Sciences, Tehran, Iran. <sup>382</sup>Department of Leukemia, The University of MD Anderson Cancer Center, Houston, TX, USA. <sup>383</sup>Department of Health and Safety, Dubai Municipality, Dubai, United Arab Emirates. <sup>384</sup>The World Academy of Sciences UNESCO, Trieste, Italy. <sup>385</sup>Shaanxi University of Technology, Hanzhong, China. <sup>386</sup>Department of Environmental Engineering, Islamic Azad University, Ahvaz, Iran. <sup>387</sup>Department of Public Health, Daffodil International University, Dhaka, Bangladesh. <sup>388</sup>Department of Pharmacology, Imam Mohammad Ibn Saud Islamic University, Riyadh, Saudi Arabia. <sup>389</sup>Department of Public Health, M S Ramaiah University of Applied Sciences, Bangalore, India. <sup>390</sup>Institute for Global Public Health, University of Manitoba, Winnipeg, Manitoba, Canada. <sup>391</sup>Department of Biochemistry, Government Medical College, Mysuru, India. <sup>392</sup>Department of Epidemiology, Wachemo University, Hossana, Ethiopia. <sup>393</sup>Department of Internal Medicine, GCS Medical College, Hospital & Research Centre, Ahmedabad, India. <sup>394</sup>Faculty of Veterinary Medicine, University of Calgary, Calgary, Alberta, Canada. <sup>395</sup>Young Researchers and Elite Club, Islamic Azad University, Karaj, Iran. <sup>396</sup>Rothschild Foundation Hospital, Institut Français de Myopie, Paris, France. <sup>397</sup>Singapore Eye Research Institute, Singapore, Singapore. <sup>398</sup>Department of Community Medicine, Jubilee Mission Medical College & Research Institute, Thrissur, India. <sup>399</sup>Department of Community Medicine, Manipal Academy of Higher Education, Mangalore, India. <sup>400</sup>Department of Economics, National Open University, Benin City, Nigeria. <sup>401</sup>Gastrointestinal and Liver Diseases Research Center, Guilan University of Medical Sciences, Rasht, Iran. <sup>402</sup>Caspian Digestive Disease Research Center, Guilan University of Medical Sciences, Rasht, Iran. <sup>403</sup>Department of Family Medicine and Public Health, University of Opole, Opole, Poland. <sup>404</sup>Minimally Invasive Surgery Research Center, Iran University of Medical Sciences, Tehran, Iran. <sup>405</sup>School of Public Health, University College Cork, Cork, Ireland. <sup>406</sup>Department of Statistics, Salahaddin University-Erbil, Erbil, Iraq. <sup>407</sup>Department of Business Administrations, Cihan University-Erbil, Erbil, Iraq. <sup>408</sup>Department of Internal Medicine, Wayne State University, Detroit, MI, USA. <sup>409</sup>School of Management and Medical Informatics, Tabriz University of Medical Sciences, Tabriz, Iran. <sup>410</sup>Russell H. Morgan Department of Radiology and Radiological Science, Johns Hopkins University, Baltimore, MD, USA. <sup>411</sup>Faculty of Dentistry, University of Puthisastra, Phnom Penh, Cambodia. <sup>412</sup>Office of the Executive Director, Cephas Health Research Initiative Inc, Ibadan, Nigeria. <sup>413</sup>The Hansjörg Wyss Department of Plastic and Reconstructive Surgery, NYU Langone Health, New York, NY, USA. <sup>414</sup>Cleft Lip and Palate Surgery Division, Global Smile Foundation, Norwood, MA, USA. <sup>415</sup>Laboratory Science Department, Khomein University of Medical Sciences, Khomein, Iran. <sup>416</sup>Department of Immunology, Tehran University of Medical Sciences, Tehran, Iran. <sup>417</sup>Endocrine Research Center, Iran University of Medical Sciences, Tehran, Iran. <sup>418</sup>Department of Echocardiography, Iran University of Medical Sciences, Tehran, Iran. <sup>419</sup>Surgery Research Unit, University of Oulu, Oulu, Finland. <sup>420</sup>Department of Molecular Medicine and Surgery, Karolinska Institute, Stockholm, Sweden. <sup>421</sup>Department of ENT, Dr. B. R. Ambedkar State Institute of Medical Sciences (AIMS), Mohali, India. <sup>422</sup>International Research Center of Excellence, Institute of Human Virology Nigeria, Abuja, Nigeria. <sup>423</sup>Julius Centre for Health Sciences and Primary Care, Utrecht University, Utrecht, The Netherlands. <sup>424</sup>Department of Biomedical Science, Arba Minch University, Arbaminch, Ethiopia. <sup>425</sup>School of Nursing and Midwifery, Haramaya University, Harar, Ethiopia. <sup>426</sup>Department of Healthcare Services Management, Alborz University of Medical Sciences, Karaj, Iran. <sup>427</sup>Open, Distance and eLearning Campus, University of Nairobi, Nairobi, Kenya. <sup>428</sup>Amity Institute of Forensic Sciences, Amity University, Noida, India. <sup>429</sup>School of Medicine, Tehran University of Medical Sciences, Tehran, Iran. <sup>430</sup>Endocrinology and Metabolism Research Institute, Non-Communicable Diseases Research Center (NCDRC), Tehran, Iran. <sup>431</sup>College of Health Sciences, Abu Dhabi University, Abu Dhabi, United Arab Emirates. <sup>432</sup>School of Food and Agricultural Sciences, University of Management and Technology, Lahore, Pakistan. <sup>433</sup>Department of Biostatistics, Mazandaran University of Medical Sciences, Sari, Iran. <sup>434</sup>Epidemiology Program, Jazan University, Jazan, Saudi Arabia. <sup>435</sup>Department of Health, Nepal Development Society, Chitwan, Nepal. <sup>436</sup>Department of Preventable Non Communicable Disease, Menzies School of Health Research, Alice Springs, Northern Territory, Australia. <sup>437</sup>Faculty of Nursing, Yarmouk University, Irbid, Jordan. <sup>438</sup>Department of Basic Medical Sciences, Yarmouk University, Irbid, Jordan. <sup>439</sup>Department of Public Health, Jordan University of Science and Technology, Irbid, Jordan. <sup>440</sup>Department of Biochemistry, Liaquat University Of Medical and Health Sciences, Jamshoro, Pakistan. <sup>441</sup>Department of Biochemistry, All India Institute of Medical Sciences, Jodhpur, India. <sup>442</sup>Department of Internal Medicine, Corewell Health East William Beaumont University Hospital, Royal Oak, MI, USA. <sup>443</sup>Department of Health Management and Economics, Qom University of Medical Sciences, Qom, Iran. <sup>444</sup>Department of Health Economics, Iran University of Medical Sciences, Tehran, Iran. <sup>445</sup>Faculty of Health Sciences, University of Muhammadiyah Prof. Dr. Hamka, Jakarta, Indonesia. <sup>446</sup>Program Division, SEAMEO Regional Center for Food and Nutrition, Jakarta, Indonesia. <sup>447</sup>School of Traditional Chinese Medicine, Xiamen University Malaysia, Sepang, Malaysia. <sup>448</sup>Millennium Prevention, Inc., Westwood, MA, USA. <sup>449</sup>School of Health Sciences, Kristiania University College, Oslo, Norway. <sup>450</sup>Department of International Health and Sustainable Development, Tulane University, New Orleans, LA, USA. <sup>451</sup>Department of Nursing and Health Promotion, Oslo Metropolitan University, Oslo, Norway. <sup>452</sup>Department of Public Health Dentistry, Krishna Vishwa Vidyapeeth, Karad, India. <sup>453</sup>Social Determinants of Health Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran. <sup>454</sup>Children's Medical Center, Tehran University of Medical Sciences, Tehran, Iran. <sup>455</sup>Social Determinants of Health Research Center, Saveh University of Medical Sciences, Saveh, Iran. <sup>456</sup>Independent Consultant, Jakarta, Indonesia. <sup>457</sup>Department of Anthropology, Panjab University, Chandigarh, India. <sup>458</sup>Department of Biochemistry, University of Hail, Hail, Saudi Arabia. <sup>459</sup>Research and Publication Activity Division, Kazakh National Medical University, Almaty, Kazakhstan. <sup>460</sup>Center of Medicine and Public Health, Asfendiyarov Kazakh National Medical University, Almaty, Kazakhstan. <sup>461</sup>Department of Community Medicine, Rajendra Institute of Medical Sciences, Ranchi, India. <sup>462</sup>Centre for Digital Transformation, Indian Institute of Management, Ahmedabad, India. <sup>463</sup>Centre for Studies in Economics and Planning, Central University of Gujarat, Gandhinagar, India.

<sup>464</sup>Public Health, School of Medicine and Dentistry, Griffith University, Gold Coast, Queensland, Australia. <sup>465</sup>Faculty of Health and Life Sciences, Coventry University, Coventry, UK. <sup>466</sup>Department of Medicine, McMaster University, Hamilton, Ontario, Canada. <sup>467</sup>Department of Environment and Public Health, University of Environment and Sustainable Development, Somanya, Ghana. <sup>468</sup>Integrated Department of Epidemiology, Health Policy, Preventive Medicine and Pediatrics, Foundation for People-centric Health Systems, New Delhi, India. <sup>469</sup>Centre for Health: The Specialty Practice, New Delhi, India. <sup>470</sup>Indian Council of Medical Research, New Delhi, India. <sup>471</sup>Unidad de Genética y Salud Pública, Instituto de Ciencias Médicas, Las Tablas, Panama. <sup>472</sup>Ministry of Health, Hospital Joaquín Pablo Franco Sayas, Las Tablas, Panama. <sup>473</sup>Centre for Family Welfare, University of Indonesia, Depok, Indonesia. <sup>474</sup>Department of Global Health and Health Security, Taipei Medical University, Taipei, Taiwan. <sup>475</sup>Faculty of Medicine, University of Medicine and Pharmacy at Ho Chi Minh City, Ho Chi Minh City, Viet Nam. <sup>476</sup>Department of Cardiovascular Research, Methodist Hospital, Merrillville, IN, USA. <sup>477</sup>University of Medicine and Pharmacy at Ho Chi Minh City, Ho Chi Minh City, Viet Nam. <sup>478</sup>Department of Medical Science, Ajou University School of Medicine, Suwon, South Korea. <sup>479</sup>Department of Precision Medicine, Sungkyunkwan University, Suwon-si, South Korea. <sup>480</sup>Department of Family Medicine, The University of Texas Medical Branch, Galveston, TX, USA. <sup>481</sup>Center for Dentistry and Oral Hygiene, University of Groningen, Groningen, The Netherlands. <sup>482</sup>Stomatological Hospital, Southern Medical University, Guangzhou, China. <sup>483</sup>Department of Health Promotion and Health Education, National Taiwan Normal University, Taipei, Taiwan. <sup>484</sup>International Centre for Future Health Systems, University of New South Wales, Sydney, New South Wales, Australia. <sup>485</sup>Lerner Research Institute, Cleveland Clinic, Cleveland, OH, USA. <sup>486</sup>Department of Quantitative Health Science, Case Western Reserve University, Cleveland, OH, USA. <sup>487</sup>Department of Community Medicine, Government Medical College, Bengaluru, India. <sup>488</sup>One Health Research Group, Universidad de Las Américas (University of the Americas), Quito, Ecuador. <sup>489</sup>Department of Epidemiology and Evidence-Based Medicine, I.M. Sechenov First Moscow State Medical University, Moscow, Russia. <sup>490</sup>Department of Health Economics, Syreon Research Romania, Targu Mures, Romania. <sup>491</sup>Department of Doctoral Studies, George Emil Palade University of Medicine, Pharmacy, Science and Technology of Targu Mures, Targu Mures, Romania. <sup>492</sup>Department of Medicine, University of São Paulo, São Paulo, Brazil. <sup>493</sup>Geospatial Health and Development Team, Child Health Analytics, Telethon Kids Institute, Perth, Western Australia, Australia. <sup>494</sup>Scientific Research and Surveillance Systems, Macha Research Trust, Choma, Zambia. <sup>495</sup>Department of Chemistry, Salahaddin University-Erbil, Erbil, Iraq. <sup>496</sup>Department of Medical Biochemical Analysis, Cihan University-Erbil, Erbil, Iraq. <sup>497</sup>Centre for Public Health and Wellbeing, University of the West of England, Bristol, UK. <sup>498</sup>Department of Pediatric Cardiology, Tehran University of Medical Sciences, Tehran, Iran. <sup>499</sup>Rabigh Faculty of Medicine, King Abdulaziz University, Jeddah, Saudi Arabia. <sup>500</sup>Department of Maternal-Child Nursing and Public Health, Federal University of Minas Gerais, Belo Horizonte, Brazil. <sup>501</sup>Smidt Heart Institute, Cedars-Sinai Medical Center, Los Angeles, CA, USA. <sup>502</sup>Department of Nutrition and Dietetics, University of Concepción, Concepción, Chile. <sup>503</sup>Centre for Healthy Living, University of Concepción, Concepción, Chile. <sup>504</sup>Faculty of Humanities and Health Sciences, Curtin University, Sarawak, Malaysia. <sup>505</sup>Jeffrey Cheah School of Medicine and Health Sciences, Monash University, Subang Jaya, Malaysia. <sup>506</sup>Department of Anatomy and Developmental Biology, Monash University, Clayton, Victoria, Australia. <sup>507</sup>Department of Anatomy, Genetics and Biomedical Informatics, University of Colombo, Colombo, Sri Lanka. <sup>508</sup>Department of Public Health and Community Medicine, Central University of Kerala, Kasaragod, India. <sup>509</sup>Department of Community Medicine, Geetanjali Medical College and Hospital, Udaipur, India. <sup>510</sup>Nuffield Department of Population Health, University of Oxford, London, UK. <sup>511</sup>SSCANS, Symbiosis International University, Pune, India. <sup>512</sup>Department of Epidemiology and Biostatistics, University of California, San Francisco, San Francisco, CA, USA. <sup>513</sup>Department of Public Health, Arba Minch University, Arba Minch, Ethiopia. <sup>514</sup>School of Nursing and Midwifery, Deakin University, Melbourne, Victoria, Australia. <sup>515</sup>Department of Medical Laboratory Sciences, Adigrat University, Adigrat, Ethiopia. <sup>516</sup>Department of Internal Medicine, University of Gondar, Gondar, Ethiopia. <sup>517</sup>Johns Hopkins University, Baltimore, MD, USA. <sup>518</sup>Department of Medical Oncology and Hematology, Kantonsspital St. Gallen, St. Gallen, Switzerland. <sup>519</sup>Universidad Nacional Mayor de San Marcos, Lima, Peru. <sup>520</sup>University Centre Varazdin, University North, Varazdin, Croatia. <sup>521</sup>Department of Pharmacology, University of Kelaniya, Ragama, Sri Lanka. <sup>522</sup>Clinical Medicine Department, Colombo North Teaching Hospital, Ragama, Sri Lanka. <sup>523</sup>Department of Paediatrics, University of Kelaniya, Ragama, Sri Lanka. <sup>524</sup>University Paediatrics Unit, Colombo North Teaching Hospital, Ragama, Sri Lanka. <sup>525</sup>Department of Propedeutics of Internal Diseases & Arterial Hypertension, Pomeranian Medical University, Szczecin, Poland. <sup>526</sup>National Cancer Registry, Maria Skłodowska-Curie National Research Institute of Oncology, Warsaw, Poland. <sup>527</sup>Department of Pathology, Maria Skłodowska-Curie National Research Institute of Oncology, Warsaw, Poland. <sup>528</sup>International Ph.D. Program in Medicine, Taipei Medical University, Taipei, Taiwan. <sup>529</sup>Research Center for Artificial Intelligence in Medicine, Taipei Medical University, Taipei, Taiwan. <sup>530</sup>Department of Public Health Dentistry, Saveetha Institute of Medical and Technical Sciences (SIMATS), Chennai, India. <sup>531</sup>Global Institute of Public Health, Ananthapuri Hospitals and Research Institute, Trivandrum, India. <sup>532</sup>Faculty of Nursing and Midwifery, Tabriz University of Medical Sciences, Tabriz, Iran. <sup>533</sup>National Data Management Center for Health, Ethiopian Public Health Institute, Addis Ababa, Ethiopia. <sup>534</sup>Department of Endocrinology & Metabolism, All India Institute of Medical Sciences, Bhopal, India. <sup>535</sup>College of Health Science, University of Hargeisa, Hargeisa, Somalia. <sup>536</sup>Institute of Health Science, Jimma University, Jimma, Ethiopia. <sup>537</sup>College of Applied and Natural Science, University of Hargeisa, Hargeisa, Somalia. <sup>538</sup>Molecular Biology Unit, Sirius Training and Research Centre, Khartoum, Sudan. <sup>539</sup>Bio-Statistical and Molecular Biology Department, Sirius Training and Research Centre, Khartoum, Sudan. <sup>540</sup>College of Medicine, University of Duhok, Duhok, Iraq. <sup>541</sup>Social Determinants of Health Research Center, Tabriz University of Medical Sciences, Tabriz, Iran. <sup>542</sup>Midwifery Department, Tabriz University of Medical Sciences, Tabriz, Iran. <sup>543</sup>Infectious Diseases Research Center, Golestan University of Medical Sciences, Gorgan, Iran. <sup>544</sup>Health Systems and Policy Research Unit, Ahmadu Bello University, Zaria, Nigeria. <sup>545</sup>Heidelberg Institute of Global Health (HIGH), Heidelberg University, Heidelberg, Germany. <sup>546</sup>Department of Biostatistics, Shiraz University of Medical Sciences, Shiraz, Iran. <sup>547</sup>AI & Cyber Futures Institute, Charles Sturt University, Bathurst, New South Wales, Australia. <sup>548</sup>The University of Queensland, Brisbane, Queensland, Australia. <sup>549</sup>Faculty of Medicine, Birjand University of Medical Sciences, Birjand, Iran. <sup>550</sup>Department of Clinical Biochemistry, Babol University of Medical Sciences, Babol, Iran. <sup>551</sup>Department of Clinical Biochemistry, Tarbiat Modares University, Tehran, Iran. <sup>552</sup>Faculty of Biotechnologies (BioTech), ITMO University, Saint Petersburg, Russia. <sup>553</sup>Halal Research Center of IRI, Iran Food and Drug Administration, Ministry of Health and Medical Education, Tehran, Iran. <sup>554</sup>Unit of Pharmacotherapy, Epidemiology and Economics, University of Groningen (Rijksuniversiteit Groningen), Groningen, The Netherlands. <sup>555</sup>Department of Epidemiology and Biostatistics, Wuhan University, Wuhan, China. <sup>556</sup>Knowledge Management Department, Prahlad Omkarwati Foundation (POF), Mumbai, India. <sup>557</sup>Independent Consultant, Changescape Consulting, New Delhi, India. <sup>558</sup>Department of Surgery, General University Hospital of Patras, Patras, Greece. <sup>559</sup>Faculty of Medicine, University of Thessaly, Larissa, Greece. <sup>560</sup>Amity Institute of Pharmacy, Amity University, Noida, India. <sup>561</sup>Department of Community and Global Health, The University of Tokyo, Tokyo, Japan. <sup>562</sup>Surgery Department, University of Medicine and Pharmacy "Grigore T Popa" Iasi, Iasi, Romania. <sup>563</sup>Second Surgical Unit, Regional Institute of Oncology, Iasi, Romania. <sup>564</sup>College of Medicine, Shaqra University, Riyadh, Saudi Arabia. <sup>565</sup>Department of Pediatrics & Pediatric Pulmonology, Institute of Mother & Child Care, Multan, Pakistan. <sup>566</sup>Department of Research Methods, Orthopaedic Research Group, Coimbatore, India. <sup>567</sup>Department of Biotechnology, Karpagam Academy of Higher Education (Deemed to be University), Coimbatore, India. <sup>568</sup>Department of Technology, The University of Lahore, Lahore, Pakistan. <sup>569</sup>Research Centre for Health Sciences (RCHS), The University of Lahore, Lahore, Pakistan. <sup>570</sup>Research and Analytics Department, Initiative for Financing Health and Human Development, Chennai, India.

<sup>571</sup>Department of Research and Analytics, Bioinsilico Technologies, Chennai, India. <sup>572</sup>Department of Computer Science, University of Illinois, Urbana, IL, USA. <sup>573</sup>Department Health Services Research, University of Alabama at Birmingham, Birmingham, AL, USA. <sup>574</sup>Heart Failure Research Center, Isfahan University of Medical Sciences, Isfahan, Iran. <sup>575</sup>Neuroscience Research Center, Isfahan University of Medical Sciences, Isfahan, Iran. <sup>576</sup>Department of Medical Laboratory Analysis, Cihan University Sulaimaniya, Sulaimaniya, Iraq. <sup>577</sup>DY Patil School of Public Health, Dr. D. Y. Patil University, Navi Mumbai, India. <sup>578</sup>Mysore Medical College and Research Institute, Government Medical College, Mysore, India. <sup>579</sup>Nursing & Midwifery Research Department (NMRD), Hamad Medical Corporation, Doha, Qatar. <sup>580</sup>Department of Dental Public Health, King Abdulaziz University, Jeddah, Saudi Arabia. <sup>581</sup>Department of Health Policy and Oral Epidemiology, Harvard University, Boston, MA, USA. <sup>582</sup>College of Medicine and Health Sciences, United Arab Emirates University, Al Ain, United Arab Emirates. <sup>583</sup>Department of Circulation and Medical Imaging, Norwegian University of Science and Technology, Trondheim, Norway. <sup>584</sup>Department of Biotechnology, University of Central Punjab, Lahore, Pakistan. <sup>585</sup>Department of Health Promotion, Zahedan University of Medical Sciences, Zahedan, Iran. <sup>586</sup>Department of Research, TroDDIVaT Initiative, Buea, Cameroon. <sup>587</sup>Department of Microbiology and Parasitology, University of Buea, Buea, Cameroon. <sup>588</sup>Medical Biochemistry, Mekelle University, Mekelle, Ethiopia. <sup>589</sup>Department of General Surgery, Emergency University Hospital Bucharest, Bucharest, Romania. <sup>590</sup>Department of Anatomy and Embryology, Carol Davila University of Medicine and Pharmacy, Bucharest, Romania. <sup>591</sup>Department of Cardiology, Cardio-Aid, Bucharest, Romania. <sup>592</sup>HIV/STI Surveillance Research Center, Kerman University of Medical Sciences, Kerman, Iran. <sup>593</sup>Department of Epidemiology, Non-Communicable Diseases Research Center (NCDRC), Tehran, Iran. <sup>594</sup>School of Nursing, University of Gondar, Gondar, Ethiopia. <sup>595</sup>Department of Public Health, University of Yaoundé I, Yaoundé, Cameroon. <sup>596</sup>Department of Biological Sciences, University of Embu, Embu, Kenya. <sup>597</sup>Cardiovascular Laboratory, Methodist Hospital, Merrillville, IN, USA. <sup>598</sup>Department of Allergy, Immunology and Dermatology, Hanoi Medical University, Hanoi, Viet Nam. <sup>599</sup>Faculty of Medicine, Duy Tan University, Da Nang, Viet Nam. <sup>600</sup>Institute for Research and Training in Medicine, Biology and Pharmacy, Duy Tan University, Da Nang, Viet Nam. <sup>601</sup>Radiology Department, University of California, Los Angeles, Los Angeles, CA, USA. <sup>602</sup>Department of General Medicine, Pham Ngoc Thach University of Medicine, Ho Chi Minh City, Viet Nam. <sup>603</sup>Faculty of Medicine, Pham Ngoc Thach University of Medicine, Ho Chi Minh City, Viet Nam. <sup>604</sup>Department of Surgery, Danang Family Hospital, Danang, Viet Nam. <sup>605</sup>Tuberculosis Group, Oxford University Clinical Research Unit, Vietnam, Ho Chi Minh City, Viet Nam. <sup>606</sup>Department of General Medicine, University of Medicine and Pharmacy at Ho Chi Minh City, Ho Chi Minh City, Viet Nam. <sup>607</sup>International Islamic University Islamabad, Islamabad, Pakistan. <sup>608</sup>School of Chemical & Biomolecular Engineering, University of Sydney, Sydney, New South Wales, Australia. <sup>609</sup>Faculty of Applied Sciences, Taiz University, Taiz, Yemen. <sup>610</sup>Global Research Institute, Keio University, Tokyo, Japan. <sup>611</sup>Department of Global Health Policy, University of Tokyo, Tokyo, Japan. <sup>612</sup>Department of Statistics, Shahjalal University of Science and Technology, Sylhet, Bangladesh. <sup>613</sup>School of Medicine, Xiamen University, Xiamen, China. <sup>614</sup>Health Policy Research Center, Shiraz University of Medical Sciences, Shiraz, Iran. <sup>615</sup>Health Research Institute, Babol University of Medical Sciences, Babol, Iran. <sup>616</sup>School of Health, Bam University of Medical Sciences, Bam, Iran. <sup>617</sup>Department of Paediatrics, Nnamdi Azikiwe University, Awka, Nigeria. <sup>618</sup>Department of Public Health, Banten School of Health Science, South Tangerang, Indonesia. <sup>619</sup>Ministry of Research, Technology and Higher Education, Higher Education Service Institutions (LL-DIKTI) Region IV, Bandung, Indonesia. <sup>620</sup>Center of Excellence in Reproductive Health Innovation (CERHI), University of Benin, Benin City, Nigeria. <sup>621</sup>Department of Physiology, University of Benin, Edo, Nigeria. <sup>622</sup>Department of Physiology, Benson Idahosa University, Benin City, Nigeria. <sup>623</sup>Sheffield Centre for Health and Related Research, University of Sheffield, Sheffield, UK. <sup>624</sup>Department of Gynecology and Obstetrics, Emory University, Atlanta, GA, USA. <sup>625</sup>University of Sydney, Sydney, New South Wales, Australia. <sup>626</sup>Department of Food and Nutrition, Seoul National University, Seoul, South Korea. <sup>627</sup>College of Medicine, University of Ibadan, Ibadan, Nigeria. <sup>628</sup>School of Pharmacy, University of the Western Cape, Cape Town, South Africa. <sup>629</sup>Department of Psychiatry and Behavioural Neurosciences, McMaster University, Hamilton, Ontario, Canada. <sup>630</sup>Department of Psychiatry, University of Lagos, Lagos, Nigeria. <sup>631</sup>Department of Nursing Science, Bowen University, Iwo, Nigeria. <sup>632</sup>Diplomacy and Public Relations Department, University of Human Development, Sulaymaniyah, Iraq. <sup>633</sup>Department of Pharmacotherapy and Pharmaceutical Care, Medical University of Warsaw, Warsaw, Poland. <sup>634</sup>Department of Medicine, Universidad Autónoma de Madrid (Autonomous University of Madrid), Madrid, Spain. <sup>635</sup>Department of Nephrology and Hypertension, The Institute for Health Research Foundation Jiménez Díaz University Hospital, Madrid, Spain. <sup>636</sup>One Health Global Research Group, Universidad de las Americas (University of the Americas), Quito, Ecuador. <sup>637</sup>School of Medicine, Western Sydney University, Bathurst, New South Wales, Australia. <sup>638</sup>Department of Optometry and Vision Science, University of KwaZulu-Natal, KwaZulu-Natal, South Africa. <sup>639</sup>Faculty of Medicine, University Ferhat Abbas of Setif, Sétif, Algeria. <sup>640</sup>Division of Infectious Diseases, University Hospital of Setif, Sétif, Algeria. <sup>641</sup>Department of Medicine, University of Ibadan, Ibadan, Nigeria. <sup>642</sup>Department of Medicine, University College Hospital, Ibadan, Ibadan, Nigeria. <sup>643</sup>Department of Respiratory Medicine, Jagadguru Sri Shivarathreeswara University, Mysore, India. <sup>644</sup>National School of Public Health, Institute of Health Carlos III, Madrid, Spain. <sup>645</sup>Department of Forensic Medicine and Toxicology, Manipal Academy of Higher Education, Mangalore, India. <sup>646</sup>Department of Ophthalmology, Heidelberg University, Heidelberg, Germany. <sup>647</sup>Department of Science and Mathematics, Deree-The American College of Greece, Athens, Greece. <sup>648</sup>Department of Biophysics, University of Athens, Athens, Greece. <sup>649</sup>Vision and Eye Research Institute, Anglia Ruskin University, Cambridge, UK. <sup>650</sup>Department of Community Medicine, All India Institute of Medical Sciences, Jammu, India. <sup>651</sup>Division of Health Policy and Management, University of Minnesota, Minneapolis, MN, USA. <sup>652</sup>Department of Biomedical Data Science, Stanford University, Stanford, CA, USA. <sup>653</sup>Department of Medical Sciences, University of Torino, Torino, Italy. <sup>654</sup>Department of Imaging, AOU Città della Salute e della Scienza di Torino, Torino, Italy. <sup>655</sup>Global Health Governance Programme, University of Edinburgh, Edinburgh, UK. <sup>656</sup>School of Dentistry, University of Leeds, Leeds, UK. <sup>657</sup>Natnov Bioscience, Tahalia, India. <sup>658</sup>College of Dental Medicine, Roseman University of Health Sciences, South Jordan, UT, USA. <sup>659</sup>Second Department of Cardiology, Aristotle University of Thessaloniki, Thessaloniki, Greece. <sup>660</sup>Department of Genetics, Yale University, New Haven, CT, USA. <sup>661</sup>Department of Outpatient, Taihe Hospital, Hubei University of Medicine, Shiyan, China. <sup>662</sup>Australian Institute of Health Innovation, Macquarie University, Sydney, New South Wales, Australia. <sup>663</sup>School of Population Health, Curtin University, Bentley, Western Australia, Australia. <sup>664</sup>Centre for Fertility and Health, Norwegian Institute of Public Health, Oslo, Norway. <sup>665</sup>Social and Economic Survey Research Institute, Qatar University, Doha, Qatar. <sup>666</sup>Mario Negri Institute for Pharmacological Research, Bergamo, Italy. <sup>667</sup>School of Pharmacy, University of Nizwa, Nizwa, Oman. <sup>668</sup>International Center of Medical Sciences Research, International Center of Medical Sciences Research, Islamabad, Pakistan. <sup>669</sup>Riphah International University, Islamabad, Pakistan. <sup>670</sup>College of Health Sciences (CHS), VinUniversity, Hanoi, Viet Nam. <sup>671</sup>Clinical Academic Department of Pediatrics, University Medical Center (UMC), Astana, Kazakhstan. <sup>672</sup>Department of Data Management and Analysis, The International Clinical Epidemiology Network (INCLIN) Trust International, New Delhi, India. <sup>673</sup>Non-communicable Diseases Research Center, Bam University of Medical Sciences, Bam, Iran. <sup>674</sup>Department of Community Medicine and Public Health, Tribhuvan University, Kathmandu, Nepal. <sup>675</sup>T.H. Chan School of Public Health, Harvard University, Boston, MA, USA. <sup>676</sup>Department of Humanities and Social Sciences, National Institute of Technology Rourkela, Rourkela, India. <sup>677</sup>Department of Clinical Research and Epidemiology, Institute of Liver and Biliary Sciences, New Delhi, India. <sup>678</sup>Centre for Dental Education and Research, All India Institute of Medical Sciences, New Delhi, India. <sup>679</sup>Department of Biostatistics, Epidemiology, and Informatics, University of Pennsylvania, Philadelphia, PA, USA. <sup>680</sup>Cihan University-Sulaimaniya Research Center, Cihan University-Sulaimaniya, Sulaymaniyah, Iraq. <sup>681</sup>Department of Neonatology, Case Western Reserve University, Cleveland, OH, USA. <sup>682</sup>Department of Community Medicine, ESI

Post Graduate Institute of Medical Science and Research, Bengaluru, India. <sup>683</sup>Department of Medical Oncology, Cancer Institute (W.I.A), Chennai, India. <sup>684</sup>Osh State University, Osh, Kyrgyzstan. <sup>685</sup>Director of Central Asia Research Collaboration Group, Asfendiyarov Kazakh National Medical University, Almaty, Kazakhstan. <sup>686</sup>Sina Trauma and Surgery Research Center, Tehran University of Medical Sciences, Tehran, Iran. <sup>687</sup>Department of Population Science and Human Resource Development, University of Rajshahi, Rajshahi, Bangladesh. <sup>688</sup>College of Medicine and Health Sciences, National University of Science and Technology, Sohar, Oman. <sup>689</sup>Institute of Health and Wellbeing, Federation University Australia, Berwick, Victoria, Australia. <sup>690</sup>School of Nursing and Midwifery, La Trobe University, Melbourne, Victoria, Australia. <sup>691</sup>Future Technology Research Center, National Yunlin University of Science and Technology, Yunlin, Taiwan. <sup>692</sup>Student Research Committee, Shahid Beheshti University of Medical Sciences, Tehran, Iran. <sup>693</sup>Department of Public Health, Torbat Jam Faculty of Medical Sciences, Torbat Jam, Iran. <sup>694</sup>Department of Nutrition Science, Muhammadiyah University of Surakarta, Surakarta, Indonesia. <sup>695</sup>Department of Cardiology, Dow University of Health Sciences, Karachi, Pakistan. <sup>696</sup>Department of Medicine, Dow University of Health Sciences, Karachi, Pakistan. <sup>697</sup>Centre for Chronic Disease Control, New Delhi, India. <sup>698</sup>Department of General Medicine, Manipal Academy of Higher Education, Manipal, India. <sup>699</sup>Department of Clinical Sciences, University of Sharjah, Sharjah, United Arab Emirates. <sup>700</sup>Department of Cardiology, Mansoura University, Mansoura, Egypt. <sup>701</sup>Department of Radiology, Stanford University, Stanford, CA, USA. <sup>702</sup>Translational Health Research Institute, Western Sydney University, Sydney, New South Wales, Australia. <sup>703</sup>Department of Community Medicine, Shaheed Nirmal Mahto Medical College and Hospital, Dhanbad, India. <sup>704</sup>Health Economics and Outcomes Research Department, Agios Pharmaceuticals, Cambridge, MA, USA. <sup>705</sup>Department of Pharmaceutical Economics and Policy, Massachusetts College of Pharmacy and Health Sciences, Boston, MA, USA. <sup>706</sup>Department of Community Medicine, Manipal Academy of Higher Education, Manipal, India. <sup>707</sup>Department of Oral Pathology, Microbiology and Forensic Odontology, Sharavathi Dental College and Hospital, Shimogga, India. <sup>708</sup>Department of Family Medicine, Rajarata University of Sri Lanka, Anuradhapura, Sri Lanka. <sup>709</sup>Department of Primary Care and Public Health, Imperial College London, London, UK. <sup>710</sup>Academic Public Health England, Public Health England, London, UK. <sup>711</sup>Department of Biological Sciences, King Abdulaziz University, Jeddah, Egypt. <sup>712</sup>Department of Protein Research, Research and Academic Institution, Alexandria, Egypt. <sup>713</sup>Saveetha Medical College and Hospital, Saveetha University, Chennai, India. <sup>714</sup>Centre for Excellence in Pharmaceutical Sciences, North-West University, Potchefstroom, South Africa. <sup>715</sup>Department of Epidemiology and Biostatistics, Rafsanjan University of Medical Sciences, Rafsanjan, Iran. <sup>716</sup>Department of Pharmacy, Shaheed Benazir Bhutto University Sheringal Pakistan, Dir Upper, Pakistan. <sup>717</sup>Department of Surgery, University of Minnesota, Minneapolis, MN, USA. <sup>718</sup>Department of Surgery, University Teaching Hospital of Kigali, Kigali, Rwanda. <sup>719</sup>Community Health Department, Federal University of Ceará, Fortaleza, Brazil. <sup>720</sup>Department of Clinical Research, University of Sao Paulo, Ribeirão Preto, Brazil. <sup>721</sup>Gilbert and Rose-Marie Chagoury School of Medicine, Lebanese American University, Beirut, Lebanon. <sup>722</sup>Department of Community Medicine, Government Medical College, Chandigarh, India. <sup>723</sup>Department of Public Health, Bangladesh Open University, Gazipur, Bangladesh. <sup>724</sup>Department of Analytical and Applied Economics, Utkal University, Bhubaneswar, India. <sup>725</sup>RUSA Centre of Excellence in Public Policy and Governance, Utkal University, Bhubaneswar, India. <sup>726</sup>Faculty of Medicine, Quest International University Perak, Ipoh, Malaysia. <sup>727</sup>Advanced Campus Governador Valadares, Juiz de Fora Federal University, Governador Valadares, Brazil. <sup>728</sup>Department of Internal Medicine, University of Botswana, Gaborone, Botswana. <sup>729</sup>Cardiovascular Department, Zagazig University, Zagazig, Egypt. <sup>730</sup>Department of Epidemiology, Shahid Beheshti University of Medical Sciences, Tehran, Iran. <sup>731</sup>Department of Pharmaceuticals, Bihar College of Pharmacy, Patna, India. <sup>732</sup>College of Medicine, University of Sharjah, Sharjah, United Arab Emirates. <sup>733</sup>School of Population Health, University of New South Wales, Sydney, New South Wales, Australia. <sup>734</sup>Operational Research Center in Healthcare, Near East University (NEU), Nicosia Cyprus, Turkey. <sup>735</sup>International Center of Medical Sciences Research (ICMSR), Islamabad, Pakistan. <sup>736</sup>Sharjah Institute of Medical Sciences, University of Sharjah, Sharjah, United Arab Emirates. <sup>737</sup>Center for Global Health Research, Saveetha University, Chennai, India. <sup>738</sup>Biotechnology Research Center, Mashhad University of Medical Sciences, Mashhad, Iran. <sup>739</sup>Department of Analytical & Applied Economics, Utkal University, Bhubaneswar, India. <sup>740</sup>Department of Community Medicine and Family Medicine, All India Institute of Medical Sciences, Bathinda, India. <sup>741</sup>Department of Nutrition and Dietetics, Cihan University, Erbil, Iraq. <sup>742</sup>Department of Oral and Maxillofacial Surgery, University College Hospital, Ibadan, Ibadan, Nigeria. <sup>743</sup>Department of Integrated Health Education, Federal University of Espirito Santo, Vitória, Brazil. <sup>744</sup>Faculty of Allied Health Sciences, Health Services Academy, Islamabad, Pakistan. <sup>745</sup>Public Health and Community Medicine Department, Cairo University, Giza, Egypt. <sup>746</sup>Institute of Epidemiology and Preventive Medicine, National Taiwan University, Taipei, Taiwan. <sup>747</sup>Benang Merah Research Center (BMRC), Minahasa Utara, Indonesia. <sup>748</sup>Department of Anatomy, Ras Al Khaimah Medical and Health Sciences University, Ras Al Khaimah, United Arab Emirates. <sup>749</sup>Department of Entomology, Ain Shams University, Cairo, Egypt. <sup>750</sup>Medical Ain Shams Research Institute (MASRI), Ain Shams University, Cairo, Egypt. <sup>751</sup>Department of Pediatrics, SRM Medical College Hospital And Research Centre, Chennai, India. <sup>752</sup>Center for Clinical and Epidemiological Research, University of São Paulo, São Paulo, Brazil. <sup>753</sup>Independent Consultant, Thiruvananthapuram, India. <sup>754</sup>Department of Pharmacology, All India Institute of Medical Sciences, Jodhpur, India. <sup>755</sup>Indira Gandhi Medical College and Research Institute, Puducherry, India. <sup>756</sup>Department of Food Processing Technology, West Bengal State Council of Technical Education, Malda, India. <sup>757</sup>Department of Oral Pathology and Microbiology, Dr. D. Y. Patil Vidyapeeth, Pune, India. <sup>758</sup>Department of Geriatric and Long Term Care, Hamad Medical Corporation, Doha, Qatar. <sup>759</sup>Faculty of Health & Social Sciences, Bournemouth University, Bournemouth, UK. <sup>760</sup>UGC Centre of Advanced Study in Psychology, Utkal University, Bhubaneswar, India. <sup>761</sup>Udyam-Global Association for Sustainable Development, Bhubaneswar, India. <sup>762</sup>Department of Public Health Sciences, University of North Carolina at Charlotte, Charlotte, NC, USA. <sup>763</sup>Maternal and Child Health Division (MCHD), International Centre for Diarrhoeal Disease Research, Dhaka, Bangladesh. <sup>764</sup>Department of Post-Harvest Technology and Marketing, Patuakhali Science and Technology University, Patuakhali, Bangladesh. <sup>765</sup>Faculty of Medicine, Katholieke Universiteit Leuven, Leuven, Belgium. <sup>766</sup>Department of Cardiovascular Sciences, Katholieke Universiteit Leuven, Leuven, Belgium. <sup>767</sup>Saveetha Dental College and Hospitals, Saveetha University, Chennai, India. <sup>768</sup>Dr. D. Y. Patil Dental College and Hospital, Dr. D. Y. Patil Vidyapeeth, Pune, India. <sup>769</sup>College of Pharmacy, University of Sharjah, Sharjah, United Arab Emirates. <sup>770</sup>Research Institute of Medical & Health Sciences, University of Sharjah, Sharjah, United Arab Emirates. <sup>771</sup>Department of Biomedical Sciences, Gulf Medical University, Ajman, United Arab Emirates. <sup>772</sup>Digestive Diseases Research Institute, Tehran University of Medical Sciences, Tehran, Iran. <sup>773</sup>Non-communicable Disease Research Center, Shiraz University of Medical Sciences, Shiraz, Iran. <sup>774</sup>Department of Medicine and Surgery, Government Doon Medical College, Dehradun, India. <sup>775</sup>Pharmacy Department, Sanata Dharma University, Yogyakarta, Indonesia. <sup>776</sup>Department of Neurology, Tehran University of Medical Sciences, Tehran, Iran. <sup>777</sup>Department of Physics, The University of Lahore, Lahore, Pakistan. <sup>778</sup>Center for Medical and Bio-Allied Health Sciences Research, Ajman University, Ajman, United Arab Emirates. <sup>779</sup>Independent Consultant, Karachi, Pakistan. <sup>780</sup>Department of Anesthesiology, Shahid Beheshti University of Medical Sciences, Tehran, Iran. <sup>781</sup>Department of Pathology and Laboratory Medicine, Northwell Health, New York, NY, USA. <sup>782</sup>School of Medicine, Alborz University of Medical Sciences, Karaj, Iran. <sup>783</sup>Centre for Interdisciplinary Research in Basic Sciences (CIRBSc), Jamia Millia Islamia, New Delhi, India. <sup>784</sup>Science Department, Kazakh National Medical University, Almaty, Kazakhstan. <sup>785</sup>College of Nursing and Health Sciences, Jazan University, Jazan, Saudi Arabia. <sup>786</sup>Amity Institute of Public Health, Amity University, Noida, India. <sup>787</sup>Department for Evidence-based Medicine and Evaluation, University for Continuing Education Krems, Krems, Austria. <sup>788</sup>Sina Hospital, Tehran University of Medical Sciences, Tehran, Iran. <sup>789</sup>Department of Hemato-oncology, Fortis Hospital, Noida, India. <sup>790</sup>Institute of Forensic Science & Criminology, Panjab

University, Chandigarh, India. <sup>791</sup>Department of Ophthalmology, Harvard University, Boston, MA, USA. <sup>792</sup>Ophthalmic Research Center (ORC), Shahid Beheshti University of Medical Sciences, Tehran, Iran. <sup>793</sup>Department of Health in Disasters and Emergencies, Shahrekord University of Medical Sciences, Shahrekord, Iran. <sup>794</sup>Department of Pharmacology, Manipal Academy of Higher Education, Manipal, India. <sup>795</sup>K S Hegde Medical Academy, Nitte University, Mangalore, India. <sup>796</sup>Manipal College of Dental Sciences, Mangalore, Manipal Academy of Higher Education, Mangalore, India. <sup>797</sup>Department of Gastroenterology and Hepatology, Manipal Academy of Higher Education, Udupi, India. <sup>798</sup>Finnish Institute of Occupational Health, Helsinki, Finland. <sup>799</sup>Department of Veterinary Public Health and Preventive Medicine, Usmanu Danfodiyo University, Sokoto, Nigeria. <sup>800</sup>Department of Experimental Research, Medical University Pleven, Pleven, Bulgaria. <sup>801</sup>Department of Genetics, Sofia University 'St. Kliment Ohridski', Sofia, Bulgaria. <sup>802</sup>Department of Research and Academics, Kathmandu Cancer Center, Bhaktapur, Nepal. <sup>803</sup>Unit of Basic Medical Sciences, University of Khartoum, Khartoum, Sudan. <sup>804</sup>Department of Medical Microbiology and Infectious Diseases, Erasmus University, Rotterdam, The Netherlands. <sup>805</sup>Sport Physical Activity and Health Research & Innovation Center (SPRINT), Polytechnic Institute of Guarda, Guarda, Portugal. <sup>806</sup>CICS-UBI Health Sciences Research Center, University of Beira Interior, Covilhã, Portugal. <sup>807</sup>Department of Biochemistry, Central University of Punjab, Bathinda, India. <sup>808</sup>Department of Pharmacology, Government Medical College and Hospital, Chandigarh, India. <sup>809</sup>Department of Paediatrics, All India Institute of Medical Sciences, Bilaspur, India. <sup>810</sup>School of Medicine, Baylor College of Medicine, Houston, TX, USA. <sup>811</sup>Department of Medicine Service, US Department of Veterans Affairs (VA), Houston, TX, USA. <sup>812</sup>Faculty of Medicine and Health Sciences, Shree Guru Gobind Singh Tricentenary University, Gurugram, India. <sup>813</sup>Department of Radiodiagnosis, All India Institute of Medical Sciences, Bathinda, India. <sup>814</sup>Department of Human Genetics, Punjabi University, Patiala, India. <sup>815</sup>Department of Systemic Pathology, Touro College of Osteopathic Medicine, Middletown, NY, USA. <sup>816</sup>Department of Pathology, American University of the Caribbean School of Medicine, Cupecoy, Sint Maarten. <sup>817</sup>Department of Biochemistry, American University of Integrative Sciences, Bridgetown, Barbados. <sup>818</sup>Student Research Committee, Urmia University of Medical Sciences, Urmia, Iran. <sup>819</sup>School of Medicine, Babol University of Medical Sciences, Babol, Iran. <sup>820</sup>Department of Global Health, University of Washington, Seattle, WA, USA. <sup>821</sup>Department of Pharmacology, RAK Medical and Health Sciences University, Ras Al Khaimah, United Arab Emirates. <sup>822</sup>Department of Medical Sciences, Sunway University, Subang Jaya, Malaysia. <sup>823</sup>Department of Clinical Pathology, Brawijaya University, Malang, Indonesia. <sup>824</sup>Hospital Central Laboratory, Dr. Saiful Anwar General Hospital, Malang, Indonesia. <sup>825</sup>Department of Clinical Research and Development, LUXMED Group, Warsaw, Poland. <sup>826</sup>Collegium Medicum, John Paul II Catholic University of Lublin, Lublin, Poland. <sup>827</sup>Northwestern University, Chicago, IL, USA. <sup>828</sup>Department of Pharmacology, All India Institute of Medical Sciences, Deoghar, India. <sup>829</sup>Department of Medical Informatics, Mashhad University of Medical Sciences, Mashhad, Iran. <sup>830</sup>Clinical Research Development Unit, Mashhad University of Medical Sciences, Mashhad, Iran. <sup>831</sup>Department of Basic Medical Sciences, Islamic Azad University, Mashhad, Iran. <sup>832</sup>Department of Internal Medicine, Islamic Azad University, Mashhad, Iran. <sup>833</sup>Department of Environmental, Agricultural and Occupational Health, University of Nebraska Medical Center, Omaha, NE, USA. <sup>834</sup>Sri Ramachandra Medical College and Research Institute, Chennai, India. <sup>835</sup>Department of Pathology, Alexandria University, Alexandria, Egypt. <sup>836</sup>Department of Dermatology, Carol Davila University of Medicine and Pharmacy, Bucharest, Romania. <sup>837</sup>Department of Dermato-Venereology, Dr. Victor Babes Clinical Hospital of Infectious Diseases and Tropical Diseases, Bucharest, Romania. <sup>838</sup>Department of Epidemiology, Stellenbosch University, Cape Town, South Africa. <sup>839</sup>Department of Medicine, Northlands Medical Group, Omuthiya, Namibia. <sup>840</sup>National Research and Innovation Agency, Jakarta, Indonesia. <sup>841</sup>Nutrition and Clinical Services Division, International Centre for Diarrhoeal Disease Research, Dhaka, Bangladesh. <sup>842</sup>Taking Our Best Shot, Houston, TX, USA. <sup>843</sup>Department of Research and Innovation, Enventure Medical Innovation, Houston, TX, USA. <sup>844</sup>School of Nursing and Midwifery, Debre Berhan University, Debre Berhan, Ethiopia. <sup>845</sup>Department of Public Health, Dire Dawa University, Dire Dawa, Ethiopia. <sup>846</sup>Pediatric Intensive Care Unit, King Saud University, Riyadh, Saudi Arabia. <sup>847</sup>Rheumatology and Immunology Unit, Mansoura University, Mansoura, Egypt. <sup>848</sup>Department of Family and Preventive Medicine, Emory University, Atlanta, GA, USA. <sup>849</sup>Department of Gastroenterology, St. Luke's Hospital, Patanamthitta, India. <sup>850</sup>Faculty of Public Health, Universitas Sam Ratulangi (Sam Ratulangi University), Manado, Indonesia. <sup>851</sup>Timiryazev Institute of Plant Physiology, Russian Academy of Sciences, Moscow, Russia. <sup>852</sup>Laboratory of Public Health Indicators Analysis and Health Digitalization, Moscow Institute of Physics and Technology, Moscow, Russia. <sup>853</sup>Department of Health, Children's Hospital 1, Ho Chi Minh City, Viet Nam. <sup>854</sup>School of Biomedical Engineering, University of Technology Sydney, Sydney, New South Wales, Australia. <sup>855</sup>Department of Internal Medicine, University of Medicine and Pharmacy at Ho Chi Minh City, Ho Chi Minh City, Viet Nam. <sup>856</sup>Department of Business Analytics, University of Massachusetts Dartmouth, Dartmouth, MA, USA. <sup>857</sup>Department of Clinical and Experimental Medicine, University of Pisa, Pisa, Italy. <sup>858</sup>Faculty of Medicine, Tan Tao University, Long An, Viet Nam. <sup>859</sup>Department of Occupational Health and Safety, University of Development, Surabaya, Indonesia. <sup>860</sup>Department of Internal Medicine, Wake Forest University, Winston-Salem, NC, USA. <sup>861</sup>Department of Biosciences and Biotechnology, University of Medical Sciences, Ondo, Nigeria. <sup>862</sup>International Center for Chemical and Biological Sciences, University of Karachi, Karachi, Pakistan. <sup>863</sup>College of Medicine and Public Health, Flinders University, Adelaide, South Australia, Australia. <sup>864</sup>Lahore Business School, The University of Lahore, Lahore, Pakistan. <sup>865</sup>Department of Dietetics and Nutrition, Jimma University, Jimma, Ethiopia. <sup>866</sup>Faculty of Medicine, Tehran University of Medical Sciences, Tehran, Iran. <sup>867</sup>Department of Informatics and Radiology, Mayo Clinic, Rochester, MN, USA. <sup>868</sup>College of Health and Sport Sciences, University of Bahrain, Zallaq, Bahrain. <sup>869</sup>Urmia University of Medical Sciences, Urmia, Iran. <sup>870</sup>College of Public Health and Tropical Medicine, Jazan University, Jazan, Saudi Arabia. <sup>871</sup>UKK Institute, Tampere, Finland. <sup>872</sup>Faculty of Medicine and Health Technology, Tampere University, Tampere, Finland. <sup>873</sup>Department of Zoology, Central University of Punjab, Bathinda, India. <sup>874</sup>Department of Human Genetics & Molecular Biology, Bharathiar University, Coimbatore, India. <sup>875</sup>Department of Surgery, University of Southampton, Southampton, UK. <sup>876</sup>College of Medicine and Veterinary Medicine, University of Edinburgh, Edinburgh, UK. <sup>877</sup>Department of Health Policy and Management, Johns Hopkins University, Baltimore, MD, USA. <sup>878</sup>Department of Cardiology, Icahn School of Medicine at Mount Sinai, New York, NY, USA. <sup>879</sup>Department of Medical Oncology, University of Medicine and Pharmacy 'Grigore T Popa' Iasi, Iasi, Romania. <sup>880</sup>Department of Medical Oncology, Regional Institute of Oncology, Iasi, Romania. <sup>881</sup>NUST School of Health Sciences, National University of Sciences and Technology (NUST), Islamabad, Pakistan. <sup>882</sup>Operational Research Center in Healthcare, Near East University, Nicosia, Turkey. <sup>883</sup>School of Life Course and Population Sciences, King's College London, London, UK. <sup>884</sup>Department of Parasitology, Rajarata University of Sri Lanka, Anuradhapura, Sri Lanka. <sup>885</sup>Institute of Clinical Epidemiology, Public Health, Health Economics, Medical Statistics and Informatics, Medical University Innsbruck, Innsbruck, Austria. <sup>886</sup>Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK. <sup>887</sup>Department of Public Health, Samara University, Samara, Ethiopia. <sup>888</sup>Department of Emergency Medicine and Critical Care Nursing, Bahir Dar University, Bahir Dar, Ethiopia. <sup>889</sup>Cardiovascular Program, The George Institute for Global Health, Sydney, New South Wales, Australia. <sup>890</sup>Department of Environmental Health and Epidemiology, National Institute for Research in Environmental Health, Bhopal, India. <sup>891</sup>Department of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran. <sup>892</sup>Department of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran. <sup>893</sup>The George Institute for Global Health, Imperial College London, London, UK. <sup>894</sup>Department of Family Medicine, St. Paul's Hospital Millennium Medical College, Addis Ababa, Ethiopia. <sup>895</sup>Independent Consultant, Addis Ababa, Ethiopia. <sup>896</sup>Public Health Department (Public Health Nutrition Unit), Dire Dawa University, Dire Dawa, Ethiopia. <sup>897</sup>Department of Epidemiology, Xuzhou Medical University, Xuzhou, China. <sup>898</sup>Department of Biostatistics, University of Toyama, Toyama, Japan. <sup>899</sup>Department of Public Health, Juntendo University, Tokyo, Japan.

<sup>900</sup>Basic Sciences Department, University of Duhok, Duhok, Iraq. <sup>901</sup>Nephology Department, Hospital San Juan de Dios, Tarija, Bolivia. <sup>902</sup>San Pablo Catholic University Tarija Bolivia, Tarija, Bolivia. <sup>903</sup>Research and Development Department, Sina Medical Biochemistry Technologies, Shiraz, Iran. <sup>904</sup>Department of Public Health, University of Hail, Hail, Saudi Arabia. <sup>905</sup>Department of Statistics, Mekelle University, Mekelle, Ethiopia. <sup>906</sup>Department of Biostatistics, Jimma University, Jimma, Ethiopia. <sup>907</sup>School of Public Health, Peking University, Beijing, China. <sup>908</sup>Department of International Health, Johns Hopkins University, Baltimore, MD, USA. <sup>909</sup>Medical Oncology Department of Gastrointestinal Cancer, Cancer Hospital of Dalian University of Technology, Shenyang, China. <sup>910</sup>School of Biomedical Engineering, Dalian University of Technology, Dalian, China. <sup>911</sup>Jockey Club School of Public Health and Primary Care, The Chinese University of Hong Kong, Hong Kong, China. <sup>912</sup>Atchabarov Scientific-Research Institute of Fundamental and Applied Medicine, Kazakh National Medical University, Almaty, Kazakhstan. <sup>913</sup>Department of Epidemiology, University of Washington, Seattle, WA, USA. <sup>914</sup>Institute of Public Health and Social Sciences, Khyber Medical University, Peshawar, Pakistan. <sup>915</sup>Endocrinology and Metabolism Research Center, Hormozgan University of Medical Sciences, Bandar Abbas, Iran. <sup>916</sup>Department of Clinical and Community Pharmacy, An-Najah National University, Nablus, Palestine. <sup>917</sup>Clinical Research Centre, An-Najah National University Hospital, Nablus, Palestine.

## Methods

### Overview

This paper was produced as part of the GBD Collaborator Network and in accordance with the GBD Protocol<sup>9</sup>. The GBD 2021 offers a comprehensive evaluation of the global disease burden, quantifying health loss from 371 diseases across 204 countries and territories between 1990 and 2021 (ref. 2). This work involved collaboration from over 10,000 experts in more than 150 countries, contributing to data provision, analysis and review necessary for generating GBD metrics, such as incidence, prevalence, cause-specific mortality, years of life lost, YLDs and DALYs. The present study specifically targeted dietary iron deficiency, distinguishing it from other forms of iron deficiency. We aimed to assess the health burden arising directly from undernutrition, thereby underscoring the importance of public health interventions focused on dietary improvement. This study adheres to the Guidelines for Accurate and Transparent Health Estimate Reporting (GATHER)<sup>2</sup> (Supplementary Table 1), with analyses conducted using R software version 4.3.2 (R Foundation) and Python version 3.11.4 (Python Software Foundation). The supplementary methods contain further details on specific methodologies.

### Case definition

Dietary iron deficiency is defined as iron deficiency directly resulting from inadequate dietary iron intake that is insufficient to meet the body's requirements<sup>2</sup>. This definition emphasizes iron deficiency as a modifiable public health burden and distinguishes it from other conditions caused by non-dietary factors such as chronic diseases, physiological conditions or metabolic disorders. This study models dietary iron deficiency as a preventable cause of anemia by focusing on cases where insufficient dietary iron intake leads to physiological complications<sup>31</sup>.

Menstruation increases physiological iron requirements, and when dietary intake does not meet these elevated needs, this is classified as dietary iron deficiency. However, pathological conditions, such as uterine fibroids or heavy menstrual bleeding, are modeled separately and are excluded from dietary iron deficiency estimates. This distinction ensures that dietary iron deficiency represents preventable cases solely attributable to insufficient dietary iron intake, independent of other pathological conditions. This study offers targeted nutritional strategies to mitigate dietary iron deficiency at the population level, providing a strong foundation for actionable public health policy. The scope of dietary iron deficiency excludes iron deficiency that occurs independent of dietary intake, and such non-dietary causes are modeled independently within the GBD framework<sup>2</sup> (Extended Data Fig. 9). Excluded causes include the following:

1. Neglected tropical diseases. Tropical diseases such as malaria, hookworm infections and schistosomiasis cause anemia through chronic blood loss or hemolysis, which is not related to dietary iron intake. These diseases are primarily addressed through disease management programs rather than nutritional interventions.
2. Gastrointestinal disorders. Gastrointestinal disorders, such as inflammatory bowel disease, chronic gastritis and ulcers, impair iron absorption or result in chronic blood loss. These conditions require clinical interventions beyond dietary therapy.
3. Chronic health conditions and metabolic disorders. Systemic diseases such as cirrhosis and chronic kidney disease affect iron metabolism or cause blood loss, which leads to anemia independent of dietary intake. These conditions require specialized medical management.
4. Reproductive health factors. Conditions such as heavy menstrual bleeding, uterine fibroids and maternal hemorrhage cause anemia through acute or chronic blood loss. Menstruation increases physiological iron requirements, but anemia that results from an inability to meet the increased need is classified as dietary iron deficiency. On the other hand, anemia caused

by a pathological condition (for example, uterine fibroids) is modeled separately.

5. Micronutrient deficiencies. Deficiencies of other micronutrients, such as vitamin A, can affect hemoglobin synthesis and iron metabolism, increasing the prevalence of anemia. To address this, a comprehensive nutritional strategy is needed, not just iron supplementation.

By restricting the study's focus to dietary iron deficiency, this analysis enables a more precise estimation that is directly preventable and manageable through dietary interventions. This approach emphasizes the need for targeted nutritional strategies to reduce dietary iron deficiency at the population level, aligning with public health policies aimed at alleviating undernutrition.

### Geographic locations of the analysis

We developed estimates for 204 countries and territories, categorized into 21 regions and 7 broad super-regions. The super-regions include Central Europe, Eastern Europe and Central Asia; High-Income; Latin America and the Caribbean; North Africa and the Middle East; South Asia; Southeast Asia, East Asia and Oceania; and sub-Saharan Africa. In GBD 2021, we continue to conduct subnational analyses for countries included in earlier cycles, such as Brazil, China, Ethiopia, India, Indonesia, Iran, Italy, Japan, Kenya, Mexico, New Zealand, Nigeria, Norway, Pakistan, Russia, the Philippines, Poland, South Africa, Sweden, the UK and the USA. Analyses are carried out at the first administrative level within each country, except for New Zealand (by Māori ethnicity), Sweden (by Stockholm and non-Stockholm areas), the UK (by local government authorities) and the Philippines (by provinces).

At the most granular spatial resolution, we generated estimates for 983 unique locations. As with GBD 2019, GBD 2021 continues to use a classification of standard and non-standard locations. Standard GBD locations encompass all subnationals from countries with high-quality data and populations exceeding 200 million, along with all other countries. This classification includes subnationals for China, India, the USA and Brazil but it excludes Indonesia; data for China, India, the USA and Brazil are also available at the national level. All other countries with subnational estimates are defined as non-standard locations.

### Input data and data processing

For GBD 2021, dietary iron deficiency estimates were derived using a substantial portion of data collected from population-based surveys, peer-reviewed studies and governmental reports<sup>32</sup>. These data sources included the Demographic and Health Survey, the Multiple Indicator Cluster Survey series, national micronutrient surveys and other nutrition surveillance systems at national and subnational levels. To account for regional and demographic variability, the modeling framework incorporated region-specific covariates, such as the Healthcare Access and Quality Index (HAQI), malaria prevalence, and child undernutrition indicators. These adjustments helped dietary iron deficiency estimates align more closely with population-specific dietary intake patterns while minimizing the influence of non-dietary factors.

During the data extraction process, we focused on demographic variables (such as location, sex and age), survey design variables (such as sampling strategy and sampling weights) and population estimate variables (such as prevalence or proportion), along with measures of uncertainty (such as standard error, confidence interval, sample size and number of cases). Using available microdata, data were categorized based on sex and age groups, including 6–11 months, 12–23 months, 2–4 years, 5–9 years, 10–14 years and similar increments up to 95 years and older. When detailed age data were unavailable or sample sizes were small, broader age categories were used to ensure robust estimates.

To adjust for altitude-related variations in hemoglobin levels, altitude-adjusted data were included following the WHO adjustment formula<sup>30</sup>. However, no additional adjustments were made for factors

such as smoking status, hemoglobin sampling techniques or other analytical methods. This systematic approach ensured the reliability and accuracy of dietary iron deficiency estimates while accommodating regional and demographic variability.

### Modeling strategy

The burden of dietary iron deficiency was estimated using a two-stage process as follows:

1. Anemia envelope estimation. Hemoglobin distributions were modeled using Spatio-Temporal Gaussian Process Regression (ST-GPR) across 204 countries, disaggregated by age, sex, location and year. The ST-GPR model integrates covariates such as age-specific fertility rates, HIV prevalence and HAQI, allowing for refined estimates in regions with limited data. The model included random effects on location to account for regional variability in dietary iron deficiency prevalence. The variability of iron deficiency across anemia severity levels was not assumed to be constant; instead, severity-level-specific estimates were independently evaluated across different population subgroups.
2. Cause attribution for dietary iron deficiency. To estimate the contribution of dietary iron deficiency, the GBD study applied a globally systematic modeling process that integrates multiple high-quality datasets, such as the WHO Vitamin and Mineral Nutrition Information System, the Demographic and Health Survey, UNICEF Multiple Indicator Cluster Surveys and the National Health and Nutrition Examination Survey. Within this framework, dietary iron deficiency was estimated to account for 80.5% of residual anemia cases globally, reflecting its important role as a preventable cause of anemia. Counterfactual hemoglobin distributions, representing hypothetical levels without specific causes, were used to proportionally allocate anemia cases to dietary iron deficiency and other etiologies.

To estimate anemia prevalence attributable to specific causes, the analysis used modeled hemoglobin distributions for each location, year, age and sex, incorporating cause-specific prevalence and associated hemoglobin shifts. The estimated cause prevalence was multiplied by hemoglobin shifts to derive prevalence-weighted distribution shifts specific to each demographic and causal factor. Subsequently, mean hemoglobin estimates were adjusted using these weighted shifts, maintaining the original variance to re-estimate distributions. The difference between initial and counterfactual distributions across anemia severity levels was calculated to determine the prevalence attributable to each cause.

Unique hemoglobin shifts for dietary iron deficiency were derived from studies on iron fortification and supplementation intervention trials, applying a standardized shift of  $4.01 \text{ g l}^{-1}$ . This approach avoids relying on fixed proportions and, instead, dynamically evaluates dietary iron deficiency contributions across regions, age groups and levels of anemia severity. Variability in dietary contributions was captured by assessing unique counterfactual hemoglobin shifts for each anemia severity level, considering population-specific dietary intake patterns and regional factors. To ensure robustness, a minimum of 10% of anemia cases for each location, year, age and sex were allocated to residual causes. These residuals were proportionally distributed among dietary iron deficiency, other neglected tropical diseases, other infectious diseases and other hemoglobinopathies or hemolytic anemias, based on global age-specific and sex-specific proportions. Additionally, the GBD framework employs adjustments for demographic and regional factors, such as malaria prevalence, healthcare access and nutritional status, to refine hemoglobin distributions and anemia attribution. These adjustments ensure that estimates of dietary iron deficiency reflect regional and demographic variability, avoiding universal assumptions and enhancing the precision of burden estimates.

This proportional approach minimized confounding effects from chronic infections and metabolic disorders, allowing for a clear

distinction between anemia due to dietary iron deficiency and other causes<sup>30</sup> (Extended Data Fig. 8). To ensure consistency, residual causes were allocated a minimum of 10% of anemia cases, reflecting findings from large-scale nutritional surveys.

### Impact of inflammation and ferritin interpretation

The GBD model accounts for inflammatory conditions such as infections and chronic diseases that may complicate ferritin interpretation and dietary iron deficiency estimates. Although covariates such as C-reactive protein and albumin are not directly incorporated, the model includes proxy indicators of inflammatory burden, such as malaria incidence, HIV prevalence, and undernutrition metrics to inform hemoglobin shift estimates, minimizing the overestimation of dietary contributions in populations with high inflammatory burdens.

### Estimating anemia burden

We estimated the prevalence of mild, moderate and severe anemia across GBD locations, age groups, sexes and years<sup>30</sup>. Primary inputs for this estimation included mean hemoglobin concentration and its standard deviation<sup>30</sup>. Mean hemoglobin levels were modeled using an ST-GPR model to capture patterns and variations across populations and geographical areas. The ST-GPR model incorporated key covariates, such as age-specific fertility rate, HIV prevalence, child malnutrition indicators (for example, stunting and wasting), malaria incidence, healthcare access and quality index, modern contraceptive prevalence and genetic hemoglobin variants (hemoglobin C and S). These covariates enhanced hemoglobin-level prediction accuracy, particularly in data-limited areas<sup>2</sup>. The standard deviation of hemoglobin concentration was optimized based on the modeled mean and anemia prevalence by severity, allowing for hemoglobin distribution estimates tailored to each demographic group<sup>30</sup>.

### Estimating dietary iron deficiency burden

The estimation of dietary iron deficiency focused on the residual health burden remaining after accounting for explicitly modeled causes of anemia, such as malaria, menstrual disorders, gastrointestinal diseases and chronic kidney disease. The GBD cause attribution model then allocated the residual burden into four categories: dietary iron deficiency, neglected tropical diseases, infectious diseases and hemoglobinopathies or hemolytic anemias<sup>30</sup>. This step ensured that dietary iron deficiency was assessed within the narrower framework of residual causes, thereby excluding explicitly modeled causes of anemia and preventing the misattribution of other anemia etiologies to dietary iron deficiency.

To quantify dietary iron deficiency, 80.5% of the residual anemia burden was specifically attributed to this cause. This attribution was based on a standardized hemoglobin shift of  $4.01 \text{ g l}^{-1}$ , derived from robust evidence from iron supplementation and fortification trials. The remaining 19.5% of the residual burden was proportionally distributed among the other three residual categories to account for contributors not explicitly measured or modeled.

This dynamic approach evaluated dietary iron deficiency contributions across different severity levels, age groups, sexes and geographic regions, reflecting population-specific dietary patterns and regional influences<sup>33</sup>. By focusing on dietary iron deficiency as a preventable condition, the model underscores its distinct role in global health, separate from other anemia-related causes. The findings, summarized in Extended Data Fig. 8 and Supplementary Tables 3–5, highlight dietary iron deficiency as a key target for nutritional interventions, distinct from broader anemia-related etiologies.

### Calculation of YLDs and DALYs

DALYs were computed to quantify the overall burden of dietary iron deficiency. DALYs combine years of life lost due to premature death with YLDs, providing a comprehensive measure of health loss within a population<sup>33</sup>.

YLDs were calculated by multiplying the prevalence of sequelae for each disease and injury—categorized by cause, age, sex, location and year—by their respective disability weights<sup>2</sup>. For dietary iron deficiency, disability weights were assigned based on anemia severity: mild anemia (0.004), moderate anemia (0.052), and severe anemia (0.149)<sup>34</sup>. This method captures the varying impact of dietary iron deficiency across severity levels. Notably, the low disability weight for mild anemia reflects its minimal contribution to overall YLDs compared to moderate and severe cases. This approach captures the varying impact of dietary iron deficiency across severity levels, reflecting both mild anemia's low disability weight and its cumulative impact across large populations. By quantifying preventable cases of dietary iron deficiency, this method ensures consistency with established GBD guidelines while providing actionable insights into targeted interventions.

The YLD and DALY calculations adhered to established GBD guidelines, ensuring consistency across global and regional populations. By summing YLDs attributable to dietary iron deficiency, this approach provides a robust estimate of the health burden directly linked to inadequate dietary iron intake.

### Uncertainty

Uncertainty was incorporated at every stage of the estimation process by random sampling from possible value ranges for each estimate<sup>2</sup>. When results were combined across different categories (for example, SDI, geography or age), each sample was treated independently. The final uncertainty interval for each result was determined using the 2.5th and 97.5th percentiles of the draws. In addition to incorporating random sampling, non-sampling variance was calculated using residuals from hierarchical models across geographic levels. For regions with sparse data, variance adjustments were applied to enhance the robustness of the estimates. Bias adjustments were implemented through network analysis, leveraging indirect comparisons to strengthen estimates where direct data comparisons were unavailable. This comprehensive approach accommodates variability across diverse populations and geographies, improving the reliability of dietary iron deficiency estimates.

### Error variance and uncertainty

Non-sampling variance was incorporated into the estimation process by analyzing weighted residuals from hierarchical models across geographic levels. This step accounts for variance even when sample sizes are small, improving estimate reliability. Error variance in the normal and logit-transformed spaces was calculated for hemoglobin levels, with variance adjustments applied where sample size information was unavailable. Uncertainty intervals for all estimates were derived from 1,000 random samples per estimate, with final 95% uncertainty intervals based on the 2.5th and 97.5th percentiles. This approach ensured robust uncertainty estimates, accommodating variability across age, sex and location.

### Data adjustments—crosswalking

Crosswalking is the process of adjusting data to account for known biases<sup>34</sup>. An observation is considered biased if it consistently differs from the standard GBD definition of the modeled parameter. Examples include measures of disease incidence that are self-reported rather than doctor-diagnosed or diagnostic tests that have lower sensitivity or specificity compared to the gold standard diagnostic method. If the difference between an alternative measurement method and the GBD definition is consistent and systematic, it can be modeled using covariates, allowing us to predict the necessary adjustment for a given alternative or non-standard observation. This process enables GBD models to incorporate data from a wider array of sources.

### Data adjustments—bias adjustment for alternative case definitions and study methods

In GBD 2021, we continued adjusting non-fatal and risk exposure data for different case definitions or study methods, a practice started in

GBD 2019. These adjustments were made before using ST-GPR, ensuring consistent data inputs and converting data for both sexes into male and female equivalents. We identified and compared alternative and reference definitions within and between studies, allowing a 5-year difference in between-study comparisons. We quantified bias by calculating differences between matched pairs of alternative and reference observations, using these as dependent variables in a mixed-effect meta-regression model. Covariates were chosen based on systematic differences, and adjustments were applied even without statistical significance if conceptual bias was likely. An open-source Python package (ihmeuw-msca, 2023) was developed to assist with these adjustments, increasing the variance of non-standard data points and reducing their influence in further modeling.

### Data adjustments—example bias adjustment calculation

To adjust for bias in data sources measuring prevalence with non-standard case definitions, we matched pairs of alternative and reference observations by age, sex, location and time period, calculating logit-scale differences to account for bias. If values were zero, data were aggregated across age groups until they became non-zero. Standard errors were computed using the delta method. Differences were modeled in a mixed-effects meta-regression with age and sex as covariates to predict bias adjustments. Adjustments were applied by subtracting the logit-space adjustment factor and using the inverse logit transformation. Uncertainty included the original observation's uncertainty, the predicted adjustment's posterior distribution and random intercepts from the model, with variances summed and transformed back to natural units.

### Data adjustments—network analysis

When multiple alternative case definitions or study methods were present, we used network analysis to utilize additional information from indirect comparisons. For example, if A is the reference and B and C are alternatives, a direct comparison is C versus A, whereas an indirect comparison combines A versus B and B versus C. This method enhances estimates by including more data. Implementing network analysis involves constructing a design matrix for the mixed-effects meta-regression model. If case definitions have subcomponents (for example, symptoms and recall periods), sparse data can make direct and indirect comparisons difficult. In such cases, we assume multiplicative effects across dimensions and use dummy variables to encode these effects. The open-source Python package (ihmeuw-msca, 2023) facilitates this process by automating the design matrix creation and accommodating multiple alternative definitions and covariates.

### Data adjustments—elevation adjustment

Hemoglobin concentration exhibits a positive correlation with elevation, representing a physiological adaptation to decreased ambient oxygen levels, thereby ensuring adequate oxygen delivery throughout the body. Below 1,000 m, the impact on hemoglobin appears negligible. However, previous research suggests an exponential relationship between elevation and hemoglobin levels, as reflected in the WHO-recommended formula for hemoglobin adjustment:

$$\Delta\text{Hb} = -0.32 \times (\text{elevation in meters} \times 0.0033) + 0.22 \times (\text{elevation in meters} \times 0.0033)^2$$

In this analytical approach, GBD modeling used survey-reported data that were either elevation adjusted or both elevation adjusted and smoking adjusted, without further modification. For individual-level data presenting unadjusted hemoglobin values but including elevation information, the equation was applied to make necessary adjustments. Further studies are necessary to test alternative elevation adjustment methodologies. In the absence of smoking-adjusted data, no additional modifications were implemented in the GBD 2021 analysis.

This methodological decision acknowledges the limitations of available data while maintaining consistency across the dataset.

**ST-GPR modeling**

GPR is a flexible statistical method used when sufficient data exists to track complex changes over time. This modeling technique is designed to detect signals in noisy data. It also functions as a powerful tool for interpolating nonlinear trends. Unlike traditional linear models that assume definitive functional forms for underlying trends, GPR considers the trend of interest as following a Gaussian process. This process is characterized by two components: a mean function  $m(\cdot)$  and a covariance function  $\text{Cov}(\cdot)$ . As an example,  $p_{c,a,s,t}$  denotes the observed data in normal, log, or logit space, observed in country  $c$ , for age group  $a$  and sex  $s$  at time  $t$ :

$$(p_{c,a,s,t}) = g_{c,a,s}(t) + \epsilon_{c,a,s,t}$$

where

$$\epsilon_{c,a,s,t} \sim \text{Normal}(0, \sigma_p^2),$$

$$g_{c,a,s}(t) \sim \text{GP}(m_{c,a,s}(t), \text{Cov}(g_{c,a,s}(t))).$$

The calculation of mean and covariance functions,  $m_{c,a,s}(t)$  and  $\text{Cov}(g_{c,a,s}(t))$ , alongside a comprehensive explanation of error variance ( $\sigma_p^2$ ), is outlined in the following section.

**Estimating mean functions**

Mean functions were calculated through a two-phase process.  $m_{c,a,s}(t)$  can be formulated, depending on the exposure transformation, as:

$$\log(p_{c,a,s}(t)) = X_{c,a,s}\beta + h(r_{c,a,s,t})$$

$$\text{logit}(p_{c,a,s}(t)) = X_{c,a,s}\beta + h(r_{c,a,s,t})$$

$$p_{c,a,s}(t) = X_{c,a,s}\beta + h(r_{c,a,s,t})$$

where  $X\beta$  represents the aggregate of hierarchical linear mixed-effects regression components, encompassing the intercept and covariate-coefficient products. The models used hierarchical linear mixed-effects regressions with random intercepts at various geographic levels. These random intercepts were applied during fitting but were excluded from predictions. An ensemble model combined these regressions. The latter part of the equation,  $h(r_{c,a,s,t})$ , functions as a residual smoothing mechanism, where  $r_{c,a,s,t}$  stems from the initial ensemble linear model. Although the linear element captures overall exposure trends over time, considerable data variability often remains unaccounted for. To resolve this, a locally weighted polynomial regression function,  $h(r_{c,a,s,t})$ , was applied. This approach systematically estimates residual variability by leveraging patterns across time, age and space—the spatiotemporal aspect of ST-GPR. The time adjustment parameter,  $\lambda$ , aims to borrow strength from adjacent timepoints, recognizing that exposure in a given year correlates strongly with the previous year but less so with earlier years. The age adjustment parameter,  $\omega$ , uses data from proximate age groups. The space adjustment parameter,  $\xi$ , aims to borrow strength across geographic hierarchies. Spatial and temporal weights merge into a unified space-time weight, allowing spatial weight for a specific point  $r_{c,a,s,t}$  to vary based on data availability at each time  $t$  and location-level  $l$  within the hierarchy.

The final weight  $w_{c,a,s,t}$  is assigned to observation  $r_{c,a,s,t}$  relative to a focal observation  $r_{c_0,a_0,s_0,t_0}$ . Initially, a temporal weight  $t.w_{c,a,s,t}$  was created for time-based smoothing, calculated using the scaled temporal distance between observations:

$$t.w_{c,a,s,t} = \frac{1}{e^{\lambda|t-t_0|}}$$

Subsequently, a spatial weight was developed for geographic smoothing. A geospatial relationship was established by categorizing data according to the GBD location hierarchy. The parameter  $\zeta$  functions as a scalar for each data point, reflecting its proximity to the target location:

$$t.w_{c,a,s,t} = \zeta^{|c-c_0|}$$

For example, estimating a country would use the following weighting scheme:

- Country data:  $\zeta^0 = 1$
- Regional data not from the country being estimated:  $\zeta^1$
- Data from other regions in the same super-region:  $\xi^2$
- Global data from other super-regions:  $\zeta^3$

In the spatial weighting framework,  $\zeta$  typically ranges from 0.001 to 0.2. This parameter indicates how much regional data are down-weighted relative to country-specific data for a given estimate. For instance, with  $\zeta = 0.01$  and a data point  $r_{c,a,s,t}$ , another point from the same region but different country would receive  $\frac{1}{100}$  the weight of an in-country data point.

Given a normalization constant,

$$K_i = \sum_{cc} s.w_{c,t} \times t.w_{c,t} + \sum_{cCR} s.w_{c,t} \times t.w_{c,t} + \sum_{ccSR} s.w_{c,t} \times t.w_{c,t}$$

the final space-time weight would then equal

$$w'_{c,a,s,t} = \frac{s.w_{c,t} \times t.w_{c,t}}{K_i}$$

Lastly, a weight  $w''_{c,a,s,t}$  was computed for age smoothing, based on the age difference between two observations. For a point between the age  $a$  of the observation  $r_{c,a,s,t}$  and a focal observation  $r_{c_0,a_0,s_0,t_0}$ , the weight is defined as follows:

$$w''_{c,a,s,t} = \frac{1}{e^{|\omega|a-a_0|}}$$

The final weights are calculated by multiplying space-time weights with age weights and then normalizing to ensure all weights for a specific time period  $t$  sum to 1. A comprehensive derivation of weights for each category, assuming a country-level estimation, is as follows:

1) When the observation  $r_{c,t}$  originates from the identical country  $c_0$  as the focal observation  $r_{c_0,t_0}$ :

$$w_{c,a,s,t} = \frac{(w'_{c,a,s,t} w''_{c,a,s,t})}{\sum_{c=c_0} (w'_{c,a,s,t} w''_{c,a,s,t})} \quad \forall c = c_0$$

2) If observation  $r_{c,t}$  comes from a different country than focal observation  $r_{c_0,t_0}$ , but both belong to the same region  $R$ :

$$w_{c,a,s,t} = \frac{(w'_{c,a,s,t} w''_{c,a,s,t})}{\sum_{c \neq c_0} (w'_{c,a,s,t} w''_{c,a,s,t})} \quad \forall c \neq c_0 \cap R[c] = R[c_0]$$

3) When observation  $r_{c,t}$  is from the same super-region  $SR$  but differs in both country  $c_0$  and region  $R[c_0]$  from the focal observation  $r_{c_0,t_0}$ :

$$w_{c,a,s,t} = \frac{(w'_{c,a,s,t} w''_{c,a,s,t})}{\sum_{c \neq c_0} (w'_{c,a,s,t} w''_{c,a,s,t})} \quad \forall c \neq c_0 \cap R[c] \neq R[c_0] \cap SR[c] = SR[c_0]$$

4) When observation  $r_{c,t}$  originates from a different super-region than focal observation  $r_{c_0,c_0}$  (encompassing all remaining data not yet assigned a weight):

$$w_{c,a,s,t} = \frac{(w'_{c,a,s,t} w''_{c,a,s,t})}{\sum_{c \neq c_0} (w'_{c,a,s,t} w''_{c,a,s,t})}$$

$$\forall c \neq c_0 \cap R[c] \neq R[c_0] \cap SR[c] \neq SR[c_0]$$

The final weights underwent normalization, ensuring that the total sum of weights across age, time and geographic hierarchy for each reference group equaled 1.

**Estimating error variance**

$\sigma_p^2$  denotes error variance in normal or transformed space, encompassing sampling variance of estimates and prediction error from cross-walks. The variance was systematically imputed when data extraction lacked uncertainty measures. For available sample sizes, missing ones were imputed using the 5th percentile of existing samples. Missing variances were calculated as  $\sigma_p^2 = \frac{p(1-p)}{n}$  for proportions or predicted via regression for continuous values. When sample sizes were completely absent, the 95th percentile of available variances at the most granular geographic level served as imputation. For proportions where  $p \times n$  or  $(1-p) \times n$  is less than 20, the variance was replaced using the Wilson interval score method.

For exposures modeled with log transformation, the error variance was converted to log space using the delta method approximation:

$$\sigma_p^2 \cong \frac{\sigma_{p'}^2}{p_{c,a,s,t}^2}$$

where  $\sigma_{p'}$  denotes the error variance in normal space. For exposures modeled with logit transformation, the error variance was converted to logit space using the delta method approximation as follows:

$$\sigma_p^2 \cong \frac{\sigma_{p'}^2}{(p_{c,a,s,t} \times (1-p_{c,a,s,t}))^2}$$

Before applying GPR, an estimate of non-sampling variance was incorporated into the error variance. These calculations were performed on normal-space variances. Non-sampling variance was derived from the variance of inverse-variance weighted residuals from space-time estimates at each location-level hierarchy. For levels with fewer than 10 data points, the non-sampling variance was substituted with that from the next higher geographic level containing more than 10 data points.

**Estimating the covariance function**

The final input into GPR is the covariance function, determining trend shape and distribution. The Matern–Euclidian covariance function was selected for its versatility in modeling diverse trends with different smoothness levels. This function is defined as:

$$M(t, t') = \sigma^2 \frac{2^{1-\nu}}{\Gamma(\nu)} \left( \frac{d(t, t') \sqrt{2\nu}}{l} \right)^\nu K_\nu \left( \frac{d(t, t') \sqrt{2\nu}}{l} \right)$$

where  $d(\cdot)$  represents a distance measure. The hyperparameters  $\sigma^2$ ,  $\nu$ ,  $l$  and  $K_\nu$  are defined as key aspects of the covariance function. Specifically,  $\sigma^2$  denotes marginal variance;  $\nu$  determines function smoothness and differentiability;  $l$  represents the length scale (indicating when two points become uncorrelated); and  $K_\nu$  is the Bessel function.  $\sigma^2$  was approximated using the normalized median absolute deviation (MADN) ( $r_c$ ) of the difference, which is the normalized absolute deviation of the difference between first-stage linear regression and second-stage spatiotemporal smoothing estimates for each country.

The mean of these country-level MADN estimates was calculated for nations with more than 10 years of data, ensuring robust information on model uncertainty. All models employed a parameter specification of  $\nu = 2$ .

**Covariates**

The modeling approach for dietary iron deficiency employed ST-GPR to estimate mean hemoglobin levels and anemia prevalence across various severities. This multi-step modeling process generated comprehensive estimates for each combination of location, year, age, and sex within the GBD framework. Covariates included factors known to impact hemoglobin levels and anemia prevalence, allowing the model to adjust for demographic, socioeconomic and health-related influences on iron deficiency.

1. Step 1: Ensemble linear mixed-effects regression. The initial phase used ensemble linear mixed-effects regression to account for potential predictive covariates drawn from the GBD study database. These covariates were tested in various combinations with nested random intercepts applied at different geographic levels. This approach allowed the model to capture hierarchical variability by location, improving accuracy in regions with sparse data. The covariate models with the lowest out-of-sample root mean squared error (RMSE) were selected and averaged to produce baseline estimates.
2. Step 2: Spatiotemporal smoothing of residuals. After the initial regression, residuals (differences between the observed data points and the model estimates) were smoothed over space, age and time. This spatiotemporal smoothing process enhanced the precision of the initial estimates by utilizing patterns in adjacent time points, neighboring age groups and nearby geographical areas. This stage drew strength across data points, filling gaps and reducing variability where data coverage was limited.
3. Step 3: Gaussian process regression refinement. In the final step, GPR was applied to refine the smoothed estimates further. GPR is particularly effective for tracking complex, nonlinear trends over time, as it adapts to regional data patterns without imposing a rigid trend form. This refinement produced robust estimates of mean hemoglobin and anemia prevalence across diverse population groups and settings.

Throughout the modeling process, mean hemoglobin data were log transformed, whereas anemia prevalence data were logit transformed to enhance normality and stabilize variance in the data. Covariates were carefully selected based on both their statistical significance and their anticipated influence on hemoglobin levels and anemia prevalence, ensuring that each covariate aligned with established biological or socioeconomic determinants of dietary iron deficiency. Supplementary Table 2 outlines the specific covariates used in the model, along with their expected direction of influence on mean hemoglobin and anemia prevalence. Covariate-specific insights include:

1. Age-specific fertility rate and HIV prevalence are associated with increased anemia prevalence and decreased hemoglobin levels due to higher physiological demands and the immune impact of HIV, respectively.
2. Child underweight and wasting indicate malnutrition, impacting growth and increasing susceptibility to iron deficiency.
3. Malaria incidence contributes to anemia by causing hemolysis, which disrupts iron homeostasis, and hemoglobin C and S traits represent genetic factors impacting hemoglobin levels, especially in regions with high sickle cell prevalence.
4. SDI serves as a broad indicator of socioeconomic development, where higher SDI is typically linked to improved nutrition, higher hemoglobin levels and lower anemia prevalence.

5. HAQI and modern contraception prevalence reflect access to health services and reproductive care, which impact maternal anemia risk.

This carefully curated set of covariates allowed the ST-GPR model to differentiate trends in dietary iron deficiency prevalence from changes in broader anemia prevalence driven by other health or environmental factors. This robust covariate inclusion strategy supports the validity and applicability of the model's predictions across different demographic and geographic contexts.

### Prediction using GPR

Integration over  $g_{c,a,t}(t_*)$  was performed to predict a complete time series for country  $c$ , age  $a$ , sex  $s$  and prediction time  $t_*$  as follows:

$$p_{c,a,s}(t_*) \sim N(m_{c,a,s,t}(t_*), \sigma_p^2 I + \text{Cov}(g_{c,a,s,t}(t_*)))$$

For each country and indicator, 1,000 random samples were drawn from the specified distributions. The final estimated mean for each country was calculated as the average of these draws. Additionally, 95% uncertainty intervals were determined using the 2.5th and 97.5th percentiles of the sample distribution.

### Subnational scaling and aggregation

Internal consistency between national and subnational estimates was maintained through two methods, depending on data coverage. When national data coverage surpassed subnational coverage, subnational estimates were adjusted using population-weighted scaling to align with national estimates. When subnational data coverage was superior, national estimates were derived using population-weighted aggregation of subnational data. This approach was applied across age, sex and time dimensions for each country and risk factor. Additionally, scaling could be performed in logit space, ensuring that subnational proportion estimates remained below 1 after adjustment to align with national levels.

### Reporting summary

Further information on research design is available in the Nature Portfolio Reporting Summary linked to this article.

### Data availability

The findings from this study were produced using data available in public online repositories or in the published literature; data that are publicly available upon reasonable request from the data provider; and data that are not publicly available due to restrictions by the data provider and that were used under license for the current study. Details on data sources can be found on the Global Health Data Exchange website, including information about the data provider and links to where the data can be accessed or requested (where available). Citations and metadata for all input sources used in this analysis are available for download at <https://ghdx.healthdata.org/gbd-2021/sources> (to access all sources, select non-fatal health outcomes as the component and dietary iron deficiency as the cause).

### Code availability

Our study follows the Guidelines for Accurate and Transparent Health Estimate Reporting (GATHER; Supplementary Table 1). All code used for this analysis is publicly available online at [https://github.com/ihmeuw/anemia\\_gbd2021](https://github.com/ihmeuw/anemia_gbd2021).

### References

31. Safiri, S. et al. Burden of anemia and its underlying causes in 204 countries and territories, 1990–2019: results from the Global Burden of Disease Study 2019. *J. Hematol. Oncol.* **14**, 185 (2021).
32. Stevens, G. A. et al. Guidelines for Accurate and Transparent Health Estimates Reporting: the GATHER statement. *Lancet* **388**, e19–e23 (2016).

33. Kassebaum, N. J. et al. A systematic analysis of global anemia burden from 1990 to 2010. *Blood* **123**, 615–624 (2014).
34. Young, M. F., Luo, H. & Suchdev, P. S. The challenge of defining the global burden of iron deficiency anaemia. *Lancet Haematol.* **10**, e702–e704 (2023).

### Acknowledgements

This study was funded by the Bill and Melinda Gates Foundation, the Australian National Health and Medical Research Council and the Queensland Department of Health, Australia. This work was supported by the Yonsei Fellowship, funded by Lee Youn Jae (J.I.S.). This research was supported by the Ministry of Science and ICT (MIST), Korea, under the Information Technology Research Center (ITRC) support program (IITP-2024-RS-2024-00438239 to D.K.Y.), supervised by the Institute for Information & Communications Technology Planning & Evaluation (IITP). The funders of the study had no role in study design, data collection, data analysis, data interpretation or writing of the report. All authors had full access to the study data and had final responsibility for the decision to submit for publication.

### Author contributions

Conceptualization and design: S.L., Y.S., J.H., M.S.K., J.I.S., D.K.Y. and N.J.K.; methodology: S.L., Y.S., J.H., M.S.K., J.I.S., D.K.Y. and N.J.K.; data acquisition: S.L., Y.S., J.H., M.S.K., J.I.S., D.K.Y. and N.J.K.; statistical analysis and data curation: S.L., Y.S., J.H., M.S.K., J.I.S., D.K.Y. and N.J.K.; validation: S.L., Y.S., J.H., M.S.K., J.I.S., D.K.Y. and N.J.K.; data interpretation: S.L., Y.S., J.H., M.S.K., J.I.S., D.K.Y. and N.J.K.; visualization: S.L., Y.S., J.H. and M.S.K.; managing the estimation or publications process: N.J.K., M.S.K. and J.I.S.; writing—original draft preparation: S.L., Y.S., J.H. and M.S.K.; writing—review and editing: all authors provided critical revision to the paper; supervision: J.I.S., D.K.Y. and N.J.K.; project administration: J.I.S., D.K.Y. and N.J.K.; funding acquisition: J.I.S., D.K.Y. and N.J.K. N.J.K. is the senior author. Contributions by the GBD 2021 Dietary Iron Deficiency Collaborators are described in Supplementary Note 1.

### Competing interests

N.J.K. reports grants or contracts from the Bill & Melinda Gates Foundations as well as grant funding for anemia-related research. N.J.K. also reports payment or honoraria for lectures, presentations, speakers' bureaus, manuscript writing or educational events from Bristol Myers Squibb (presentation on GBD 2021 findings for anemia and dietary iron deficiency), outside the submitted work. Competing interests for the GBD 2021 Dietary Iron Deficiency Collaborators are listed in Supplementary Note 2.

### Additional information


**Extended data** is available for this paper at <https://doi.org/10.1038/s41591-025-03624-8>.

**Supplementary information** The online version contains supplementary material available at <https://doi.org/10.1038/s41591-025-03624-8>.

**Correspondence and requests for materials** should be addressed to Jae Il Shin or Dong Keon Yon.


**Peer review information** *Nature Medicine* thanks Jiahong Sun, Jessica Grieger and the other, anonymous, reviewer(s) for their contribution to the peer review of this work. Primary Handling Editor: Ming Yang, in collaboration with the *Nature Medicine* team.

**Reprints and permissions information** is available at [www.nature.com/reprints](http://www.nature.com/reprints).




**Extended Data Fig. 1 | Global rank of dietary iron deficiency in DALYs rate by sex, age group, and SDI.** Global ranks of dietary iron deficiency in DALYs per 100,000 population are displayed by sex, age group, and SDI quintile. The three panels represent data for both sexes (left), males (middle), and females (right).

Rows correspond to age groups (70 years), while columns represent SDI quintiles from low to high. Each cell contains the global rank of dietary iron deficiency for the corresponding subgroup, with color gradients reflecting the rank, ranging from red (higher rank) to green (lower rank).




**Extended Data Fig. 2 | Global distributions for dietary iron deficiency in individuals under 5 years.** Global maps display the prevalence rate and DALY rate (per 100,000 population) for dietary iron deficiency among individuals under 5 years of age, both sexes, in 2021 (**a**). Global maps show the percent change in

prevalence rate and DALY rate (per 100,000 population) for the same population group from 1990 to 2021 (**b**). The maps use color gradients to illustrate geographical variations, with darker shades representing higher values in (**a**) or larger percent changes in (**b**).




**Extended Data Fig. 3 | Global distributions for dietary iron deficiency in individuals aged 5–14 years.** Global maps display the prevalence rate and DALY rate (per 100,000 population) for dietary iron deficiency among individuals aged 5–14 years, both sexes, in 2021 (**a**). Global maps show the percent change in

prevalence rate and DALY rate (per 100,000 population) for the same population group from 1990 to 2021 (**b**). The maps use color gradients to illustrate geographical variations, with darker shades representing higher values in (**a**) or larger percent changes in (**b**).




**Extended Data Fig. 4 | Global distributions for dietary iron deficiency in individuals aged 15–49 years.** Global maps display the prevalence rate and DALY rate (per 100,000 population) for dietary iron deficiency among individuals aged 15–49 years, both sexes, in 2021 (**a**). Global maps show the percent change in

prevalence rate and DALY rate (per 100,000 population) for the same population group from 1990 to 2021 (**b**). The maps use color gradients to illustrate geographical variations, with darker shades representing higher values in (**a**) or larger percent changes in (**b**).




**Extended Data Fig. 5 | Global distributions for dietary iron deficiency in individuals aged 50–69 years.** Global maps display the prevalence rate and DALY rate (per 100,000 population) for dietary iron deficiency among individuals aged 50–69 years, both sexes, in 2021 (**a**). Global maps show the

percent change in prevalence rate and DALY rate (per 100,000 population) for the same population group from 1990 to 2021 (**b**). The maps use color gradients to illustrate geographical variations, with darker shades representing higher values in (**a**) or larger percent changes in (**b**).




**Extended Data Fig. 6 | Global distributions for dietary iron deficiency in individuals aged >70 years.** Global maps display the prevalence rate and DALY rate (per 100,000 population) for dietary iron deficiency among individuals aged > 70 years, both sexes, in 2021 (a). Global maps show the percent change in

prevalence rate and DALY rate (per 100,000 population) for the same population group from 1990 to 2021 (b). The maps use color gradients to illustrate geographical variations, with darker shades representing higher values in (a) or larger percent changes in (b).



**Extended Data Fig. 7 | Global distributions for dietary iron deficiency across all age groups.** Global maps display the all-age prevalence rate and DALY rate (per 100,000 population) for dietary iron deficiency in both sexes in 2021 (**a**). Global maps show the all-age percent change in prevalence rate and DALY rate


(per 100,000 population) for dietary iron deficiency in both sexes from 1990 to 2021 (**b**). The maps use color gradients to illustrate geographical variations, with darker shades representing higher values in (**a**) or larger percent changes in (**b**).



**Extended Data Fig. 8 | Age-standardized prevalence and YLDs due to anemia causes by sex, 2021.**

Age-standardized rates of prevalence and YLDs due to anemia causes are displayed by sex for 2021. Dietary iron deficiency (pink) constitutes the largest segment of both prevalence and YLDs for males and

females, followed by contributions from hemoglobinopathies and hemolytic anemias (orange), other infectious diseases (green), and neglected tropical diseases (purple). The bars, color-coded for specific causes, illustrate the relative proportions attributed to each anemia cause across sexes.



**Extended Data Fig. 9 | Flow chart of dietary iron deficiency.** The methodological framework outlines the estimation process for the burden of dietary iron deficiency, anemia (nonfatal), and their causal attributions. It integrates data inputs such as individual hemoglobin levels, and survey data, utilizing ST-GPR to model hemoglobin mean and standard deviation. Cause-specific prevalence and hemoglobin shifts are derived for mild, moderate, and

severe anemia, incorporating counterfactual distributions to attribute cases specifically to dietary iron deficiency. Disability weights are applied to calculate YLDs and DALYs, while comorbidity corrections refine the estimates. Relative risks, population-attributable fractions, and covariates are combined to provide a structured pathway from raw data to final burden estimation within the GBD framework.

## Reporting Summary

Nature Portfolio wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Portfolio policies, see our [Editorial Policies](#) and the [Editorial Policy Checklist](#).

### Statistics

For all statistical analyses, confirm that the following items are present in the figure legend, table legend, main text, or Methods section.

n/a Confirmed

- The exact sample size ( $n$ ) for each experimental group/condition, given as a discrete number and unit of measurement
- A statement on whether measurements were taken from distinct samples or whether the same sample was measured repeatedly
- The statistical test(s) used AND whether they are one- or two-sided  
*Only common tests should be described solely by name; describe more complex techniques in the Methods section.*
- A description of all covariates tested
- A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons
- A full description of the statistical parameters including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals)
- For null hypothesis testing, the test statistic (e.g.  $F$ ,  $t$ ,  $r$ ) with confidence intervals, effect sizes, degrees of freedom and  $P$  value noted  
*Give  $P$  values as exact values whenever suitable.*
- For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings
- For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes
- Estimates of effect sizes (e.g. Cohen's  $d$ , Pearson's  $r$ ), indicating how they were calculated

*Our web collection on [statistics for biologists](#) contains articles on many of the points above.*

### Software and code

Policy information about [availability of computer code](#)

Data collection

Data analysis

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors and reviewers. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Portfolio [guidelines for submitting code & software](#) for further information.

### Data

Policy information about [availability of data](#)

All manuscripts must include a [data availability statement](#). This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A description of any restrictions on data availability
- For clinical datasets or third party data, please ensure that the statement adheres to our [policy](#)

No primary data collection was carried out for these analyses. Relevant data from population surveys, published studies, and government reports were extracted by a reviewer using a data collection. The findings of this study are based on data from multiple sources, including public online repositories, data available on request, and restricted data used under license. Comprehensive information on all data sources, including provider details and access instructions where applicable, is

available through the Global Health Data Exchange GBD 2021 website <https://ghdx.healthdata.org/gbd-2021>.

This study complies with the Guidelines for Accurate and Transparent Health Estimates Reporting (GATHER) recommendations (Table S1). All maps presented in this study are generated by the authors; no permissions are required for publication.

## Research involving human participants, their data, or biological material

Policy information about studies with [human participants or human data](#). See also policy information about [sex, gender \(identity/presentation\), and sexual orientation](#) and [race, ethnicity and racism](#).

Reporting on sex and gender	No primary data collection was carried out for this analysis, so the study does not involve human research participants.
Reporting on race, ethnicity, or other socially relevant groupings	Not applicable.
Population characteristics	Not applicable.
Recruitment	Not applicable.
Ethics oversight	This study was approved by the University of Washington IRB Committee.

Note that full information on the approval of the study protocol must also be provided in the manuscript.

## Field-specific reporting

Please select the one below that is the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

Life sciences  Behavioural & social sciences  Ecological, evolutionary & environmental sciences

For a reference copy of the document with all sections, see [nature.com/documents/nr-reporting-summary-flat.pdf](https://www.nature.com/documents/nr-reporting-summary-flat.pdf)

## Life sciences study design

All studies must disclose on these points even when the disclosure is negative.

Sample size	The sample size used in the study model is defined as the population from all locations analyzed from 1990 to 2021. 204 countries and territories, categorized into 21 regions and seven broad super-regions. The population at the state, national, regional, super-regional, and global levels was estimated as part of the Global Burden of Disease Study 2021 and represents the global population based on various demographic characteristics such as age (6-11 months, 12-23 months, 2-4 years, 5-9 years, 10-14 years, and similar increments up to 95 years and older), sex, and regions. Detailed methods are described in <a href="https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(24)00757-8/fulltext">https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(24)00757-8/fulltext</a> .
Data exclusions	We carefully evaluated all data sources for inclusion in our study. Sources were excluded if they lacked necessary survey weight factors or crucial demographic information such as sex or age variables. Additionally, we omitted data deemed unreliable, based on assessments by survey administrators or through our own detailed examination. This careful selection process ensured the quality and completeness of the information used in our analysis.
Replication	This study is an observational study based on several years of survey and report data and is, in principle, replicable. However, due to the time required to extract, process, geo-locate all data, and run the statistical models, an explicit replication analysis was not conducted.
Randomization	Randomization was not relevant to this study. This analysis is an observational mapping study and there were no experimental groups.
Blinding	Blinding was not relevant to this study, as it was an observational study using survey and surveillance data.

## Reporting for specific materials, systems and methods

We require information from authors about some types of materials, experimental systems and methods used in many studies. Here, indicate whether each material, system or method listed is relevant to your study. If you are not sure if a list item applies to your research, read the appropriate section before selecting a response.

## Materials & experimental systems

n/a	Involvement in the study
<input checked="" type="checkbox"/>	<input type="checkbox"/> Antibodies
<input checked="" type="checkbox"/>	<input type="checkbox"/> Eukaryotic cell lines
<input checked="" type="checkbox"/>	<input type="checkbox"/> Palaeontology and archaeology
<input checked="" type="checkbox"/>	<input type="checkbox"/> Animals and other organisms
<input checked="" type="checkbox"/>	<input type="checkbox"/> Clinical data
<input checked="" type="checkbox"/>	<input type="checkbox"/> Dual use research of concern
<input checked="" type="checkbox"/>	<input type="checkbox"/> Plants

## Methods

n/a	Involvement in the study
<input checked="" type="checkbox"/>	<input type="checkbox"/> ChIP-seq
<input checked="" type="checkbox"/>	<input type="checkbox"/> Flow cytometry
<input checked="" type="checkbox"/>	<input type="checkbox"/> MRI-based neuroimaging

## Plants

Seed stocks

Not applicable.

Novel plant genotypes

Not applicable.

Authentication

Not applicable.