

4-1-2025

Characterization of interaction aggregating operators setting interval-valued Pythagorean neutrosophic set

Raed Hatamleh

Ahmed Salem Heilat

M. Palanikumar

Abdallah Al-Husban

Follow this and additional works at: https://digitalrepository.unm.edu/nss_journal

Recommended Citation

Hatamleh, Raed; Ahmed Salem Heilat; M. Palanikumar; and Abdallah Al-Husban. "Characterization of interaction aggregating operators setting interval-valued Pythagorean neutrosophic set." *Neutrosophic Sets and Systems* 81, 1 (2025). https://digitalrepository.unm.edu/nss_journal/vol81/iss1/18

This Article is brought to you for free and open access by UNM Digital Repository. It has been accepted for inclusion in *Neutrosophic Sets and Systems* by an authorized editor of UNM Digital Repository. For more information, please contact disc@unm.edu.



Characterization of interaction aggregating operators setting interval-valued Pythagorean neutrosophic set

Raed Hatamleh¹, Ahmed Salem Heilat², M.Palanikumar^(3,*) and Abdallah Al-Husban^{4,5}

¹ Department of Mathematics, Faculty of Science, Jadara University, P.O. Box 733, Irbid 21110, Jordan; raed@jadara.edu.jo.

² Department of Mathematics, Faculty of Science, Jadara University, P.O. Box 733, Irbid 21110, Jordan; ahmed.heilat@yahoo.com

³ Department of Mathematics, Saveetha School of Engineering, Saveetha Institute of Medical and Technical Sciences, Chennai-602105, India; palanimaths86@gmail.com.

⁴ Department of Mathematics, Faculty of Science and Technology, Irbid National University, P.O. Box: 2600 Irbid, Jordan; dralhoshban@inu.edu.jo.

⁵ Jadara Research Center, Jadara University, Irbid 21110, Jordan.

*Correspondence: palanimaths86@gmail.com;

Abstract. In this work, we present novel techniques for the interval-valued Pythagorean neutrosophic interaction aggregating operator. A hybrid of the neutrosophic set and the interval-valued Pythagorean fuzzy set. The innovative averaging and geometric operations of interval-valued Pythagorean neutrosophic interaction numbers are studied using aggregation operator. The interval-valued Pythagorean neutrosophic interaction is boundedness compatible, idempotent, associative, and commutative. Four new aggregating operators are introduced: IPNI weighted averaging operator, interval-valued Pythagorean neutrosophic interaction weighted geometric operator, generalized interval-valued Pythagorean neutrosophic interaction weighted averaging and generalized interval-valued Pythagorean neutrosophic interaction weighted geometric.

Keywords: weighted averaging, weighted geometric, generalized weighted averaging, generalized weighted geometric.

1. Introduction

The uncertainties led to the development of the fuzzy set (FS) [1], intuitionistic FS (IFS) [2], Pythagorean FS (PFS) [3, 4], neutrosophic set (NS) [5], Fuzzy extension and its application [6–14] and Fermatean FS (FFS) [15]. FS suggests that decision-makers use Zadeh's [1] with the membership degree (MD). Each object has MD $\angle$ and non-membership degree (NMD)

ϑ , and it fulfills $0 \leq \angle + \vartheta \leq 1$, for $\angle, \vartheta \in (0, 1)$. Atanassov developed an IFS idea [2]. PFSs are defined by their MDs and NMDs under the condition that $\angle^2 + \vartheta^2 \leq 1$, which was devised by Yager [3]. IFSs and PFSs are widely utilized and have been explored in a wide range of fields. Cuong and associates [16] by developing the concept of picture FSs (PiFSs). PiFSs has been found to accommodate certain more ambiguity because it is an expanded version of IFSs. For $\angle, \vartheta, \varrho \in (0, 1)$, it is observed in PiFSs that the MD $\angle$, the neutral ϑ , and the NMD ϱ have $0 \leq \angle + \vartheta + \varrho \leq 1$. "Yes," "abstain," "no," and "refusal" are expert opinion messages that will be ensured to be transmitted via the PiFS. Furthermore, it will ensure consistency between the assessment data and the actual decision environment and avoid evaluation information from being left out. Despite the fact that PiFSs have been the subject of several applications and studies, the concept has not been fully explored. Raed Hatamleh et al. discussed the various applications such as Uryson's Operator, Sine-Gordon System and Traces Class Perturbations [17–22]. Shahzaib and colleagues [23] defined the spherical FS (SFS) for certain AOs with MADM. An alternative to $0 \leq \angle + \vartheta + \varrho \leq 1$ is required by the SFS: $0 \leq \angle^2 + \vartheta^2 + \varrho^2 \leq 1$. Linguistic SFS AOs were introduced by Jin et al. [24] and discussed in MADM problems. Rafiq and colleagues introduced SFSs and their applications in DM [25] with the condition that $\angle^2 + \vartheta^2 \geq 1$ and decision-making (DM) are troublesome. The concept of an FFS was first proposed by Senapati and associates [15] in 2019. When $0 \leq \angle^3 + \vartheta^3 \leq 1$, both the MD and NMD have this characteristic. Al-Husban et al. discussed the algebraic structures via complex Fuzzy Hyperring, Fuzzy Soft Groups based on Fuzzy Space and Structures of fibers of groups actions on graphs [26–33].

2. Different AOs for IPNN

Definition 2.1. Suppose that $\mathfrak{A}_1 = \langle (\angle_1^{ta}, \angle_1^{tb}), (\sigma_1^{ia}, \sigma_1^{ib}), (\varrho_1^{fa}, \varrho_1^{fb}) \rangle$ and $\mathfrak{A}_2 = \langle (\angle_2^{ta}, \angle_2^{tb}), (\sigma_2^{ia}, \sigma_2^{ib}), (\varrho_2^{fa}, \varrho_2^{fb}) \rangle$ be the any two IPNNs. Then

$$(1) \mathfrak{A}_1 \vee \mathfrak{A}_2 = \left[\begin{array}{l} \left[\begin{array}{l} \sqrt{(\angle_1^{ta})^2 + (\angle_2^{ta})^2 - (\angle_1^{ta})^2 \cdot (\angle_2^{ta})^2}, \\ \sqrt{(\angle_1^{tb})^2 + (\angle_2^{tb})^2 - (\angle_1^{tb})^2 \cdot (\angle_2^{tb})^2} \end{array} \right] \\ \left[\begin{array}{l} \sqrt{(\sigma_1^{ia})^2 + (\sigma_2^{ia})^2 - (\sigma_1^{ia})^2 \cdot (\sigma_2^{ia})^2}, \\ \sqrt{(\sigma_1^{ib})^2 + (\sigma_2^{ib})^2 - (\sigma_1^{ib})^2 \cdot (\sigma_2^{ib})^2} \end{array} \right] \\ \left[\begin{array}{l} (\varrho_1^{fa})^2 + (\varrho_2^{fa})^2 - (\varrho_1^{fa})^2 \cdot (\varrho_2^{fa})^2 \\ \sqrt{-(\varrho_1^{fa})^2 \cdot (\angle_2^{ta})^2 - (\angle_1^{ta})^2 \cdot (\varrho_2^{fa})^2}, \\ (\varrho_1^{fb})^2 + (\varrho_2^{fb})^2 - (\varrho_1^{fb})^2 \cdot (\varrho_2^{fb})^2 \\ \sqrt{-(\varrho_1^{fb})^2 \cdot (\angle_2^{tb})^2 - (\angle_1^{tb})^2 \cdot (\varrho_2^{fb})^2} \end{array} \right] \end{array} \right]$$

Raed Hatamleh, Ahmed Salem Heilat, M.Palanikumar and Abdallah Al-Husban,
Characterization of interaction aggregating operators setting interval-valued Pythagorean
neutrosophic set

$$\begin{aligned}
(2) \quad \mathfrak{A}_1 \bar{\wedge} \mathfrak{A}_2 &= \left[\begin{array}{c} \left[\begin{array}{c} (\mathcal{A}_1^{ta})^2 + (\mathcal{A}_2^{ta})^2 - (\mathcal{A}_1^{ta})^2 \cdot (\mathcal{A}_2^{ta})^2 \\ -(\mathcal{A}_1^{ta})^2 \cdot (\varrho_2^{fa})^2 - (\varrho_1^{fa})^2 \cdot (\mathcal{A}_2^{ta})^2 \\ (\mathcal{A}_1^{tb})^2 + (\mathcal{A}_2^{tb})^2 - (\mathcal{A}_1^{tb})^2 \cdot (\mathcal{A}_2^{tb})^2 \\ -(\mathcal{A}_1^{tb})^2 \cdot (\varrho_2^{fb})^2 - (\varrho_1^{fb})^2 \cdot (\mathcal{A}_2^{tb})^2 \end{array} \right] \\ \left[\begin{array}{c} \sqrt{(\sigma_1^{ia})^2 + (\sigma_2^{ia})^2 - (\sigma_1^{ia})^2 \cdot (\sigma_2^{ia})^2}, \\ \sqrt{(\sigma_1^{ib})^2 + (\sigma_2^{ib})^2 - (\sigma_1^{ib})^2 \cdot (\sigma_2^{ib})^2} \end{array} \right] \\ \left[\begin{array}{c} \sqrt{(\varrho_1^{fa})^2 + (\varrho_2^{fa})^2 - (\varrho_1^{fa})^2 \cdot (\varrho_2^{fa})^2}, \\ \sqrt{(\varrho_1^{fb})^2 + (\varrho_2^{fb})^2 - (\varrho_1^{fb})^2 \cdot (\varrho_2^{fb})^2} \end{array} \right] \end{array} \right] \\
(3) \quad \zeta \cdot \mathfrak{A}_1 &= \left[\begin{array}{c} \left[\begin{array}{c} \sqrt{1 - (1 - (\mathcal{A}_1^{ta})^2)^\zeta}, \sqrt{1 - (1 - (\mathcal{A}_1^{tb})^2)^\zeta} \\ \sqrt{1 - (1 - (\sigma_1^{ia})^2)^\zeta}, \sqrt{1 - (1 - (\sigma_1^{ib})^2)^\zeta} \end{array} \right] \\ \left[\begin{array}{c} \sqrt{(1 - (\mathcal{A}_1^{ta})^2)^\zeta - (1 - (\mathcal{A}_1^{ta} + \varrho_1^{fa})^2)^\zeta}, \\ \sqrt{(1 - (\mathcal{A}_1^{tb})^2)^\zeta - (1 - (\mathcal{A}_1^{tb} + \varrho_1^{fb})^2)^\zeta} \end{array} \right] \\ \left[\begin{array}{c} \sqrt{(1 - (\varrho_1^{fa})^2)^\zeta - (1 - (\mathcal{A}_1^{ta} + \varrho_1^{fa})^2)^\zeta}, \\ \sqrt{(1 - (\varrho_1^{fb})^2)^\zeta - (1 - (\mathcal{A}_1^{tb} + \varrho_1^{fb})^2)^\zeta} \end{array} \right] \end{array} \right] \\
(4) \quad \mathfrak{A}_1^\zeta &= \left[\begin{array}{c} \left[\begin{array}{c} \sqrt{(1 - (\varrho_1^{fa})^2)^\zeta - (1 - (\mathcal{A}_1^{ta} + \varrho_1^{fa})^2)^\zeta}, \\ \sqrt{(1 - (\varrho_1^{fb})^2)^\zeta - (1 - (\mathcal{A}_1^{tb} + \varrho_1^{fb})^2)^\zeta} \end{array} \right] \\ \left[\begin{array}{c} \sqrt{1 - (1 - (\sigma_1^{ia})^2)^\zeta}, \sqrt{1 - (1 - (\sigma_1^{ib})^2)^\zeta}, \\ \sqrt{1 - (1 - (\varrho_1^{fa})^2)^\zeta}, \sqrt{1 - (1 - (\varrho_1^{fb})^2)^\zeta} \end{array} \right] \end{array} \right]
\end{aligned}$$

2.1. IPNIWAO

Definition 2.2. Let $\mathfrak{A}_\partial = \langle (\mathcal{A}_\partial^{ta}, \mathcal{A}_\partial^{tb}), (\sigma_\partial^{ia}, \sigma_\partial^{ib}), (\varrho_\partial^{fa}, \varrho_\partial^{fb}) \rangle$ be the IPNNs, $\partial = 1, 2, \dots, \infty$, $\mathfrak{A}_\partial$ be the weight of $\mathfrak{A}_\partial$ and $\mathfrak{A}_\partial \geq 0, \diamond_{\partial=1}^\infty \mathfrak{A}_\partial = 1$. Then the IPNIWAO $(\mathfrak{A}_1, \mathfrak{A}_2, \dots, \mathfrak{A}_\infty) = \diamond_{\partial=1}^\infty \mathfrak{A}_\partial$.

Theorem 2.3. Let $\mathfrak{A}_\partial = \langle (\mathcal{A}_\partial^{ta}, \mathcal{A}_\partial^{tb}), (\sigma_\partial^{ia}, \sigma_\partial^{ib}), (\varrho_\partial^{fa}, \varrho_\partial^{fb}) \rangle$ be the IPNNs, $\partial = 1, 2, \dots, \infty$.

$$\text{Then, IPNIWAO}(\mathfrak{A}_1, \mathfrak{A}_2, \dots, \mathfrak{A}_\infty) = \left[\begin{array}{c} \left[\begin{array}{c} \sqrt{1 - \square_{\partial=1}^\infty (1 - (\mathcal{A}_\partial^{ta})^2)^{\mathfrak{A}_\partial}}, \sqrt{1 - \square_{\partial=1}^\infty (1 - (\mathcal{A}_\partial^{tb})^2)^{\mathfrak{A}_\partial}} \\ \sqrt{1 - \square_{\partial=1}^\infty (1 - (\sigma_\partial^{ia})^2)^{\mathfrak{A}_\partial}}, \sqrt{1 - \square_{\partial=1}^\infty (1 - (\sigma_\partial^{ib})^2)^{\mathfrak{A}_\partial}} \end{array} \right] \\ \left[\begin{array}{c} \sqrt{\square_{\partial=1}^\infty (1 - (\mathcal{A}_\partial^{ta})^2)^{\mathfrak{A}_\partial} - \square_{\partial=1}^\infty (1 - (\mathcal{A}_\partial^{ta} + \varrho_\partial^{fa})^2)^{\mathfrak{A}_\partial}}, \\ \sqrt{\square_{\partial=1}^\infty (1 - (\mathcal{A}_\partial^{tb})^2)^{\mathfrak{A}_\partial} - \square_{\partial=1}^\infty (1 - (\mathcal{A}_\partial^{tb} + \varrho_\partial^{fb})^2)^{\mathfrak{A}_\partial}} \end{array} \right] \end{array} \right].$$

Raed Hatamleh, Ahmed Salem Heilat, M.Palanikumar and Abdallah Al-Husban,
Characterization of interaction aggregating operators setting interval-valued Pythagorean
neutrosophic set

Proof. If $\partial = 2$, $\text{IPNIWAO}(\lrcorner_1, \lrcorner_2) = \mathfrak{A}_1 \lrcorner_1 \vee \mathfrak{A}_2 \lrcorner_2$,

where,

$$\mathfrak{A}_1 \lrcorner_1 = \left[\begin{array}{l} \left[\sqrt{1 - \left(1 - (\angle_1^{ta})^2\right)^{\mathfrak{A}_1}}, \sqrt{1 - \left(1 - (\angle_1^{tb})^2\right)^{\mathfrak{A}_1}} \right] \\ \left[\sqrt{1 - \left(1 - (\sigma_1^{ia})^2\right)^{\mathfrak{A}_1}}, \sqrt{1 - \left(1 - (\sigma_1^{ib})^2\right)^{\mathfrak{A}_1}} \right] \\ \left[\sqrt{\left(1 - (\angle_1^{ta})^2\right)^{\mathfrak{A}_1} - \left(1 - (\angle_1^{ta} + \varrho_1^{fa})^2\right)^{\mathfrak{A}_1}}, \right. \\ \left. \sqrt{\left(1 - (\angle_1^{tb})^2\right)^{\mathfrak{A}_1} - \left(1 - (\angle_1^{tb} + \varrho_1^{fb})^2\right)^{\mathfrak{A}_1}} \right] \end{array} \right]$$

and

$$\mathfrak{A}_2 \lrcorner_2 = \left[\begin{array}{l} \left[\sqrt{1 - \left(1 - (\angle_2^{ta})^2\right)^{\mathfrak{A}_2}}, \sqrt{1 - \left(1 - (\angle_2^{tb})^2\right)^{\mathfrak{A}_2}} \right] \\ \left[\sqrt{1 - \left(1 - (\sigma_2^{ia})^2\right)^{\mathfrak{A}_2}}, \sqrt{1 - \left(1 - (\sigma_2^{ib})^2\right)^{\mathfrak{A}_2}} \right] \\ \left[\sqrt{\left(1 - (\angle_2^{ta})^2\right)^{\mathfrak{A}_2} - \left(1 - (\angle_2^{ta} + \varrho_2^{fa})^2\right)^{\mathfrak{A}_2}}, \right. \\ \left. \sqrt{\left(1 - (\angle_2^{tb})^2\right)^{\mathfrak{A}_2} - \left(1 - (\angle_2^{tb} + \varrho_2^{fb})^2\right)^{\mathfrak{A}_2}} \right] \end{array} \right]$$

We get

$$\mathfrak{A}_1 \lrcorner_1 \vee \mathfrak{A}_2 \lrcorner_2 = \left[\begin{array}{l} \left[\begin{array}{l} \left(1 - \left(1 - (\angle_1^{ta})^2\right)^{\mathfrak{A}_1}\right) + \left(1 - \left(1 - (\angle_2^{ta})^2\right)^{\mathfrak{A}_2}\right) \\ - \left(1 - \left(1 - (\angle_1^{ta})^2\right)^{\mathfrak{A}_1}\right) \cdot \left(1 - \left(1 - (\angle_2^{ta})^2\right)^{\mathfrak{A}_2}\right), \\ \left(1 - \left(1 - (\angle_1^{tb})^2\right)^{\mathfrak{A}_1}\right) + \left(1 - \left(1 - (\angle_2^{tb})^2\right)^{\mathfrak{A}_2}\right) \\ - \left(1 - \left(1 - (\angle_1^{tb})^2\right)^{\mathfrak{A}_1}\right) \cdot \left(1 - \left(1 - (\angle_2^{tb})^2\right)^{\mathfrak{A}_2}\right), \end{array} \right] \\ \left[\begin{array}{l} \left(1 - \left(1 - (\sigma_1^{ia})^2\right)^{\mathfrak{A}_1}\right) + \left(1 - \left(1 - (\sigma_2^{ia})^2\right)^{\mathfrak{A}_2}\right) \\ - \left(1 - \left(1 - (\sigma_1^{ia})^2\right)^{\mathfrak{A}_1}\right) \cdot \left(1 - \left(1 - (\sigma_2^{ia})^2\right)^{\mathfrak{A}_2}\right), \\ \left(1 - \left(1 - (\sigma_1^{ib})^2\right)^{\mathfrak{A}_1}\right) + \left(1 - \left(1 - (\sigma_2^{ib})^2\right)^{\mathfrak{A}_2}\right) \\ - \left(1 - \left(1 - (\sigma_1^{ib})^2\right)^{\mathfrak{A}_1}\right) \cdot \left(1 - \left(1 - (\sigma_2^{ib})^2\right)^{\mathfrak{A}_2}\right), \end{array} \right] \\ \left[\begin{array}{l} \left(1 - \left(1 - \varrho_1^{fa}\right)^2\right)^{\mathfrak{A}_1} + \left(1 - \left(1 - \varrho_2^{fa}\right)^2\right)^{\mathfrak{A}_2} \\ - \left(1 - \left(1 - \varrho_1^{fa}\right)^2\right)^{\mathfrak{A}_1} \cdot \left(1 - \left(1 - \varrho_2^{fa}\right)^2\right)^{\mathfrak{A}_2} \\ - \left(1 - \left(\angle_1^{ta} + \varrho_1^{fa}\right)^2\right)^{\mathfrak{A}_1} \cdot \left(1 - \left(\angle_2^{ta} + \varrho_2^{fa}\right)^2\right)^{\mathfrak{A}_2} \\ \left(1 - \left(1 - \varrho_1^{fb}\right)^2\right)^{\mathfrak{A}_1} + \left(1 - \left(1 - \varrho_2^{fb}\right)^2\right)^{\mathfrak{A}_2} \\ - \left(1 - \left(1 - \varrho_1^{fb}\right)^2\right)^{\mathfrak{A}_1} \cdot \left(1 - \left(1 - \varrho_2^{fb}\right)^2\right)^{\mathfrak{A}_2} \\ - \left(1 - \left(\angle_1^{tb} + \varrho_1^{fb}\right)^2\right)^{\mathfrak{A}_1} \cdot \left(1 - \left(\angle_2^{tb} + \varrho_2^{fb}\right)^2\right)^{\mathfrak{A}_2} \end{array} \right] \end{array} \right]$$

$$= \left[\begin{array}{c} \left[\sqrt{1 - \square_{\partial=1}^2 \left(1 - (\angle_{\partial}^{ta})^2\right)^{3\partial}}, \sqrt{1 - \square_{\partial=1}^2 \left(1 - (\angle_{\partial}^{tb})^2\right)^{3\partial}} \right] \\ \left[\sqrt{1 - \square_{\partial=1}^2 \left(1 - (\sigma_{\partial}^{ia})^2\right)^{3\partial}}, \sqrt{1 - \square_{\partial=1}^2 \left(1 - (\sigma_{\partial}^{ib})^2\right)^{3\partial}} \right] \\ \left[\sqrt{\square_{\partial=1}^2 \left(1 - (\angle_{\partial}^{ta})^2\right)^{3\partial} - \square_{\partial=1}^2 \left(1 - (\angle_{\partial}^{ta} + \varrho_{\partial}^{fa})^2\right)^{3\partial}}, \right. \\ \left. \sqrt{\square_{\partial=1}^2 \left(1 - (\angle_{\partial}^{tb})^2\right)^{3\partial} - \square_{\partial=1}^2 \left(1 - (\angle_{\partial}^{tb} + \varrho_{\partial}^{fb})^2\right)^{3\partial}} \right] \end{array} \right].$$

Using induction $\partial \geq 3, IPNIWAO(\downarrow_1, \downarrow_2, \dots, \downarrow_{\alpha})$

$$= \left[\begin{array}{c} \left[\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - (\angle_{\partial}^{ta})^2\right)^{3\partial}}, \sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - (\angle_{\partial}^{tb})^2\right)^{3\partial}} \right] \\ \left[\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - (\sigma_{\partial}^{ia})^2\right)^{3\partial}}, \sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - (\sigma_{\partial}^{ib})^2\right)^{3\partial}} \right] \\ \left[\sqrt{\square_{\partial=1}^{\infty} \left(1 - (\angle_{\partial}^{ta})^2\right)^{3\partial} - \square_{\partial=1}^{\infty} \left(1 - (\angle_{\partial}^{ta} + \varrho_{\partial}^{fa})^2\right)^{3\partial}}, \right. \\ \left. \sqrt{\square_{\partial=1}^{\infty} \left(1 - (\angle_{\partial}^{tb})^2\right)^{3\partial} - \square_{\partial=1}^{\infty} \left(1 - (\angle_{\partial}^{tb} + \varrho_{\partial}^{fb})^2\right)^{3\partial}} \right] \end{array} \right].$$

If $\partial = \alpha + 1$, then $IPNIWAO(\downarrow_1, \downarrow_2, \dots, \downarrow_{\alpha}, \downarrow_{\alpha+1})$

$$= \left[\begin{array}{c} \left[\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - (\angle_{\partial}^{ta})^2\right)^{3\partial} \cdot \left(1 - (\angle_{\alpha+1}^{ta})^2\right)^{3\alpha+1}}, \right. \\ \left. \sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - (\angle_{\partial}^{tb})^2\right)^{3\partial} \cdot \left(1 - (\angle_{\alpha+1}^{tb})^2\right)^{3\alpha+1}} \right] \\ \left[\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - (\sigma_{\partial}^{ia})^2\right)^{3\partial} \cdot \left(1 - (\sigma_{\alpha+1}^{ia})^2\right)^{3\alpha+1}}, \right. \\ \left. \sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - (\sigma_{\partial}^{ib})^2\right)^{3\partial} \cdot \left(1 - (\sigma_{\alpha+1}^{ib})^2\right)^{3\alpha+1}} \right] \\ \left[\sqrt{\left(\square_{\partial=1}^{\infty} \left(1 - (\angle_{\partial}^{ta})^2\right)^{3\partial} - \square_{\partial=1}^{\infty} \left(1 - (\angle_{\partial}^{ta} + \varrho_{\partial}^{fa})^2\right)^{3\partial}\right) \cdot \left(\left((\angle_{\alpha+1}^{ta})^2\right)^{3\alpha+1} - (\angle_{\alpha+1}^{ta} + \varrho_{\alpha+1}^{fa})^2\right)^{3\alpha+1}}, \right. \\ \left. \sqrt{\left(\square_{\partial=1}^{\infty} \left(1 - (\angle_{\partial}^{tb})^2\right)^{3\partial} - \square_{\partial=1}^{\infty} \left(1 - (\angle_{\partial}^{tb} + \varrho_{\partial}^{fb})^2\right)^{3\partial}\right) \cdot \left(\left((\angle_{\alpha+1}^{tb})^2\right)^{3\alpha+1} - (\angle_{\alpha+1}^{tb} + \varrho_{\alpha+1}^{fb})^2\right)^{3\alpha+1}} \right] \end{array} \right] \\ = \left[\begin{array}{c} \left[\sqrt{1 - \square_{\partial=1}^{\alpha+1} \left(1 - (\angle_{\partial}^{ta})^2\right)^{3\partial}}, \sqrt{1 - \square_{\partial=1}^{\alpha+1} \left(1 - (\angle_{\partial}^{tb})^2\right)^{3\partial}} \right] \\ \left[\sqrt{1 - \square_{\partial=1}^{\alpha+1} \left(1 - (\sigma_{\partial}^{ia})^2\right)^{3\partial}}, \sqrt{1 - \square_{\partial=1}^{\alpha+1} \left(1 - (\sigma_{\partial}^{ib})^2\right)^{3\partial}} \right] \\ \left[\sqrt{\square_{\partial=1}^{\alpha+1} \left(1 - (\angle_{\partial}^{ta})^2\right)^{3\partial} - \square_{\partial=1}^{\alpha+1} \left(1 - (\angle_{\partial}^{ta} + \varrho_{\partial}^{fa})^2\right)^{3\partial}}, \right. \\ \left. \sqrt{\square_{\partial=1}^{\alpha+1} \left(1 - (\angle_{\partial}^{tb})^2\right)^{3\partial} - \square_{\partial=1}^{\alpha+1} \left(1 - (\angle_{\partial}^{tb} + \varrho_{\partial}^{fb})^2\right)^{3\partial}} \right] \end{array} \right]$$

Theorem 2.4. If $\downarrow_{\partial} = \langle (\angle_{\partial}^{ta}, \angle_{\partial}^{tb}), (\sigma_{\partial}^{ia}, \sigma_{\partial}^{ib}), (\varrho_{\partial}^{fa}, \varrho_{\partial}^{fb}) \rangle$ be the IPNNs and $\downarrow_{\partial} = \downarrow$ and $\angle_{\partial}^{tb} \cdot \varrho_{\partial}^{fb} = 0$, then the $IPNIWAO(\downarrow_1, \downarrow_2, \dots, \downarrow_{\alpha}) = \downarrow, \partial = 1, 2, \dots, \alpha$.

Raed Hatamleh, Ahmed Salem Heilat, M.Palanikumar and Abdallah Al-Husban,
Characterization of interaction aggregating operators setting interval-valued Pythagorean
neutrosophic set

Proof. Note that, $\langle (\angle_{\partial}^{ta}, \angle_{\partial}^{tb}), (\sigma_{\partial}^{ia}, \sigma_{\partial}^{ib}), (\varrho_{\partial}^{fa}, \varrho_{\partial}^{fb}) \rangle = \langle (\angle^{ta}, \angle^{tb}), (\sigma^{ia}, \sigma^{ib}), (\varrho^{fa}, \varrho^{fb}) \rangle$, $\partial = 1, 2, \dots, \alpha$ and $\diamond_{\partial=1}^{\infty} \vartheta = 1$. We get, $IPNIWAO(\lrcorner_1, \lrcorner_2, \dots, \lrcorner_{\alpha})$

$$\begin{aligned}
 &= \left[\left[\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - (\angle^{ta})^2\right)^{\vartheta}}, \sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - (\angle^{tb})^2\right)^{\vartheta}} \right] \right. \\
 &\quad \left[\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - (\sigma^{ia})^2\right)^{\vartheta}}, \sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - (\sigma^{ib})^2\right)^{\vartheta}} \right] \\
 &\quad \left[\sqrt{\square_{\partial=1}^{\infty} \left(1 - (\angle^{ta})^2\right)^{\vartheta} - \square_{\partial=1}^{\infty} \left(1 - (\angle^{ta} + \varrho^{fa})^2\right)^{\vartheta}}, \right. \\
 &\quad \left. \left. \sqrt{\square_{\partial=1}^{\infty} \left(1 - (\angle^{tb})^2\right)^{\vartheta} - \square_{\partial=1}^{\infty} \left(1 - (\angle^{tb} + \varrho^{fb})^2\right)^{\vartheta}} \right] \right] \\
 &= \left[\left[\sqrt{1 - \left(1 - (\angle^{ta})^2\right)^{\diamond_{\partial=1}^{\infty} \vartheta}}, \sqrt{1 - \left(1 - (\angle^{tb})^2\right)^{\diamond_{\partial=1}^{\infty} \vartheta}} \right] \right. \\
 &\quad \left[\sqrt{1 - \left(1 - (\sigma^{ia})^2\right)^{\diamond_{\partial=1}^{\infty} \vartheta}}, \sqrt{1 - \left(1 - (\sigma^{ib})^2\right)^{\diamond_{\partial=1}^{\infty} \vartheta}} \right] \\
 &\quad \left[\sqrt{\left(1 - (\angle^{ta})^2\right)^{\diamond_{\partial=1}^{\infty} \vartheta} - \left(1 - (\angle^{ta} + \varrho^{fa})^2\right)^{\diamond_{\partial=1}^{\infty} \vartheta}}, \right. \\
 &\quad \left. \left. \sqrt{\left(1 - (\angle^{tb})^2\right)^{\diamond_{\partial=1}^{\infty} \vartheta} - \left(1 - (\angle^{tb} + \varrho^{fb})^2\right)^{\diamond_{\partial=1}^{\infty} \vartheta}} \right] \right] \\
 &= \left[\left[\sqrt{1 - \left(1 - (\angle^{ta})^2\right)}, \sqrt{1 - \left(1 - (\angle^{tb})^2\right)} \right] \right. \\
 &\quad \left[\sqrt{1 - \left(1 - (\sigma^{ia})^2\right)}, \sqrt{1 - \left(1 - (\sigma^{ib})^2\right)} \right] \\
 &\quad \left[\sqrt{\left(1 - (\angle^{ta})^2\right) - \left(1 - (\angle^{ta} + \varrho^{fa})^2\right)}, \right. \\
 &\quad \left. \left. \sqrt{\left(1 - (\angle^{tb})^2\right) - \left(1 - (\angle^{tb} + \varrho^{fb})^2\right)} \right] \right] \\
 &= \left((\angle^{ta}, \angle^{tb}), (\sigma^{ia}, \sigma^{ib}), (\varrho^{fa}, \varrho^{fb}) \right) = \lrcorner
 \end{aligned}$$

2.2. Interaction weighted geometric(IPNIWGO)

Definition 2.5. Let $\lrcorner_{\partial} = \langle (\angle_{\partial}^{ta}, \angle_{\partial}^{tb}), (\sigma_{\partial}^{ia}, \sigma_{\partial}^{ib}), (\varrho_{\partial}^{fa}, \varrho_{\partial}^{fb}) \rangle$ be the IPNNs, ϑ_{∂} be the weight of $\lrcorner_{\partial}$. Then the IPNIWGO $(\lrcorner_1, \lrcorner_2, \dots, \lrcorner_{\alpha}) = \square_{\partial=1}^{\infty} \lrcorner_{\partial}^{\vartheta_{\partial}}$.

Theorem 2.6. If $\lrcorner_{\partial} = \langle (\angle_{\partial}^{ta}, \angle_{\partial}^{tb}), (\sigma_{\partial}^{ia}, \sigma_{\partial}^{ib}), (\varrho_{\partial}^{fa}, \varrho_{\partial}^{fb}) \rangle$ be the IPNNs. Then,

$$IPNIWGO(\lrcorner_1, \lrcorner_2, \dots, \lrcorner_{\alpha}) = \left[\left[\sqrt{\square_{\partial=1}^{\infty} \left(1 - (\varrho_{\partial}^{fa})^2\right)^{\vartheta_{\partial}} - \square_{\partial=1}^{\infty} \left(1 - (\angle_{\partial}^{ta} + \varrho_{\partial}^{fa})^2\right)^{\vartheta_{\partial}}}, \right. \right. \\
 \left. \left[\sqrt{\square_{\partial=1}^{\infty} \left(1 - (\varrho_{\partial}^{fb})^2\right)^{\vartheta_{\partial}} - \square_{\partial=1}^{\infty} \left(1 - (\angle_{\partial}^{tb} + \varrho_{\partial}^{fb})^2\right)^{\vartheta_{\partial}}} \right] \right. \\
 \left[\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - (\sigma_{\partial}^{ia})^2\right)^{\vartheta_{\partial}}}, \sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - (\sigma_{\partial}^{ib})^2\right)^{\vartheta_{\partial}}} \right] \\
 \left. \left[\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - (\varrho_{\partial}^{fa})^2\right)^{\vartheta_{\partial}}}, \sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - (\varrho_{\partial}^{fb})^2\right)^{\vartheta_{\partial}}} \right] \right]$$

Proof. If $\partial = 2$, then $\text{IPNIWGO}(\mathfrak{J}_1, \mathfrak{J}_2) = \mathfrak{J}_1^{\mathfrak{J}_1} \bar{\wedge} \mathfrak{J}_2^{\mathfrak{J}_2}$,

where,

$$\mathfrak{J}_1^{\mathfrak{J}_1} = \begin{bmatrix} \left[\sqrt{\left(1 - (\varrho_1^{fa})^2\right)^{\mathfrak{J}_1}} - \left(1 - (\angle_1^{ta} + \varrho_1^{fa})^2\right)^{\mathfrak{J}_1}, \right. \\ \left. \sqrt{\left(1 - (\varrho_1^{fb})^2\right)^{\mathfrak{J}_1}} - \left(1 - (\angle_1^{tb} + \varrho_1^{fb})^2\right)^{\mathfrak{J}_1} \right] \\ \left[\sqrt{1 - \left(1 - (\sigma_1^{ia})^2\right)^{\mathfrak{J}_1}}, \sqrt{1 - \left(1 - (\sigma_1^{ib})^2\right)^{\mathfrak{J}_1}} \right] \\ \left[\sqrt{1 - \left(1 - (\varrho_1^{fa})^2\right)^{\mathfrak{J}_1}}, \sqrt{1 - \left(1 - (\varrho_1^{fb})^2\right)^{\mathfrak{J}_1}} \right] \end{bmatrix}$$

$$\mathfrak{J}_2^{\mathfrak{J}_2} = \begin{bmatrix} \left[\sqrt{\left(1 - (\varrho_2^{fa})^2\right)^{\mathfrak{J}_2}} - \left(1 - (\angle_2^{ta} + \varrho_2^{fa})^2\right)^{\mathfrak{J}_2}, \right. \\ \left. \sqrt{\left(1 - (\varrho_2^{fb})^2\right)^{\mathfrak{J}_2}} - \left(1 - (\angle_2^{tb} + \varrho_2^{fb})^2\right)^{\mathfrak{J}_2} \right] \\ \left[\sqrt{1 - \left(1 - (\sigma_2^{ia})^2\right)^{\mathfrak{J}_2}}, \sqrt{1 - \left(1 - (\sigma_2^{ib})^2\right)^{\mathfrak{J}_2}} \right] \\ \left[\sqrt{1 - \left(1 - (\varrho_2^{fa})^2\right)^{\mathfrak{J}_2}}, \sqrt{1 - \left(1 - (\varrho_2^{fb})^2\right)^{\mathfrak{J}_2}} \right] \end{bmatrix}$$

We get,

$$\mathfrak{J}_1^{\mathfrak{J}_1} \bar{\wedge} \mathfrak{J}_2^{\mathfrak{J}_2} = \begin{bmatrix} \left[\begin{aligned} &\left(1 - \left(1 - (\varrho_1^{fa})^2\right)^{\mathfrak{J}_1}\right) + \left(1 - \left(1 - (\varrho_2^{fa})^2\right)^{\mathfrak{J}_2}\right) \\ &- \left(1 - \left(1 - (\varrho_1^{fa})^2\right)^{\mathfrak{J}_1}\right) \cdot \left(1 - \left(1 - (\varrho_2^{fa})^2\right)^{\mathfrak{J}_2}\right) \\ &- \left(\left(1 - (\angle_1^{ta} + \varrho_1^{fa})^2\right)^{\mathfrak{J}_1} \cdot \left(1 - (\angle_2^{ta} + \varrho_2^{fa})^2\right)^{\mathfrak{J}_2}\right) \end{aligned} \right] \\ \left[\begin{aligned} &\left(1 - \left(1 - (\varrho_1^{fb})^2\right)^{\mathfrak{J}_1}\right) + \left(1 - \left(1 - (\varrho_2^{fb})^2\right)^{\mathfrak{J}_2}\right) \\ &- \left(1 - \left(1 - (\varrho_1^{fb})^2\right)^{\mathfrak{J}_1}\right) \cdot \left(1 - \left(1 - (\varrho_2^{fb})^2\right)^{\mathfrak{J}_2}\right) \\ &- \left(\left(1 - (\angle_1^{tb} + \varrho_1^{fb})^2\right)^{\mathfrak{J}_1} \cdot \left(1 - (\angle_2^{tb} + \varrho_2^{fb})^2\right)^{\mathfrak{J}_2}\right) \end{aligned} \right] \\ \left[\begin{aligned} &\left(1 - \left(1 - (\sigma_1^{ia})^2\right)^{\mathfrak{J}_1}\right) + \left(1 - \left(1 - (\sigma_2^{ia})^2\right)^{\mathfrak{J}_2}\right) \\ &- \left(1 - \left(1 - (\sigma_1^{ia})^2\right)^{\mathfrak{J}_1}\right) \cdot \left(1 - \left(1 - (\sigma_2^{ia})^2\right)^{\mathfrak{J}_2}\right) \\ &- \left(1 - \left(1 - (\sigma_1^{ib})^2\right)^{\mathfrak{J}_1}\right) + \left(1 - \left(1 - (\sigma_2^{ib})^2\right)^{\mathfrak{J}_2}\right) \\ &- \left(1 - \left(1 - (\sigma_1^{ib})^2\right)^{\mathfrak{J}_1}\right) \cdot \left(1 - \left(1 - (\sigma_2^{ib})^2\right)^{\mathfrak{J}_2}\right) \end{aligned} \right] \\ \left[\begin{aligned} &\left(1 - \left(1 - (\varrho_1^{fa})^2\right)^{\mathfrak{J}_1}\right) + \left(1 - \left(1 - (\varrho_2^{fa})^2\right)^{\mathfrak{J}_2}\right) \\ &- \left(1 - \left(1 - (\varrho_1^{fa})^2\right)^{\mathfrak{J}_1}\right) \cdot \left(1 - \left(1 - (\varrho_2^{fa})^2\right)^{\mathfrak{J}_2}\right), \\ &\left(1 - \left(1 - (\varrho_1^{fb})^2\right)^{\mathfrak{J}_1}\right) + \left(1 - \left(1 - (\varrho_2^{fb})^2\right)^{\mathfrak{J}_2}\right) \\ &- \left(1 - \left(1 - (\varrho_1^{fb})^2\right)^{\mathfrak{J}_1}\right) \cdot \left(1 - \left(1 - (\varrho_2^{fb})^2\right)^{\mathfrak{J}_2}\right) \end{aligned} \right] \end{bmatrix}$$

$$\text{Hence, IPNIWGO}(\mathfrak{J}_1, \mathfrak{J}_2) = \left[\begin{array}{c} \left[\sqrt{\square_{\partial=1}^2 \left(1 - (\varrho_{\partial}^{fa})^2\right)^{3\partial}} - \square_{\partial=1}^2 \left(1 - (\angle_{\partial}^{ta} + \varrho_{\partial}^{fa})^2\right)^{3\partial}}, \right. \\ \left. \sqrt{\square_{\partial=1}^2 \left(1 - (\varrho_{\partial}^{fb})^2\right)^{3\partial}} - \square_{\partial=1}^2 \left(1 - (\angle_{\partial}^{tb} + \varrho_{\partial}^{fb})^2\right)^{3\partial}, \right. \\ \left. \sqrt{1 - \square_{\partial=1}^2 \left(1 - (\sigma_{\partial}^{ia})^2\right)^{3\partial}}, \sqrt{1 - \square_{\partial=1}^2 \left(1 - (\sigma_{\partial}^{ib})^2\right)^{3\partial}} \right. \\ \left. \sqrt{1 - \square_{\partial=1}^2 \left(1 - (\varrho_{\partial}^{fa})^2\right)^{3\partial}}, \sqrt{1 - \square_{\partial=1}^2 \left(1 - (\varrho_{\partial}^{fb})^2\right)^{3\partial}} \right] \end{array} \right]$$

$$\text{IPNIWGO}(\mathfrak{J}_1, \mathfrak{J}_2, \dots, \mathfrak{J}_{\infty}) = \left[\begin{array}{c} \left[\sqrt{\square_{\partial=1}^{\infty} \left(1 - (\varrho_{\partial}^{fa})^2\right)^{3\partial}} - \square_{\partial=1}^{\infty} \left(1 - (\angle_{\partial}^{ta} + \varrho_{\partial}^{fa})^2\right)^{3\partial}, \right. \\ \left. \sqrt{\square_{\partial=1}^{\infty} \left(1 - (\varrho_{\partial}^{fb})^2\right)^{3\partial}} - \square_{\partial=1}^{\infty} \left(1 - (\angle_{\partial}^{tb} + \varrho_{\partial}^{fb})^2\right)^{3\partial} \right. \\ \left. \sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - (\sigma_{\partial}^{ia})^2\right)^{3\partial}}, \sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - (\sigma_{\partial}^{ib})^2\right)^{3\partial}} \right. \\ \left. \sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - (\varrho_{\partial}^{fa})^2\right)^{3\partial}}, \sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - (\varrho_{\partial}^{fb})^2\right)^{3\partial}} \right] \end{array} \right]$$

If $\partial = \infty + 1$, then $\text{IPNIWGO}(\mathfrak{J}_1, \dots, \mathfrak{J}_{\infty}, \mathfrak{J}_{\infty+1})$

$$= \left[\begin{array}{c} \left[\begin{array}{c} \diamond_{\partial=1}^{\infty} \left(1 - \left(1 - (\varrho_{\partial}^{fa})^2\right)^{3\partial}\right) + \left(1 - \left(1 - (\varrho_{\infty+1}^{fa})^2\right)^{3\infty+1}\right) \\ - \square_{\partial=1}^{\infty} \left(1 - \left(1 - (\varrho_{\partial}^{fa})^2\right)^{3\partial}\right) \cdot \left(1 - \left(1 - (\varrho_{\infty+1}^{fa})^2\right)^{3\infty+1}\right) \\ - \square_{\partial=1}^{\infty} \left(1 - (\angle_{\partial}^{ta} + \varrho_{\partial}^{fa})^2\right)^{3\partial} \cdot \left(1 - (\angle_{\infty+1}^{ta} + \varrho_{\infty+1}^{fa})^2\right)^{3\infty+1} \end{array} \right. \\ \left[\begin{array}{c} \diamond_{\partial=1}^{\infty} \left(1 - \left(1 - (\varrho_{\partial}^{fb})^2\right)^{3\partial}\right) + \left(1 - \left(1 - (\varrho_{\infty+1}^{fb})^2\right)^{3\infty+1}\right) \\ - \square_{\partial=1}^{\infty} \left(1 - \left(1 - (\varrho_{\partial}^{fb})^2\right)^{3\partial}\right) \cdot \left(1 - \left(1 - (\varrho_{\infty+1}^{fb})^2\right)^{3\infty+1}\right) \\ - \square_{\partial=1}^{\infty} \left(1 - (\angle_{\partial}^{tb} + \varrho_{\partial}^{fb})^2\right)^{3\partial} \cdot \left(1 - (\angle_{\infty+1}^{tb} + \varrho_{\infty+1}^{fb})^2\right)^{3\infty+1} \end{array} \right. \\ \left[\begin{array}{c} \diamond_{\partial=1}^{\infty} \left(1 - \left(1 - (\sigma_{\partial}^{ia})^2\right)^{3\partial}\right) + \left(1 - \left(1 - (\sigma_{\infty+1}^{ia})^2\right)^{3\infty+1}\right) \\ - \square_{\partial=1}^{\infty} \left(1 - \left(1 - (\sigma_{\partial}^{ia})^2\right)^{3\partial}\right) \cdot \left(1 - \left(1 - (\sigma_{\infty+1}^{ia})^2\right)^{3\infty+1}\right) \end{array} \right. \\ \left[\begin{array}{c} \diamond_{\partial=1}^{\infty} \left(1 - \left(1 - (\sigma_{\partial}^{ib})^2\right)^{3\partial}\right) + \left(1 - \left(1 - (\sigma_{\infty+1}^{ib})^2\right)^{3\infty+1}\right) \\ - \square_{\partial=1}^{\infty} \left(1 - \left(1 - (\sigma_{\partial}^{ib})^2\right)^{3\partial}\right) \cdot \left(1 - \left(1 - (\sigma_{\infty+1}^{ib})^2\right)^{3\infty+1}\right) \end{array} \right. \\ \left[\begin{array}{c} \diamond_{\partial=1}^{\infty} \left(1 - \left(1 - (\varrho_{\partial}^{fa})^2\right)^{3\partial}\right) + \left(1 - \left(1 - (\varrho_{\infty+1}^{fa})^2\right)^{3\infty+1}\right) \\ - \square_{\partial=1}^{\infty} \left(1 - \left(1 - (\varrho_{\partial}^{fa})^2\right)^{3\partial}\right) \cdot \left(1 - \left(1 - (\varrho_{\infty+1}^{fa})^2\right)^{3\infty+1}\right) \end{array} \right. \\ \left[\begin{array}{c} \diamond_{\partial=1}^{\infty} \left(1 - \left(1 - (\varrho_{\partial}^{fb})^2\right)^{3\partial}\right) + \left(1 - \left(1 - (\varrho_{\infty+1}^{fb})^2\right)^{3\infty+1}\right) \\ - \square_{\partial=1}^{\infty} \left(1 - \left(1 - (\varrho_{\partial}^{fb})^2\right)^{3\partial}\right) \cdot \left(1 - \left(1 - (\varrho_{\infty+1}^{fb})^2\right)^{3\infty+1}\right) \end{array} \right] \end{array} \right]$$

$$\begin{aligned}
&= \left[\left[\sqrt{\frac{\left(\square_{\partial=1}^{\infty} \left(1 - (\varrho_{\partial}^{fa})^2\right)^{3\partial} - \square_{\partial=1}^{\infty} \left(1 - (\angle_{\partial}^{ta} + \varrho_{\partial}^{fa})^2\right)^{3\partial}\right)}{\left(\left((\varrho_{\alpha+1}^{fa})^2\right)^{3\alpha+1} - (\angle_{\alpha+1}^{ta} + \varrho_{\alpha+1}^{fa})^2\right)^{3\alpha+1}}}, \right. \right. \\
&\quad \left. \left[\sqrt{\frac{\left(\square_{\partial=1}^{\infty} \left(1 - (\varrho_{\partial}^{fb})^2\right)^{3\partial} - \square_{\partial=1}^{\infty} \left(1 - (\angle_{\partial}^{tb} + \varrho_{\partial}^{fb})^2\right)^{3\partial}\right)}{\left(\left((\varrho_{\alpha+1}^{fb})^2\right)^{3\alpha+1} - (\angle_{\alpha+1}^{tb} + \varrho_{\alpha+1}^{fb})^2\right)^{3\alpha+1}}}, \right. \right. \\
&\quad \left. \left[\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - (\sigma_{\partial}^{ia})^2\right)^{3\partial} \cdot \left(1 - (\sigma_{\alpha+1}^{ia})^2\right)^{3\alpha+1}}, \right. \right. \\
&\quad \left. \left[\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - (\sigma_{\partial}^{ib})^2\right)^{3\partial} \cdot \left(1 - (\sigma_{\alpha+1}^{ib})^2\right)^{3\alpha+1}}, \right. \right. \\
&\quad \left. \left[\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - (\varrho_{\partial}^{fa})^2\right)^{3\partial} \cdot \left(1 - (\varrho_{\alpha+1}^{fa})^2\right)^{3\alpha+1}}, \right. \right. \\
&\quad \left. \left[\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - (\varrho_{\partial}^{fb})^2\right)^{3\partial} \cdot \left(1 - (\varrho_{\alpha+1}^{fb})^2\right)^{3\alpha+1}} \right] \right] \\
&= \left[\left[\sqrt{\square_{\partial=1}^{\infty+1} \left(1 - (\varrho_{\partial}^{fa})^2\right)^{3\partial} - \square_{\partial=1}^{\infty+1} \left(1 - (\angle_{\partial}^{ta} + \varrho_{\partial}^{fa})^2\right)^{3\partial}}, \right. \right. \\
&\quad \left[\sqrt{\square_{\partial=1}^{\infty+1} \left(1 - (\varrho_{\partial}^{fb})^2\right)^{3\partial} - \square_{\partial=1}^{\infty+1} \left(1 - (\angle_{\partial}^{tb} + \varrho_{\partial}^{fb})^2\right)^{3\partial}} \right] \\
&\quad \left[\sqrt{1 - \square_{\partial=1}^{\infty+1} \left(1 - (\sigma_{\partial}^{ia})^2\right)^{3\partial}}, \sqrt{1 - \square_{\partial=1}^{\infty+1} \left(1 - (\sigma_{\partial}^{ib})^2\right)^{3\partial}} \right] \\
&\quad \left[\sqrt{1 - \square_{\partial=1}^{\infty+1} \left(1 - (\varrho_{\partial}^{fa})^2\right)^{3\partial}}, \sqrt{1 - \square_{\partial=1}^{\infty+1} \left(1 - (\varrho_{\partial}^{fb})^2\right)^{3\partial}} \right] \right]
\end{aligned}$$

Corollary 2.7. Let $\downarrow_{\partial} = \langle (\angle_{\partial}^{ta}, \angle_{\partial}^{tb}), (\sigma_{\partial}^{ia}, \sigma_{\partial}^{ib}), (\varrho_{\partial}^{fa}, \varrho_{\partial}^{fb}) \rangle$ be the IPNNs and all are equal and $\angle^{tb} \cdot \varrho^{fb} = 0$. Then $IPNIWGO(\downarrow_1, \downarrow_2, \dots, \downarrow_{\alpha}) = \downarrow$.

2.3. generalized IPNIWAO (GIPNIWAO)

Definition 2.8. Let $\downarrow_{\partial} = \langle (\angle_{\partial}^{ta}, \angle_{\partial}^{tb}), (\sigma_{\partial}^{ia}, \sigma_{\partial}^{ib}), (\varrho_{\partial}^{fa}, \varrho_{\partial}^{fb}) \rangle$ be the IPNNs, ϑ_{∂} be a weight of $\downarrow_{\partial}$. Then, the GIPNIWAO $(\downarrow_1, \downarrow_2, \dots, \downarrow_{\alpha}) = \left(\diamond_{\partial=1}^{\infty} \vartheta_{\partial} \downarrow_{\partial}^2 \right)^{1/2}$.

Theorem 2.9. Let $\downarrow_{\partial} = \langle (\angle_{\partial}^{ta}, \angle_{\partial}^{tb}), (\sigma_{\partial}^{ia}, \sigma_{\partial}^{ib}), (\varrho_{\partial}^{fa}, \varrho_{\partial}^{fb}) \rangle$ be the IPNNs. Then GIPNIWAO $(\downarrow_1, \downarrow_2, \dots, \downarrow_{\alpha}) =$

$$\left[\left[\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\angle_{\partial}^{ta})^2\right)^2\right)^{3\partial}}, \sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\angle_{\partial}^{tb})^2\right)^2\right)^{3\partial}} \right] \right. \\
\left[\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\sigma_{\partial}^{ia})^2\right)^2\right)^{3\partial}}, \sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\sigma_{\partial}^{ib})^2\right)^2\right)^{3\partial}} \right] \\
\left[\sqrt{\square_{\partial=1}^{\infty} \left(1 - \left((\angle_{\partial}^{ta})^2\right)^2\right)^{3\partial} - \square_{\partial=1}^{\infty} \left(1 - \left((\angle_{\partial}^{ta} + \varrho_{\partial}^{fa})^2\right)^2\right)^{3\partial}}, \right. \\
\left. \left[\sqrt{\square_{\partial=1}^{\infty} \left(1 - \left((\angle_{\partial}^{tb})^2\right)^2\right)^{3\partial} - \square_{\partial=1}^{\infty} \left(1 - \left((\angle_{\partial}^{tb} + \varrho_{\partial}^{fb})^2\right)^2\right)^{3\partial}} \right] \right] \right].$$

Raed Hatamleh, Ahmed Salem Heilat, M.Palanikumar and Abdallah Al-Husban,
Characterization of interaction aggregating operators setting interval-valued Pythagorean
neutrosophic set

Proof. First, we have to prove that

$$\diamond_{\partial=1}^{\infty} \mathfrak{A}_{\partial}^2 = \left[\begin{array}{c} \left[\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\angle_{\partial}^{ta})^2 \right)^2 \right)^{\mathfrak{A}_{\partial}}}, \sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\angle_{\partial}^{tb})^2 \right)^2 \right)^{\mathfrak{A}_{\partial}}} \right] \\ \left[\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\sigma_{\partial}^{ia})^2 \right)^2 \right)^{\mathfrak{A}_{\partial}}}, \sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\sigma_{\partial}^{ib})^2 \right)^2 \right)^{\mathfrak{A}_{\partial}}} \right] \\ \left[\sqrt{\square_{\partial=1}^{\infty} \left(1 - \left((\angle_{\partial}^{ta})^2 \right)^2 \right)^{\mathfrak{A}_{\partial}}} - \square_{\partial=1}^{\infty} \left(1 - \left((\angle_{\partial}^{ta} + \varrho_{\partial}^{fa})^2 \right)^2 \right)^{\mathfrak{A}_{\partial}} \right] \\ \left[\sqrt{\square_{\partial=1}^{\infty} \left(1 - \left((\angle_{\partial}^{tb})^2 \right)^2 \right)^{\mathfrak{A}_{\partial}}} - \square_{\partial=1}^{\infty} \left(1 - \left((\angle_{\partial}^{tb} + \varrho_{\partial}^{fb})^2 \right)^2 \right)^{\mathfrak{A}_{\partial}} \right] \end{array} \right].$$

$$\text{If } \partial = 2, \text{ then } \mathfrak{A}_1^2 = \left[\begin{array}{c} \left[\sqrt{1 - \left(1 - \left((\angle_1^{ta})^2 \right)^2 \right)^{\mathfrak{A}_1}}, \sqrt{1 - \left(1 - \left((\angle_1^{tb})^2 \right)^2 \right)^{\mathfrak{A}_1}} \right] \\ \left[\sqrt{1 - \left(1 - \left((\sigma_1^{ia})^2 \right)^2 \right)^{\mathfrak{A}_1}}, \sqrt{1 - \left(1 - \left((\sigma_1^{ib})^2 \right)^2 \right)^{\mathfrak{A}_1}} \right] \\ \left[\sqrt{\left(1 - \left((\angle_1^{ta})^2 \right)^2 \right)^{\mathfrak{A}_1}} - \left(1 - \left((\angle_1^{ta} + \varrho_1^{fa})^2 \right)^2 \right)^{\mathfrak{A}_1} \right] \\ \left[\sqrt{\left(1 - \left((\angle_1^{tb})^2 \right)^2 \right)^{\mathfrak{A}_1}} - \left(1 - \left((\angle_1^{tb} + \varrho_1^{fb})^2 \right)^2 \right)^{\mathfrak{A}_1} \right] \end{array} \right]$$

and

$$\mathfrak{A}_2^2 = \left[\begin{array}{c} \left[\sqrt{1 - \left(1 - \left((\angle_2^{ta})^2 \right)^2 \right)^{\mathfrak{A}_2}}, \sqrt{1 - \left(1 - \left((\angle_2^{tb})^2 \right)^2 \right)^{\mathfrak{A}_2}} \right] \\ \left[\sqrt{1 - \left(1 - \left((\sigma_2^{ia})^2 \right)^2 \right)^{\mathfrak{A}_2}}, \sqrt{1 - \left(1 - \left((\sigma_2^{ib})^2 \right)^2 \right)^{\mathfrak{A}_2}} \right] \\ \left[\sqrt{\left(1 - \left((\angle_2^{ta})^2 \right)^2 \right)^{\mathfrak{A}_2}} - \left(1 - \left((\angle_2^{ta} + \varrho_2^{fa})^2 \right)^2 \right)^{\mathfrak{A}_2} \right] \\ \left[\sqrt{\left(1 - \left((\angle_2^{tb})^2 \right)^2 \right)^{\mathfrak{A}_2}} - \left(1 - \left((\angle_2^{tb} + \varrho_2^{fb})^2 \right)^2 \right)^{\mathfrak{A}_2} \right] \end{array} \right].$$

We get, $\mathfrak{A}_1 \sqcup_1 \vee \mathfrak{A}_2 \sqcup_2 =$

$$\left[\begin{array}{l} \sqrt{\left(\sqrt{1 - \left(1 - \left((\angle_1^{ta})^2 \right)^2} \right)^{\mathfrak{A}_1}} \right)^2 + \left(\sqrt{1 - \left(1 - \left((\angle_2^{ta})^2 \right)^2} \right)^{\mathfrak{A}_1}} \right)^2,} \\ - \left(\sqrt{1 - \left(1 - \left((\angle_1^{ta})^2 \right)^2} \right)^{\mathfrak{A}_1}} \right)^2 \cdot \left(\sqrt{1 - \left(1 - \left((\angle_2^{ta})^2 \right)^2} \right)^{\mathfrak{A}_1}} \right)^2 \\ \sqrt{\left(\sqrt{1 - \left(1 - \left((\angle_1^{tb})^2 \right)^2} \right)^{\mathfrak{A}_1}} \right)^2 + \left(\sqrt{1 - \left(1 - \left((\angle_2^{tb})^2 \right)^2} \right)^{\mathfrak{A}_1}} \right)^2,} \\ - \left(\sqrt{1 - \left(1 - \left((\angle_1^{tb})^2 \right)^2} \right)^{\mathfrak{A}_1}} \right)^2 \cdot \left(\sqrt{1 - \left(1 - \left((\angle_2^{tb})^2 \right)^2} \right)^{\mathfrak{A}_1}} \right)^2 \\ \sqrt{\left(\sqrt{1 - \left(1 - \left((\sigma_1^{ia})^2 \right)^2} \right)^{\mathfrak{A}_1}} \right)^2 + \left(\sqrt{1 - \left(1 - \left((\sigma_2^{ia})^2 \right)^2} \right)^{\mathfrak{A}_1}} \right)^2,} \\ - \left(\sqrt{1 - \left(1 - \left((\sigma_1^{ia})^2 \right)^2} \right)^{\mathfrak{A}_1}} \right)^2 \cdot \left(\sqrt{1 - \left(1 - \left((\sigma_2^{ia})^2 \right)^2} \right)^{\mathfrak{A}_1}} \right)^2 \\ \sqrt{\left(\sqrt{1 - \left(1 - \left((\sigma_1^{ib})^2 \right)^2} \right)^{\mathfrak{A}_1}} \right)^2 + \left(\sqrt{1 - \left(1 - \left((\sigma_2^{ib})^2 \right)^2} \right)^{\mathfrak{A}_1}} \right)^2,} \\ - \left(\sqrt{1 - \left(1 - \left((\sigma_1^{ib})^2 \right)^2} \right)^{\mathfrak{A}_1}} \right)^2 \cdot \left(\sqrt{1 - \left(1 - \left((\sigma_2^{ib})^2 \right)^2} \right)^{\mathfrak{A}_1}} \right)^2 \\ \sqrt{\left(\sqrt{1 - \left(1 - \left((\varrho_1^{fa})^2 \right)^2} \right)^{\mathfrak{A}_1}} \right)^2 + \left(\sqrt{1 - \left(1 - \left((\varrho_2^{fa})^2 \right)^2} \right)^{\mathfrak{A}_1}} \right)^2,} \\ - \left(\sqrt{1 - \left(1 - \left((\varrho_1^{fa})^2 \right)^2} \right)^{\mathfrak{A}_1}} \right)^2 \cdot \left(\sqrt{1 - \left(1 - \left((\varrho_2^{fa})^2 \right)^2} \right)^{\mathfrak{A}_1}} \right)^2 \\ - \left[\left(\sqrt{\left(1 - \left((\angle_1^{ta})^2 \right)^2 \right)^{\mathfrak{A}_1}} - \left(1 - \left((\angle_1^{ta} + \varrho_1^{fa})^2 \right)^2 \right)^{\mathfrak{A}_1} \right)^2 \cdot \right. \\ \left. \left(\sqrt{\left(1 - \left((\angle_2^{ta})^2 \right)^2 \right)^{\mathfrak{A}_1}} - \left(1 - \left((\angle_2^{ta} + \varrho_2^{fa})^2 \right)^2 \right)^{\mathfrak{A}_1} \right)^2 \right] \\ \sqrt{\left(\sqrt{1 - \left(1 - \left((\varrho_1^{fb})^2 \right)^2} \right)^{\mathfrak{A}_1}} \right)^2 + \left(\sqrt{1 - \left(1 - \left((\varrho_2^{fb})^2 \right)^2} \right)^{\mathfrak{A}_1}} \right)^2,} \\ - \left(\sqrt{1 - \left(1 - \left((\varrho_1^{fb})^2 \right)^2} \right)^{\mathfrak{A}_1}} \right)^2 \cdot \left(\sqrt{1 - \left(1 - \left((\varrho_2^{fb})^2 \right)^2} \right)^{\mathfrak{A}_1}} \right)^2 \\ - \left[\left(\sqrt{\left(1 - \left((\angle_1^{tb})^2 \right)^2 \right)^{\mathfrak{A}_1}} - \left(1 - \left((\angle_1^{tb} + \varrho_1^{fb})^2 \right)^2 \right)^{\mathfrak{A}_1} \right)^2 \cdot \right. \\ \left. \left(\sqrt{\left(1 - \left((\angle_2^{tb})^2 \right)^2 \right)^{\mathfrak{A}_1}} - \left(1 - \left((\angle_2^{tb} + \varrho_2^{fb})^2 \right)^2 \right)^{\mathfrak{A}_1} \right)^2 \right] \end{array} \right]$$

Raed Hatamleh, Ahmed Salem Heilat, M.Palanikumar and Abdallah Al-Husban,
Characterization of interaction aggregating operators setting interval-valued Pythagorean
neutrosophic set

$$= \left[\begin{array}{c} \left[\sqrt{1 - \square_{\partial=1}^2 \left(1 - \left((\angle_1^{ta})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}}, \sqrt{1 - \square_{\partial=1}^2 \left(1 - \left((\angle_1^{tb})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} \right] \\ \left[\sqrt{1 - \square_{\partial=1}^2 \left(1 - \left((\sigma_1^{ia})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}}, \sqrt{1 - \square_{\partial=1}^2 \left(1 - \left((\sigma_1^{ib})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} \right] \\ \left[\sqrt{\square_{\partial=1}^2 \left(1 - \left((\angle_1^{ta})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} - \square_{\partial=1}^2 \left(1 - \left((\angle_1^{ta} + \varrho_1^{fa})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}} \right] \\ \left[\sqrt{\square_{\partial=1}^2 \left(1 - \left((\angle_1^{tb})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} - \square_{\partial=1}^2 \left(1 - \left((\angle_1^{tb} + \varrho_1^{fb})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}} \right] \end{array} \right]$$

In general,

$$= \left[\begin{array}{c} \left[\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\angle_1^{ta})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}}, \sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\angle_1^{tb})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} \right] \\ \left[\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\sigma_1^{ia})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}}, \sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\sigma_1^{ib})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} \right] \\ \left[\sqrt{\square_{\partial=1}^{\infty} \left(1 - \left((\angle_1^{ta})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} - \square_{\partial=1}^{\infty} \left(1 - \left((\angle_1^{ta} + \varrho_1^{fa})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}} \right] \\ \left[\sqrt{\square_{\partial=1}^{\infty} \left(1 - \left((\angle_1^{tb})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} - \square_{\partial=1}^{\infty} \left(1 - \left((\angle_1^{tb} + \varrho_1^{fb})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}} \right] \end{array} \right].$$

If $\partial = \infty + 1$, then $\diamond_{\partial=1}^{\infty} \mathfrak{z}_{\partial} \mathfrak{z}_{\partial}^2 + \mathfrak{z}_{\infty+1} \mathfrak{z}_{\infty+1}^2 = \diamond_{\partial=1}^{\infty+1} \mathfrak{z}_{\partial} \mathfrak{z}_{\partial}^2$.

Now, $\diamond_{\partial=1}^{\infty} \mathfrak{z}_{\partial} \mathfrak{z}_{\partial}^2 + \mathfrak{z}_{\infty+1} \mathfrak{z}_{\infty+1}^2 = \mathfrak{z}_1 \mathfrak{z}_1^2 \vee \mathfrak{z}_2 \mathfrak{z}_2^2 \vee \dots \vee \mathfrak{z}_{\infty} \mathfrak{z}_{\infty}^2 \vee \mathfrak{z}_{\infty+1} \mathfrak{z}_{\infty+1}^2$

$$\begin{aligned}
&= \left[\sqrt[3]{\frac{\left(\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\angle_{\partial}^{ta})^2 \right)^2} \right)^{3_{\partial}}} \right)^2 + \left(\sqrt{1 - \left(1 - \left((\angle_{\alpha+1}^{ta})^2 \right)^2 \right)^{3_1}} \right)^2}{\left(\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\angle_{\partial}^{ta})^2 \right)^2 \right)^{3_{\partial}}} \right)^2 \cdot \left(\sqrt{1 - \left(1 - \left((\angle_{\alpha+1}^{ta})^2 \right)^2 \right)^{3_1}} \right)^2} \right.} \\
&\quad \sqrt[3]{\frac{\left(\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\angle_{\partial}^{tb})^2 \right)^2 \right)^{3_{\partial}}} \right)^2 + \left(\sqrt{1 - \left(1 - \left((\angle_{\alpha+1}^{tb})^2 \right)^2 \right)^{3_1}} \right)^2}{\left(\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\angle_{\partial}^{tb})^2 \right)^2 \right)^{3_{\partial}}} \right)^2 \cdot \left(\sqrt{1 - \left(1 - \left((\angle_{\alpha+1}^{tb})^2 \right)^2 \right)^{3_1}} \right)^2} \right.} \\
&\quad \sqrt[3]{\frac{\left(\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\sigma_{\partial}^{ia})^2 \right)^2 \right)^{3_{\partial}}} \right)^2 + \left(\sqrt{1 - \left(1 - \left((\sigma_{\alpha+1}^{ia})^2 \right)^2 \right)^{3_1}} \right)^2}{\left(\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\sigma_{\partial}^{ia})^2 \right)^2 \right)^{3_{\partial}}} \right)^2 \cdot \left(\sqrt{1 - \left(1 - \left((\sigma_{\alpha+1}^{ia})^2 \right)^2 \right)^{3_1}} \right)^2} \right.} \\
&\quad \left. \sqrt[3]{\frac{\left(\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\sigma_{\partial}^{ib})^2 \right)^2 \right)^{3_{\partial}}} \right)^2 + \left(\sqrt{1 - \left(1 - \left((\sigma_{\alpha+1}^{ib})^2 \right)^2 \right)^{3_1}} \right)^2}{\left(\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\sigma_{\partial}^{ib})^2 \right)^2 \right)^{3_{\partial}}} \right)^2 \cdot \left(\sqrt{1 - \left(1 - \left((\sigma_{\alpha+1}^{ib})^2 \right)^2 \right)^{3_1}} \right)^2} \right.} \\
&\quad - \left[\frac{\left(\sqrt{\square_{\partial=1}^{\infty} \left(1 - \left((\angle_{\partial}^{ta})^2 \right)^2 \right)^{3_{\partial}}} - \square_{\partial=1}^{\infty} \left(1 - \left((\angle_{\partial}^{ta} + \varrho_{\partial}^{fa})^2 \right)^2 \right)^{3_{\partial}}} \right)^2}{\left(\sqrt{\left(1 - \left((\angle_{\alpha+1}^{ta})^2 \right)^2 \right)^{3_1}} - \left(1 - \left((\angle_{\alpha+1}^{ta} + \varrho_{\alpha+1}^{fa})^2 \right)^2 \right)^{3_1}} \right)^2} \right.} \\
&\quad \left. \frac{\left(\sqrt{\square_{\partial=1}^{\infty} \left(1 - \left((\angle_{\partial}^{tb})^2 \right)^2 \right)^{3_{\partial}}} - \square_{\partial=1}^{\infty} \left(1 - \left((\angle_{\partial}^{tb} + \varrho_{\partial}^{fb})^2 \right)^2 \right)^{3_{\partial}}} \right)^2}{\left(\sqrt{\left(1 - \left((\angle_{\alpha+1}^{tb})^2 \right)^2 \right)^{3_1}} - \left(1 - \left((\angle_{\alpha+1}^{tb} + \varrho_{\alpha+1}^{fb})^2 \right)^2 \right)^{3_1}} \right)^2} \right] \right] \\
&= \left[\left[\sqrt{1 - \square_{\partial=1}^{\infty+1} \left(1 - \left((\angle_1^{ta})^2 \right)^2 \right)^{3_{\partial}}} \right], \left[\sqrt{1 - \square_{\partial=1}^{\infty+1} \left(1 - \left((\angle_1^{tb})^2 \right)^2 \right)^{3_{\partial}}} \right] \right] \\
&\quad \left[\sqrt{1 - \square_{\partial=1}^{\infty+1} \left(1 - \left((\sigma_1^{ia})^2 \right)^2 \right)^{3_{\partial}}} \right], \left[\sqrt{1 - \square_{\partial=1}^{\infty+1} \left(1 - \left((\sigma_1^{ib})^2 \right)^2 \right)^{3_{\partial}}} \right] \right] \\
&\quad \left[\sqrt{\square_{\partial=1}^{\infty+1} \left(1 - \left((\angle_1^{ta})^2 \right)^2 \right)^{3_{\partial}}} - \square_{\partial=1}^{\infty+1} \left(1 - \left((\angle_1^{ta} + \varrho_1^{fa})^2 \right)^2 \right)^{3_{\partial}}} \right], \\
&\quad \left[\sqrt{\square_{\partial=1}^{\infty+1} \left(1 - \left((\angle_1^{tb})^2 \right)^2 \right)^{3_{\partial}}} - \square_{\partial=1}^{\infty+1} \left(1 - \left((\angle_1^{tb} + \varrho_1^{fb})^2 \right)^2 \right)^{3_{\partial}}} \right] \right]
\end{aligned}$$

Raed Hatamleh, Ahmed Salem Heilat, M.Palanikumar and Abdallah Al-Husban,
 Characterization of interaction aggregating operators setting interval-valued Pythagorean
 neutrosophic set

$$\text{and } \diamond_{\partial=1}^{\alpha+1} \left(\begin{matrix} \mathfrak{z}_{\partial} \\ \mathfrak{z}_{\partial}^2 \end{matrix} \right) = \left[\left[\left(\sqrt{1 - \square_{\partial=1}^{\alpha+1} \left(1 - \left((\angle_{\partial}^{ta})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} \right), \left(\sqrt{1 - \square_{\partial=1}^{\alpha+1} \left(1 - \left((\angle_{\partial}^{tb})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} \right) \right] \right. \\ \left. \left[\left(\sqrt{1 - \square_{\partial=1}^{\alpha+1} \left(1 - \left((\sigma_{\partial}^{ia})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} \right), \left(\sqrt{1 - \square_{\partial=1}^{\alpha+1} \left(1 - \left((\sigma_{\partial}^{ib})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} \right) \right] \right. \\ \left. \left[\left(\sqrt{\square_{\partial=1}^{\alpha+1} \left(1 - \left((\angle_{\partial}^{ta})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} - \square_{\partial=1}^{\alpha+1} \left(1 - \left((\angle_{\partial}^{ta} + \varrho_{\partial}^{fa})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} \right), \right. \right. \\ \left. \left. \left(\sqrt{\square_{\partial=1}^{\alpha+1} \left(1 - \left((\angle_{\partial}^{tb})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} - \square_{\partial=1}^{\alpha+1} \left(1 - \left((\angle_{\partial}^{tb} + \varrho_{\partial}^{fb})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} \right) \right] \right].$$

Corollary 2.10. Let $\mathfrak{z}_{\partial} = \langle (\angle_{\partial}^{ta}, \angle_{\partial}^{tb}), (\sigma_{\partial}^{ia}, \sigma_{\partial}^{ib}), (\varrho_{\partial}^{fa}, \varrho_{\partial}^{fb}) \rangle$ be the IPNNs and all are equal. Then GIPNIWAO $(\mathfrak{z}_1, \mathfrak{z}_2, \dots, \mathfrak{z}_{\alpha}) = \mathfrak{z}$.

2.4. Generalized IPNIWGO (GIPNIWGO)

Definition 2.11. Let $\mathfrak{z}_{\partial} = \langle (\angle_{\partial}^{ta}, \angle_{\partial}^{tb}), (\sigma_{\partial}^{ia}, \sigma_{\partial}^{ib}), (\varrho_{\partial}^{fa}, \varrho_{\partial}^{fb}) \rangle$ be the IPNNs and $\mathfrak{z}_{\partial}$ be the weight of $\mathfrak{z}_{\partial}$, where $\partial = 1, 2, \dots, \alpha$. Then, the GIPNIWGO $(\mathfrak{z}_1, \mathfrak{z}_2, \dots, \mathfrak{z}_{\alpha}) = \frac{1}{\zeta} \left(\square_{\partial=1}^{\alpha} (\zeta \mathfrak{z}_{\partial})^{\mathfrak{z}_{\partial}} \right)$.

Theorem 2.12. Let $\mathfrak{z}_{\partial} = \langle (\angle_{\partial}^{ta}, \angle_{\partial}^{tb}), (\sigma_{\partial}^{ia}, \sigma_{\partial}^{ib}), (\varrho_{\partial}^{fa}, \varrho_{\partial}^{fb}) \rangle$ be the collection of IPNNs. Then the GIPNIWGO $(\mathfrak{z}_1, \mathfrak{z}_2, \dots, \mathfrak{z}_{\alpha}) =$

$$\left[\left[\sqrt{\square_{\partial=1}^{\alpha} \left(1 - \left((\varrho_{\partial}^{fa})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} - \square_{\partial=1}^{\alpha} \left(1 - \left((\angle_{\partial}^{ta} + \varrho_{\partial}^{fa})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} \right], \right. \\ \left. \left[\sqrt{\square_{\partial=1}^{\alpha} \left(1 - \left((\varrho_{\partial}^{fb})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} - \square_{\partial=1}^{\alpha} \left(1 - \left((\angle_{\partial}^{tb} + \varrho_{\partial}^{fb})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} \right] \right] \\ \left[\left[\sqrt{1 - \square_{\partial=1}^{\alpha} \left(1 - \left((\sigma_{\partial}^{ia})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} \right], \sqrt{1 - \square_{\partial=1}^{\alpha} \left(1 - \left((\sigma_{\partial}^{ib})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} \right] \right] \\ \left[\left[\sqrt{1 - \square_{\partial=1}^{\alpha} \left(1 - \left((\varrho_{\partial}^{fa})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} \right], \sqrt{1 - \square_{\partial=1}^{\alpha} \left(1 - \left((\varrho_{\partial}^{fb})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} \right] \right]$$

Proof. Using the induction method,

$$\square_{\partial=1}^{\alpha} (\zeta \mathfrak{z}_{\partial})^{\mathfrak{z}_{\partial}} = \left[\left[\sqrt{\square_{\partial=1}^{\alpha} \left(1 - \left((\varrho_{\partial}^{fa})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} - \square_{\partial=1}^{\alpha} \left(1 - \left((\angle_{\partial}^{ta} + \varrho_{\partial}^{fa})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} \right], \right. \\ \left. \left[\sqrt{\square_{\partial=1}^{\alpha} \left(1 - \left((\varrho_{\partial}^{fb})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} - \square_{\partial=1}^{\alpha} \left(1 - \left((\angle_{\partial}^{tb} + \varrho_{\partial}^{fb})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} \right] \right] \\ \left[\left[\sqrt{1 - \square_{\partial=1}^{\alpha} \left(1 - \left((\sigma_{\partial}^{ia})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} \right], \sqrt{1 - \square_{\partial=1}^{\alpha} \left(1 - \left((\sigma_{\partial}^{ib})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} \right] \right] \\ \left[\left[\sqrt{1 - \square_{\partial=1}^{\alpha} \left(1 - \left((\varrho_{\partial}^{fa})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} \right], \sqrt{1 - \square_{\partial=1}^{\alpha} \left(1 - \left((\varrho_{\partial}^{fb})^2 \right)^2 \right)^{\mathfrak{z}_{\partial}}} \right] \right]$$

If $\partial = 2$, then

$$(\zeta \lrcorner_1)^{\mathfrak{P}_1} = \left[\begin{array}{c} \left[\sqrt{\left(1 - \left((\varrho_1^{fa})^2\right)^2\right)^{\mathfrak{P}_1}} - \left(1 - \left((\angle_1^{ta} + \varrho_1^{fa})^2\right)^2\right)^{\mathfrak{P}_1}, \right. \\ \left. \sqrt{\left(1 - \left((\varrho_1^{fb})^2\right)^2\right)^{\mathfrak{P}_1}} - \left(1 - \left((\angle_1^{tb} + \varrho_1^{fb})^2\right)^2\right)^{\mathfrak{P}_1} \right] \\ \left[\sqrt{1 - \left(1 - \left((\sigma_1^{ia})^2\right)^2\right)^{\mathfrak{P}_1}}, \sqrt{1 - \left(1 - \left((\sigma_1^{ib})^2\right)^2\right)^{\mathfrak{P}_1}} \right] \\ \left[\sqrt{1 - \left(1 - \left((\varrho_1^{fa})^2\right)^2\right)^{\mathfrak{P}_1}}, \sqrt{1 - \left(1 - \left((\varrho_1^{fb})^2\right)^2\right)^{\mathfrak{P}_1}} \right] \end{array} \right]$$

and

$$(\zeta \lrcorner_2)^{\mathfrak{P}_2} = \left[\begin{array}{c} \left[\sqrt{\left(1 - \left((\varrho_2^{fa})^2\right)^2\right)^{\mathfrak{P}_1}} - \left(1 - \left((\angle_2^{ta} + \varrho_2^{fa})^2\right)^2\right)^{\mathfrak{P}_1}, \right. \\ \left. \sqrt{\left(1 - \left((\varrho_2^{fb})^2\right)^2\right)^{\mathfrak{P}_1}} - \left(1 - \left((\angle_2^{tb} + \varrho_2^{fb})^2\right)^2\right)^{\mathfrak{P}_1} \right] \\ \left[\sqrt{1 - \left(1 - \left((\sigma_2^{ia})^2\right)^2\right)^{\mathfrak{P}_1}}, \sqrt{1 - \left(1 - \left((\sigma_2^{ib})^2\right)^2\right)^{\mathfrak{P}_1}} \right] \\ \left[\sqrt{1 - \left(1 - \left((\varrho_2^{fa})^2\right)^2\right)^{\mathfrak{P}_1}}, \sqrt{1 - \left(1 - \left((\varrho_2^{fb})^2\right)^2\right)^{\mathfrak{P}_1}} \right] \end{array} \right]$$

We get, $(\zeta \downarrow_1)^{\mathfrak{P}_1} \bar{\wedge} (\zeta \downarrow_2)^{\mathfrak{P}_2}$

$$\begin{aligned}
 & \sqrt{\left[\left(\sqrt{1 - \left(1 - ((\varrho_1^{fa})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 + \left(\sqrt{1 - \left(1 - ((\varrho_2^{fa})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 \right.} \\
 & \quad \left. - \left(\sqrt{1 - \left(1 - ((\varrho_1^{fa})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 \cdot \left(\sqrt{1 - \left(1 - ((\varrho_2^{fa})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 \right.} \\
 & \quad \left. - \left[\left(\sqrt{\left(1 - ((\varrho_1^{fa})^2)^2\right)^{\mathfrak{P}_1} - \left(1 - ((\angle_1^{ta} + \angle_1^{ta})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 \cdot \right. \right. \\
 & \quad \left. \left. - \left(\sqrt{\left(1 - ((\varrho_2^{fa})^2)^2\right)^{\mathfrak{P}_1} - \left(1 - ((\angle_2^{ta} + \varrho_2^{fa})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 \right] \right]^{2'} \\
 & \sqrt{\left[\left(\sqrt{1 - \left(1 - ((\varrho_1^{fb})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 + \left(\sqrt{1 - \left(1 - ((\varrho_2^{fb})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 \right.} \\
 & \quad \left. - \left(\sqrt{1 - \left(1 - ((\varrho_1^{fb})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 \cdot \left(\sqrt{1 - \left(1 - ((\varrho_2^{fb})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 \right.} \\
 & \quad \left. - \left[\left(\sqrt{\left(1 - ((\varrho_1^{fb})^2)^2\right)^{\mathfrak{P}_1} - \left(1 - ((\angle_1^{tb} + \angle_1^{tb})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 \cdot \right. \right. \\
 & \quad \left. \left. - \left(\sqrt{\left(1 - ((\varrho_2^{fb})^2)^2\right)^{\mathfrak{P}_1} - \left(1 - ((\angle_2^{tb} + \varrho_2^{fb})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 \right] \right]^{2'} \\
 = & \sqrt{\left[\left(\sqrt{1 - \left(1 - ((\sigma_1^{ia})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 + \left(\sqrt{1 - \left(1 - ((\sigma_2^{ia})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 \right.} \\
 & \quad \left. - \left(\sqrt{1 - \left(1 - ((\sigma_1^{ia})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 \cdot \left(\sqrt{1 - \left(1 - ((\sigma_2^{ia})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 \right.} \\
 & \quad \left. - \left[\left(\sqrt{\left(1 - ((\sigma_1^{ib})^2)^2\right)^{\mathfrak{P}_1} - \left(1 - ((\sigma_2^{ib})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 \cdot \right. \right. \\
 & \quad \left. \left. - \left(\sqrt{\left(1 - ((\sigma_1^{ib})^2)^2\right)^{\mathfrak{P}_1} - \left(1 - ((\sigma_2^{ib})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 \right] \right]^{2'} \\
 & \sqrt{\left[\left(\sqrt{1 - \left(1 - ((\varrho_1^{fa})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 + \left(\sqrt{1 - \left(1 - ((\varrho_2^{fa})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 \right.} \\
 & \quad \left. - \left(\sqrt{1 - \left(1 - ((\varrho_1^{fa})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 \cdot \left(\sqrt{1 - \left(1 - ((\varrho_2^{fa})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 \right.} \\
 & \quad \left. - \left[\left(\sqrt{\left(1 - ((\varrho_1^{fb})^2)^2\right)^{\mathfrak{P}_1} - \left(1 - ((\varrho_2^{fb})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 \cdot \right. \right. \\
 & \quad \left. \left. - \left(\sqrt{\left(1 - ((\varrho_1^{fb})^2)^2\right)^{\mathfrak{P}_1} - \left(1 - ((\varrho_2^{fb})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 \right] \right]^{2'} \\
 & \sqrt{\left[\left(\sqrt{1 - \left(1 - ((\varrho_1^{fb})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 + \left(\sqrt{1 - \left(1 - ((\varrho_2^{fb})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 \right.} \\
 & \quad \left. - \left(\sqrt{1 - \left(1 - ((\varrho_1^{fb})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 \cdot \left(\sqrt{1 - \left(1 - ((\varrho_2^{fb})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 \right.} \\
 & \quad \left. - \left[\left(\sqrt{\left(1 - ((\varrho_1^{fb})^2)^2\right)^{\mathfrak{P}_1} - \left(1 - ((\varrho_2^{fb})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 \cdot \right. \right. \\
 & \quad \left. \left. - \left(\sqrt{\left(1 - ((\varrho_1^{fb})^2)^2\right)^{\mathfrak{P}_1} - \left(1 - ((\varrho_2^{fb})^2)^2\right)^{\mathfrak{P}_1}} \right)^2 \right] \right]^{2'}
 \end{aligned}$$

$$= \left[\begin{array}{c} \left[\sqrt{\square_{\partial=1}^2 \left(1 - \left((\varrho_{\partial}^{fa})^2 \right)^2 \right)^{3\partial}} - \square_{\partial=1}^2 \left(1 - \left((\angle_{\partial}^{ta} + \varrho_{\partial}^{fa})^2 \right)^2 \right)^{3\partial}}, \right. \\ \left. \sqrt{\square_{\partial=1}^2 \left(1 - \left((\varrho_{\partial}^{fb})^2 \right)^2 \right)^{3\partial}} - \square_{\partial=1}^2 \left(1 - \left((\angle_{\partial}^{tb} + \varrho_{\partial}^{fb})^2 \right)^2 \right)^{3\partial}} \right] \\ \left[\sqrt{1 - \square_{\partial=1}^2 \left(1 - \left((\sigma_{\partial}^{ia})^2 \right)^2 \right)^{3\partial}}, \sqrt{1 - \square_{\partial=1}^2 \left(1 - \left((\sigma_{\partial}^{ib})^2 \right)^2 \right)^{3\partial}} \right] \\ \left[\sqrt{1 - \square_{\partial=1}^2 \left(1 - \left((\varrho_{\partial}^{fa})^2 \right)^2 \right)^{3\partial}}, \sqrt{1 - \square_{\partial=1}^2 \left(1 - \left((\varrho_{\partial}^{fb})^2 \right)^2 \right)^{3\partial}} \right] \end{array} \right]$$

If $\partial = \infty$, then

$$= \left[\begin{array}{c} \left[\sqrt{\square_{\partial=1}^{\infty} \left(1 - \left((\varrho_{\partial}^{fa})^2 \right)^2 \right)^{3\partial}} - \square_{\partial=1}^{\infty} \left(1 - \left((\angle_{\partial}^{ta} + \varrho_{\partial}^{fa})^2 \right)^2 \right)^{3\partial}}, \right. \\ \left. \sqrt{\square_{\partial=1}^{\infty} \left(1 - \left((\varrho_{\partial}^{fb})^2 \right)^2 \right)^{3\partial}} - \square_{\partial=1}^{\infty} \left(1 - \left((\angle_{\partial}^{tb} + \varrho_{\partial}^{fb})^2 \right)^2 \right)^{3\partial}} \right] \\ \left[\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\sigma_{\partial}^{ia})^2 \right)^2 \right)^{3\partial}}, \sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\sigma_{\partial}^{ib})^2 \right)^2 \right)^{3\partial}} \right] \\ \left[\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\varrho_{\partial}^{fa})^2 \right)^2 \right)^{3\partial}}, \sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\varrho_{\partial}^{fb})^2 \right)^2 \right)^{3\partial}} \right] \end{array} \right]$$

If $\partial = \infty + 1$, then $\square_{\partial=1}^{\infty} (\zeta \downarrow_{\partial})^{3\partial} \cdot (\zeta \downarrow_{\infty+1})^{3\infty+1} = \square_{\partial=1}^{\infty+1} (\zeta \downarrow_{\partial})^{3\partial}$.

Now, $\square_{\partial=1}^{\infty} (\zeta \downarrow_{\partial})^{3\partial} \cdot (\zeta \downarrow_{\infty+1})^{3\infty+1} = (\zeta \downarrow_1)^{3^1} \bar{\wedge} (\zeta \downarrow_2)^{3^2} \bar{\wedge} \dots \bar{\wedge} (\zeta \downarrow_{\infty})^{3^{\infty}} \bar{\wedge} (\zeta \downarrow_{\infty+1})^{3^{\infty+1}}$

Raed Hatamleh, Ahmed Salem Heilat, M.Palanikumar and Abdallah Al-Husban,
Characterization of interaction aggregating operators setting interval-valued Pythagorean
neutrosophic set

$$\begin{aligned}
& \left[\left(\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\varrho_{\partial}^{fa})^2 \right)^2 \right)^{3_{\partial}}} \right)^2 + \left(\sqrt{1 - \left(1 - \left((\varrho_{\alpha+1}^{fa})^2 \right)^2 \right)^{3_1}} \right)^2 \right. \\
& \quad \left. - \left(\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\varrho_{\partial}^{fa})^2 \right)^2 \right)^{3_{\partial}}} \right)^2 \cdot \left(\sqrt{1 - \left(1 - \left((\varrho_{\alpha+1}^{fa})^2 \right)^2 \right)^{3_1}} \right)^2 \right. \\
& \quad \left. - \left[\left(\sqrt{\square_{\partial=1}^{\infty} \left(1 - \left((\varrho_{\partial}^{fa})^2 \right)^2 \right)^{3_{\partial}}} - \square_{\partial=1}^{\infty} \left(1 - \left((\varangle_{\partial}^{ta} + \varrho_{\partial}^{fa})^2 \right)^2 \right)^{3_{\partial}} \right)^2 \right. \right. \\
& \quad \left. \left. - \left(\sqrt{\left(1 - \left((\varrho_{\alpha+1}^{fa})^2 \right)^2 \right)^{3_1}} - \left(1 - \left((\varangle_{\alpha+1}^{ta} + \varrho_{\alpha+1}^{fa})^2 \right)^2 \right)^{3_1} \right)^2 \right] \right]^{2'} \\
& \sqrt{\left[\left(\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\varrho_{\partial}^{fb})^2 \right)^2 \right)^{3_{\partial}}} \right)^2 + \left(\sqrt{1 - \left(1 - \left((\varrho_{\alpha+1}^{fb})^2 \right)^2 \right)^{3_1}} \right)^2 \right.} \\
& \quad \left. - \left(\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\varrho_{\partial}^{fb})^2 \right)^2 \right)^{3_{\partial}}} \right)^2 \cdot \left(\sqrt{1 - \left(1 - \left((\varrho_{\alpha+1}^{fb})^2 \right)^2 \right)^{3_1}} \right)^2 \right. \\
& \quad \left. - \left[\left(\sqrt{\square_{\partial=1}^{\infty} \left(1 - \left((\varrho_{\partial}^{fb})^2 \right)^2 \right)^{3_{\partial}}} - \square_{\partial=1}^{\infty} \left(1 - \left((\varangle_{\partial}^{tb} + \varrho_{\partial}^{fb})^2 \right)^2 \right)^{3_{\partial}} \right)^2 \right. \right. \\
& \quad \left. \left. - \left(\sqrt{\left(1 - \left((\varrho_{\alpha+1}^{fb})^2 \right)^2 \right)^{3_1}} - \left(1 - \left((\varangle_{\alpha+1}^{tb} + \varrho_{\alpha+1}^{fb})^2 \right)^2 \right)^{3_1} \right)^2 \right] \right]^{2'} \\
& = \left[\left(\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\sigma_{\partial}^{ia})^2 \right)^2 \right)^{3_{\partial}}} \right)^2 + \left(\sqrt{1 - \left(1 - \left((\sigma_{\alpha+1}^{ia})^2 \right)^2 \right)^{3_1}} \right)^2 \right. \\
& \quad \left. - \left(\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\sigma_{\partial}^{ia})^2 \right)^2 \right)^{3_{\partial}}} \right)^2 \cdot \left(\sqrt{1 - \left(1 - \left((\sigma_{\alpha+1}^{ia})^2 \right)^2 \right)^{3_1}} \right)^2 \right. \\
& \quad \left. - \left[\left(\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\sigma_{\partial}^{ib})^2 \right)^2 \right)^{3_{\partial}}} \right)^2 + \left(\sqrt{1 - \left(1 - \left((\sigma_{\alpha+1}^{ib})^2 \right)^2 \right)^{3_1}} \right)^2 \right. \right. \\
& \quad \left. \left. - \left(\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\sigma_{\partial}^{ib})^2 \right)^2 \right)^{3_{\partial}}} \right)^2 \cdot \left(\sqrt{1 - \left(1 - \left((\sigma_{\alpha+1}^{ib})^2 \right)^2 \right)^{3_1}} \right)^2 \right] \right. \\
& \quad \left. - \left[\left(\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\varrho_{\partial}^{fa})^2 \right)^2 \right)^{3_{\partial}}} \right)^2 + \left(\sqrt{1 - \left(1 - \left((\varrho_{\alpha+1}^{fa})^2 \right)^2 \right)^{3_1}} \right)^2 \right. \right. \\
& \quad \left. \left. - \left(\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\varrho_{\partial}^{fa})^2 \right)^2 \right)^{3_{\partial}}} \right)^2 \cdot \left(\sqrt{1 - \left(1 - \left((\varrho_{\alpha+1}^{fa})^2 \right)^2 \right)^{3_1}} \right)^2 \right. \right. \\
& \quad \left. \left. - \left[\left(\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\varrho_{\partial}^{fb})^2 \right)^2 \right)^{3_{\partial}}} \right)^2 + \left(\sqrt{1 - \left(1 - \left((\varrho_{\alpha+1}^{fb})^2 \right)^2 \right)^{3_1}} \right)^2 \right. \right. \right. \\
& \quad \left. \left. - \left(\sqrt{1 - \square_{\partial=1}^{\infty} \left(1 - \left((\varrho_{\partial}^{fb})^2 \right)^2 \right)^{3_{\partial}}} \right)^2 \cdot \left(\sqrt{1 - \left(1 - \left((\varrho_{\alpha+1}^{fb})^2 \right)^2 \right)^{3_1}} \right)^2 \right] \right]^{2'}
\end{aligned}$$

Raed Hatamleh, Ahmed Salem Heilat, M.Palanikumar and Abdallah Al-Husban,
 Characterization of interaction aggregating operators setting interval-valued Pythagorean
 neutrosophic set

$$= \left[\begin{array}{c} \left[\sqrt{\square_{\partial=1}^{\infty+1} \left(1 - \left((\varrho_1^{fa})^2 \right)^{2^{\partial}} \right)} - \square_{\partial=1}^{\infty+1} \left(1 - \left((\angle_1^{ta} + \varrho_1^{fa})^2 \right)^{2^{\partial}} \right) \right], \\ \left[\sqrt{\square_{\partial=1}^{\infty+1} \left(1 - \left((\varrho_1^{fb})^2 \right)^{2^{\partial}} \right)} - \square_{\partial=1}^{\infty+1} \left(1 - \left((\angle_1^{tb} + \varrho_1^{fb})^2 \right)^{2^{\partial}} \right) \right] \\ \left[\sqrt{1 - \square_{\partial=1}^{\infty+1} \left(1 - \left((\sigma_1^{ia})^2 \right)^{2^{\partial}} \right)} , \sqrt{1 - \square_{\partial=1}^{\infty+1} \left(1 - \left((\sigma_1^{ib})^2 \right)^{2^{\partial}} \right)} \right] \\ \left[\sqrt{1 - \square_{\partial=1}^{\infty+1} \left(1 - \left((\varrho_1^{fa})^2 \right)^{2^{\partial}} \right)} , \sqrt{1 - \square_{\partial=1}^{\infty+1} \left(1 - \left((\varrho_1^{fb})^2 \right)^{2^{\partial}} \right)} \right] \end{array} \right]$$

Hence

$$\frac{1}{\zeta} \left(\square_{\partial=1}^{\infty+1} (\zeta \downarrow_{\partial})^{2^{\partial}} \right) = \left[\begin{array}{c} \left(\sqrt{\square_{\partial=1}^{\infty+1} \left(1 - \left((\varrho_{\partial}^{fa})^2 \right)^{2^{\partial}} \right)} - \square_{\partial=1}^{\infty+1} \left(1 - \left((\angle_{\partial}^{ta} + \varrho_{\partial}^{fa})^2 \right)^{2^{\partial}} \right) \right), \\ \left(\sqrt{1 - \square_{\partial=1}^{\infty+1} \left(1 - \left((\sigma_{\partial}^{ia})^2 \right)^{2^{\partial}} \right)} \right) \left(\sqrt{1 - \square_{\partial=1}^{\infty+1} \left(1 - \left((\varrho_{\partial}^{fa})^2 \right)^{2^{\partial}} \right)} \right) \end{array} \right]$$

Corollary 2.13. Let $\downarrow_{\partial} = \langle (\angle_{\partial}^{ta}, \angle_{\partial}^{tb}), (\sigma_{\partial}^{ia}, \sigma_{\partial}^{ib}), (\varrho_{\partial}^{fa}, \varrho_{\partial}^{fb}) \rangle$ be the collection of IPNNs and all are equal. Then the GIPNIWGO $(\downarrow_1, \downarrow_2, \dots, \downarrow_{\infty}) = \downarrow$.

Acknowledgments: The authors would like to thank the Editor-InChief and the anonymous referees for their various suggestions and helpful comments that have led to the improved in the quality and clarity version of the paper.

Conflicts of Interest: The author declares no conflict of interest.

References

1. L. A. Zadeh, Fuzzy sets, Information and control, 8(3), (1965), 338-353.
2. K. Atanassov, Intuitionistic fuzzy sets, Fuzzy sets and Systems, 20(1), (1986), 87-96.
3. R.R. Yager, Pythagorean membership grades in multi criteria decision making, IEEE Trans. Fuzzy Systems, 22, (2014), 958-965.
4. D. Liang and Z. Xu, The new extension of TOPSIS method for multiple criteria decision making with hesitant Pythagorean fuzzy sets, Applied Soft Computing, 60, (2017), 167-179.
5. F. Smarandache, A unifying field in logics, Neutrosophy neutrosophic probability, set and logic, American Research Press, Rehoboth, (1999).
6. Shihadeh, A., Matarneh, K. A. M., Hatamleh, R., Al-Qadri, M. O., & Al-Husban, A., On The Two-Fold Fuzzy n-Refined Neutrosophic Rings For $2 \leq n \leq 3$. Neutrosophic Sets and Systems, 68, (2024), 8-25.
7. Abdallah Shihadeh, Khaled Ahmad Mohammad Matarneh, Raed Hatamleh, Randa Bashir Yousef Hijazeen, Mowafaq Omar Al-Qadri, Abdallah Al-Husban, An Example of Two-Fold Fuzzy Algebras Based On Neutrosophic Real Numbers, Neutrosophic Sets and Systems, 67, (2024), 169-178.

Raed Hatamleh, Ahmed Salem Heilat, M. Palanikumar and Abdallah Al-Husban, Characterization of interaction aggregating operators setting interval-valued Pythagorean neutrosophic set

8. A. Rajalakshmi, Raed Hatamleh, Abdallah Al-Husban, K. Lenin Muthu Kumaran, M. S. Malchijah raj, Various (ζ_1, ζ_2) neutrosophic ideals of an ordered ternary semigroups. Communications on Applied Nonlinear Analysis, 32 (3), (2025), 400-417.
9. Raed Hatamleh, Abdallah Al-Husban, N. Sundarakannan, M. S. Malchijah Raj, Complex cubic intuitionistic fuzzy set applied to subbisemirings of bisemirings using homomorphism. Communications on Applied Nonlinear Analysis, 32 (3), (2025), 418-435.
10. Abubaker, Ahmad A, Hatamleh, Raed, Matarneh, Khaled, Al-Husban, Abdallah, On the Numerical Solutions for Some Neutrosophic Singular Boundary Value Problems by Using (LPM) Polynomials, International Journal of Neutrosophic Science, 25(2), (2024), 197-205.
11. A., Ahmad. , Hatamleh, Raed. , Matarneh, Khaled. , Al-Husban, Abdallah. On the Irreversible k-Threshold Conversion Number for Some Graph Products and Neutrosophic Graphs, International Journal of Neutrosophic Science, 25(2), (2025), 183-196.
12. Raed Hatamleh, Abdallah Al-Husban, K. Sundareswari, G.Balaj, M.Palanikumar, Complex Tangent Trigonometric Approach Applied to (α, β) -Rung Fuzzy Set using Weighted Averaging, Geometric Operators and its Extension. Communications on Applied Nonlinear Analysis, 32 (5), (2025), 133-144.
13. Raed Hatamleh, Abdallah Al-Husban, M. Palanikumar, K. Sundareswari, Different Weighted Operators such as Generalized Averaging and Generalized Geometric based on Trigonometric q-rung Interval-Valued Approach, Communications on Applied Nonlinear Analysis, 32 (5), (2025), 91-101.
14. Abdallah Shihadeh, Raed Hatamleh, M.Palanikumar, Abdallah Al-Husban, New algebraic structures towards different $(, \ell)$ intuitionistic fuzzy ideals and its characterization of an ordered ternary semigroups. Communications on Applied Nonlinear Analysis, 32 (6), (2025), 568-578.
15. T. Senapati, R.R. Yager, Fermatean fuzzy sets. J. Ambient Intell. Humaniz. Comput. 11, (2020), 663-674.
16. Cuong B.C, Kreinovich V, Picture fuzzy sets a new concept for computational intelligence problems, third world congress on information and communication technologies, WICT 2013. IEEE, 1-6.
17. R.Hatamleh, On the Compactness and Continuity of Uryson's Operator in Orlicz Space, International Journal of Neutrosophic Science, 24 (3), (2024), 233-239.
18. R.Hatamleh, On the Numerical Solutions of the Neutrosophic One-Dimensional Sine-Gordon System, International Journal of Neutrosophic Science, 25 (3), (2024), 25-36.
19. Raed Hatamleh, On The Continuous and Differentiable Two-Fold Neutrosophic and Fuzzy Real Functions, Neutrosophic Sets and Systems, 75, (2025), 196-209,
20. R.Hatamleh, V.A.Zolotarev, On the Universal Models of Commutative Systems of Linear Operators. Journal of Mathematical Physics, Analysis, Geometry, 8(3), (2012), 248-259.
21. R.Hatamleh, V.A. Zolotarev. (2017). On the Abstract Inverse Scattering Problem for Traces Class Perturbations. Journal of Mathematical Physics, Analysis, Geometry, 13(1), (2017), 1-32.
22. Raed Hatamleh, On a Novel Topological Space Based on Partially Ordered Ring of Weak Fuzzy Complex Numbers and its Relation with the Partially ordered Neutrosophic Ring of Real Numbers, Neutrosophic Sets and Systems, 78, (2025), 578-590.
23. Ashraf S, Abdullah S and Mahmood T, Spherical fuzzy Dombi aggregation operators and their application in group decision making problems, Journal of Ambient Intelligence and Humanized Computing, (2019), 1-19.
24. Jin, H., Ashraf, S., Abdullah, S., Qiyas, M., Bano, M. and Zeng, S, Linguistic spherical fuzzy aggregation operators and their applications in multi-attribute decision making problems, Mathematics, 7(5), 2019, 413.
25. Rafiq, M., Ashraf, S., Abdullah, S., Mahmood, T. and Muhammad, S, The cosine similarity measures of spherical fuzzy sets and their applications in decision making. Journal of Intelligent & Fuzzy Systems, 36(6), 2019, 6059-6073.
26. Abdallah Al-Husban & Abdul Razak Salleh, Complex fuzzy ring. Proceedings of 2nd International Conference on Computing, Mathematics and Statistics, IEEE, 2015, 241-245.

Raed Hatamleh, Ahmed Salem Heilat, M.Palanikumar and Abdallah Al-Husban,
Characterization of interaction aggregating operators setting interval-valued Pythagorean
neutrosophic set

27. Abdallah Al-Husban & Abdul Razak Salleh, Complex Fuzzy Hyperring Based on Complex Fuzzy Spaces. Proceedings of 2nd Innovation and Analytics Conference & Exhibition (IACE). Vol. 1691. AIP Publishing 2015, 040009-040017.
28. Al-Husban, A., & Salleh, A. R. Complex fuzzy hyper groups based on complex fuzzy spaces. International Journal of Pure and Applied Mathematics, 107(4), (2016), 949-958.
29. Alsarahead, M. O., & Al-Husban, A, Complex multi-fuzzy subgroups. Journal of discrete mathematical sciences and cryptography, 25(8), (2022), 2707-2716.
30. Al-Husban, A, Multi-fuzzy hyper groups. Italian Journal of Pure and Applied Mathematics, 46, (2021), 382-390.
31. Al-Husban, A, Fuzzy Soft Groups based on Fuzzy Space, Wseas Transactions on Mathematics. 21, (2021), 53-57.
32. Al-Husban,. Abdallah, Al-Sharoa, Doaa., Al-Kaseasbeh, Mohammad., & Mahmood, R.M.S, Structures of fibers of groups actions on graphs. Wseas Transactions on Mathematics, 7, (2022), 650-658.
33. Hatamleh, R., Heilat, A. S., Palanikumar, M., & Al-Husban, A. Different operators via weighted averaging and geometric approach using trigonometric neutrosophic interval-valued set and its extension, Neutrosophic Sets and Systems, 80, 2025, 194-213.

Received: Oct 20, 2024. Accepted: Feb 5, 2025