

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/382115137>

Collection of Bi-Univalent Functions Using Bell Distribution Associated With Jacobi Polynomials

Article · July 2024
DOI: 10.54216/IJNS.250121

CITATIONS
0

READS
202

4 authors:



Ala. A. Amourah

Irbid National University

63 PUBLICATIONS 681 CITATIONS

SEE PROFILE



Tariq Al-Hawary

Al-Balqa' Applied University

62 PUBLICATIONS 618 CITATIONS

SEE PROFILE



Feras Yousef

University of Jordan

51 PUBLICATIONS 829 CITATIONS

SEE PROFILE



Jamal Salah

A'Sharqiyah University

82 PUBLICATIONS 414 CITATIONS

SEE PROFILE



Collection of Bi-Univalent Functions Using Bell Distribution Associated With Jacobi Polynomials

Ala Amourah^{1,4 *}, Tariq Al-Hawary^{2,4}, Feras Yousef^{3,4}, Jamal Salah^{5 *}

¹Mathematics Education Program, Faculty of Education and Arts, Sohar University, Sohar 3111, Oman.

²Department of Applied Science, Ajloun College, Al-Balqa Applied University, Ajloun 26816. Jordan.

³Department of Mathematics, The University of Jordan, Amman 11942, Jordan.

⁴Jadara Research Center, Jadara University, Irbid 21110, Jordan.

⁵College of Applied and Health Sciences, A'Sharqiyah University, Post Box No. 42, Post Code No. 400 Ibra, Sultanate of Oman.

Emails: AAmourah@su.edu.om; tariq-amh@bau.edu.jo; fyousef@ju.edu.jo; damous73@yahoo.com

Abstract

The aim of this study is to present novel collections of bi-univalent functions, which are characterized using the Bell Distribution. These collections are delineated through the application of Jacobi polynomials. We have established bounds for the Taylor-Maclaurin coefficients, particularly $|a_2|$ and $|a_3|$. Additionally, we have investigated the Fekete-Szegő functional issues pertinent to functions within these subclasses. By concentrating on particular parameters in our principal findings, we have identified numerous new insights.

Keywords: Jacobi polynomials; analytic functions; univalent functions; bi-univalent functions; Fekete-Szegő problem.

1 Introduction and preliminaries

Legendre introduced orthogonal polynomials in 1784.¹ These polynomials are frequently employed for solving ordinary differential equations under specific model constraints. Additionally, they have a crucial role in approximation theory.²

Two polynomials, Y_n and Y_m , are said to be orthogonal if they have orders n and m respectively

$$\int_{\epsilon}^{\iota} Y_n(x) Y_m(x) v(x) dx = 0, \quad \text{for } n \neq m. \quad (1)$$

Assuming $v(x)$ is non-negative within the interval (ϵ, ι) , all polynomials of finite order $Y_n(x)$ possess a clearly defined integral. Jacobi polynomials belong to the class of orthogonal polynomials.

This paper analytically examines a newly defined subclass $\mathfrak{G}_{\Sigma}(\varrho, \Upsilon, \varsigma, x)$ of bi-univalent functions utilizing Jacobi polynomials.

Let $\mathcal{A}$ represent the class of analytic functions Θ within the open unit disk $\mathbb{U} = \{z \in \mathbb{C} : |z| < 1\}$, normalized such that $\Theta(0) = 0$ and $\Theta'(0) = 1$. Consequently, each $\Theta \in \mathcal{A}$ can be expressed as:

$$\Theta(z) = z + a_2 z^2 + a_3 z^3 + \cdots, \quad (z \in \mathbb{U}). \quad (2)$$

Moreover, the set of all univalent functions $\Theta \in \mathcal{A}$ is denoted by $\mathcal{S}$ (for more information, refer to³).

The field of geometric function theory can greatly benefit from the powerful tools provided by differential subordination of analytic functions. Miller and Mocanu⁴ originally introduced the concept of differential subordination. Further references can be found in reference.⁵ Miller and Mocanu's book⁶ provides a comprehensive overview of the advancements in this area, including publication dates.

It is well established that for an analytic and univalent function $\Theta(z)$ mapping a domain $\mathbb{D}_1$ onto a domain $\mathbb{D}_2$, the inverse function $g(z) = \Theta^{-1}(z)$ is defined as

$$g(\Theta(z)) = z, \quad (z \in \mathbb{D}_1).$$

This function is analytic and univalent. Additionally, according to,³ every function $\Theta \in \mathcal{S}$ has an inverse map Θ^{-1} which satisfies

$$\Theta^{-1}(\Theta(z)) = z \quad (z \in \mathbb{U}),$$

and

$$\Theta(\Theta^{-1}(\varpi)) = \varpi \quad \left(|\varpi| < r_0(\Theta); r_0(\Theta) \geq \frac{1}{4} \right).$$

In fact, the inverse function is given by

$$\Theta^{-1}(\varpi) = \varpi - a_2^2 \varpi + (2a_2^2 - a_3) \varpi^3 - (5a_2^3 - 5a_2 a_3 + a_4) \varpi^4 + \cdots. \quad (3)$$

A function $\Theta \in \mathcal{A}$ is bi-univalent in $\mathbb{U}$ if both $\Theta(z)$ and $\Theta^{-1}(z)$ are univalent in $\mathbb{U}$. Let Σ represent the class of bi-univalent functions in $\mathbb{U}$ as defined in (2). For further details on the class Σ , refer to.⁷⁻¹⁹

2 Bell Distribution and Jacobi polynomial

In 2018, Castellares et al. introduced the Bell distribution,²⁰ which is suitable for count data with over-dispersion. The Bell distribution is an improvement over the Bell numbers.^{21,22} The probability density function of a discrete random variable X , which follows the Bell distribution, is expressed as:

$$\mathcal{P}(X = m) = \frac{\mathfrak{T}^m e^{(-\mathfrak{T}^2)+1} \mathcal{B}_m}{m!}; \quad m = 1, 2, 3, \cdots, \quad (4)$$

where $\mathcal{B}_m = \frac{1}{e} \sum_{k=0}^{\infty} \frac{k^m}{m!}$ is the Bell numbers, $m \geq 2$, and $\mathfrak{T} > 0$.

Example of the Bell numbers are $\mathcal{B}_2 = 2, \mathcal{B}_3 = 5, \mathcal{B}_4 = 15$ and $\mathcal{B}_5 = 52$.

Now, we will introduce a new power series whose coefficients represent the probabilities of the Bell distribution

$$\mathcal{B}(\mathfrak{T}, z) = z + \sum_{n=2}^{\infty} \frac{\mathfrak{T}^{n-1} \mathcal{B}_n}{(n-1)! e^{\mathfrak{T}^2-1}} z^n, \quad (z \in \mathbb{U}), \quad (5)$$

where $\mathfrak{T} > 0$.

Now, we considered the linear operator $\mathbb{P}_{\mathfrak{T}} : \mathcal{A} \rightarrow \mathcal{A}$ defined by the convolution (or Hadamard product)

$$\mathbb{P}_{\mathfrak{T}}\Theta(z) = \mathcal{B}(\mathfrak{T}, z) * f(z) \quad (6)$$

$$= z + \sum_{n=2}^{\infty} \frac{\mathfrak{T}^{n-1} e^{1-\mathfrak{T}^2} \mathcal{B}_n}{(n-1)!} a_n z^n, \quad (z \in \mathbb{U}), \quad (7)$$

$$= z + \frac{2\mathfrak{T}}{e^{\mathfrak{T}^2-1}} a_2 z^2 + \frac{5\mathfrak{T}^2}{2e^{\mathfrak{T}^2-1}} a_3 z^3 + \frac{15\mathfrak{T}^3}{3!e^{\mathfrak{T}^2-1}} a_4 z^4 + \dots \quad (8)$$

For n , $n + \vartheta$, $n + \varsigma$ are nonnegative integers, a generating function of Jacobi polynomials is defined by

$$J_n(x, z) = 2^{\mathfrak{T}+\varsigma} R^{-1} (1 - x + R)^{-\mathfrak{T}} (1 + x + R)^{-\varsigma},$$

where $R = R(x, z) = (1 - 2zx + x^2)^{0.5}$, $\mathfrak{T} > -1$, $\varsigma > -1$, $x \in [-1, 1]$ and $z \in \mathbb{U}$, (see²³).

For a fixed x , the function $J_n(x, z)$ is analytic in $\mathbb{U}$, allowing it to be represented by a Taylor series expansion as follows:

$$J_n(x, z) = \sum_{n=0}^{\infty} P_n^{(\mathfrak{T}, \varsigma)}(x) z^n, \quad (9)$$

where $P_n^{(\mathfrak{T}, \varsigma)}(x)$ is Jacobi polynomial of degree n .

The Jacobi polynomial $P_n^{(\mathfrak{T}, \varsigma)}(x)$ satisfies a second-order linear homogeneous differential equation:

$$(1 - x^2)y'' + (\varsigma - \mathfrak{T} - (\mathfrak{T} + \varsigma + 2)x)y' + n(n + \mathfrak{T} + \varsigma + 1)y = 0.$$

Jacobi polynomials can alternatively be characterized by the following recursive relationships:

$$P_n^{(\mathfrak{T}, \varsigma)}(x) = (a_{n-1}z - b_{n-1})P_{n-1}^{(\mathfrak{T}, \varsigma)}(x) - c_{n-1}P_{n-2}^{(\mathfrak{T}, \varsigma)}(x), \quad n \geq 2,$$

where

$$a_n = \frac{(2n+\mathfrak{T}+\varsigma+1)(2n+\mathfrak{T}+\varsigma+2)}{2(n+1)(n+\mathfrak{T}+\varsigma+1)}, \quad b_n = \frac{(2n+\mathfrak{T}+\varsigma+1)(\varsigma^2-\mathfrak{T}^2)}{2(n+1)(n+\mathfrak{T}+\varsigma+1)(2n+\mathfrak{T}+\varsigma)} \text{ and } c_n = \frac{(2n+\mathfrak{T}+\varsigma+2)(n+\mathfrak{T})(n+\varsigma)}{(n+1)(n+\mathfrak{T}+\varsigma+1)(2n+\mathfrak{T}+\varsigma)}, \text{ with the initial values}$$

$$P_0^{(\mathfrak{T}, \varsigma)}(x) = 1, \quad P_1^{(\mathfrak{T}, \varsigma)}(x) = (\mathfrak{T} + 1) + \frac{1}{2}(\mathfrak{T} + \varsigma + 2)(x - 1) \text{ and} \quad (10)$$

$$P_2^{(\mathfrak{T}, \varsigma)}(x) = \frac{(\mathfrak{T} + 1)(\mathfrak{T} + 2)}{2} + \frac{1}{2}(\mathfrak{T} + 2)(\mathfrak{T} + \varsigma + 3)(x - 1) + \frac{1}{8}(\mathfrak{T} + \varsigma + 3)(\mathfrak{T} + \varsigma + 4)(x - 1)^2.$$

To begin, we introduce certain special instances of the polynomials $P_n^{(\mathfrak{T}, \varsigma)}$:

1. For $\mathfrak{T} = \varsigma = 0$, we get the Legendre Polynomials.

[title=Legendre Polynomials $L_n(x)$ within the Unit Circle, xlabel=Re(z), ylabel=Im(z), axis equal, grid=major, xmin=-1.2, xmax=1.2, ymin=-1.2, ymax=1.2, samples=100, xtick=-1, -0.5, 0, 0.5, 1, ytick=-1, -0.5, 0, 0.5, 1]
[domain=0:360, samples=200, smooth, thick, gray] (cos(x), sin(x));

[domain=-1:1, thick, blue] (x, 1); $L_0(x)$

[domain=-1:1, thick, red] (x, x); $L_1(x)$

[domain=-1:1, thick, green] (x, (1/2)*(3*x^2 - 1)); $L_2(x)$

[domain=-1:1, thick, orange] (x, (1/2)*(5*x^3 - 3*x)); $L_3(x)$

2. For $\nabla = \varsigma = -0.5$, this results in the Chebyshev Polynomials of the first kind.

[title=Chebyshev Polynomials of the First Kind $T_n(x)$ within the Unit Circle, xlabel=Re(z), ylabel=Im(z), axis equal, grid=major, xmin=-1.2, xmax=1.2, ymin=-1.2, ymax=1.2, samples=100, xtick=-1, -0.5, 0, 0.5, 1, ytick=-1, -0.5, 0, 0.5, 1] [domain=0:360, samples=200, smooth, thick, gray] (cos(x), sin(x));

[domain=-1:1, thick, blue] (x, 1); $T_0(x)$

[domain=-1:1, thick, red] (x, x); $T_1(x)$

[domain=-1:1, thick, green] (x, $2x^2 - 1$); $T_2(x)$

[domain=-1:1, thick, orange] (x, $4x^3 - 3x$); $T_3(x)$

3. For $\nabla = \varsigma = 0.5$, this results in the Chebyshev Polynomials of the second kind.

[title=Chebyshev Polynomials of the Second Kind $U_n(x)$ within the Unit Circle, xlabel=Re(z), ylabel=Im(z), axis equal, grid=major, xmin=-1.2, xmax=1.2, ymin=-1.2, ymax=1.2, samples=100, xtick=-1, -0.5, 0, 0.5, 1, ytick=-1, -0.5, 0, 0.5, 1] [domain=0:360, samples=200, smooth, thick, gray] (cos(x), sin(x));

[domain=-1:1, thick, blue] (x, 1); $U_0(x)$

[domain=-1:1, thick, red] (x, $2x$); $U_1(x)$

[domain=-1:1, thick, green] (x, $4x^2 - 1$); $U_2(x)$

[domain=-1:1, thick, orange] (x, $8x^3 - 4x$); $U_3(x)$

4. For $\nabla = \varsigma$, we get the Gegenbauer Polynomials and each is replaced by $(\nabla - 0.5)$.

In recent years, many scholars have explored the connection between bi-univalent functions and orthogonal polynomials. Some notable references in this field are.²⁴⁻³⁸ However, there seems to be a lack of research on bi-univalent functions related to Jacobi polynomials in existing literature. The main goal of this paper is to initiate an investigation into the properties of bi-univalent functions associated with Jacobi polynomials. To accomplish this, we will consider the following definitions.

3 Definition and Examples

In this section, we introduce and analyze a novel subclass of bi-univalent functions defined within the unit disk. This subclass is established by leveraging the concept of subordination. To construct this new class, we employ the Bell Distribution in conjunction with subordination via Jacobi polynomials.

Definition 3.1. Let $\nabla > -1$, $\varsigma > -1$, $\varrho \geq 0$, $x \in (\frac{1}{2}, 1]$ and $n, n + \vartheta, n + \varsigma$ are nonnegative integers. A function $\Theta \in \Sigma$ given by (2) is said to be in the class $\mathfrak{G}_{\Sigma}(\varrho, \nabla, \varsigma, x)$ if the following subordinations are satisfied:

$$(1 - \varrho) \frac{\mathbb{P}_{\nabla} \Theta(z)}{z} + \varrho (\mathbb{P}_{\nabla} \Theta(z))' \prec J_n(x, z) \quad (11)$$

and

$$(1 - \varrho) \frac{\mathbb{P}_{\nabla} g(\varpi)}{\varpi} + \varrho (\mathbb{P}_{\nabla} g(\varpi))' \prec J_n(x, \varpi), \quad (12)$$

where the function J_n is given by (9) and the function $g(\varpi) = \Theta^{-1}(\varpi)$ is defined by (3).

Example 3.2. Let $\nabla > -1$, $\varsigma > -1$, $x \in (\frac{1}{2}, 1]$ and $n, n + \vartheta, n + \varsigma$ are nonnegative integers. A function $\Theta \in \Sigma$ given by (2) is said to be in the class $\mathfrak{G}_{\Sigma}(0, \nabla, \varsigma, x)$ if the following subordinations are satisfied:

$$\frac{\mathbb{P}_{\nabla} \Theta(z)}{z} \prec J_n(x, z)$$

and

$$\frac{\mathbb{P}\Upsilon g(\varpi)}{\varpi} \prec J_n(x, \varpi),$$

where the function J_n is given by (9) and the function $g(\varpi) = \Theta^{-1}(\varpi)$ is defined by (3).

Example 3.3. Let $\Upsilon > -1$, $\varsigma > -1$, $x \in (\frac{1}{2}, 1]$ and $n, n + \vartheta, n + \varsigma$ are nonnegative integers. A function $\Theta \in \Sigma$ given by (2) is said to be in the class $\mathfrak{G}_\Sigma(1, \Upsilon, \varsigma, x)$ if the following subordinations are satisfied:

$$(\mathbb{P}\Upsilon\Theta(z))' \prec J_n(x, z)$$

and

$$(\mathbb{P}\Upsilon g(\varpi))' \prec J_n(x, \varpi),$$

where the function J_n is given by (9) and the function $g(\varpi) = \Theta^{-1}(\varpi)$ is defined by (3).

4 Coefficient bounds of the subclass $\mathfrak{G}_\Sigma(\varrho, \Upsilon, \varsigma, x)$

This section focuses on determining initial coefficient bounds for the subclass $\mathfrak{G}_\Sigma(\varrho, \Upsilon, \varsigma, x)$.

Theorem 4.1. If $\Theta \in \Sigma$ as defined by equation (2), it belongs to the class $\mathfrak{G}_\Sigma(\varrho, \Upsilon, \varsigma, x)$. Then

$$|a_2| \leq \frac{e^{(\Upsilon^2-1)} |(\Upsilon+1) + \frac{1}{2}(\Upsilon+\varsigma+2)(x-1)| \sqrt{|(\Upsilon+1) + \frac{1}{2}(\Upsilon+\varsigma+2)(x-1)|}}{\sqrt{\Upsilon(x, \Upsilon, \varsigma)}},$$

and

$$|a_3| \leq \frac{e^{2(\Upsilon^2-1)} [(\Upsilon+1) + \frac{1}{2}(\Upsilon+\varsigma+2)(x-1)]^2}{4(1+\varrho)^2\Upsilon^2} + \frac{2e^{\Upsilon^2-1} [(\Upsilon+1) + \frac{1}{2}(\Upsilon+\varsigma+2)(x-1)]}{5(1+2\varrho)\Upsilon^2},$$

where

$$\Upsilon(x, \vartheta, \varsigma) = \Upsilon^2 \left[\begin{array}{c} 5(1+2\varrho) [(\Upsilon+1) + \frac{1}{2}(\Upsilon+\varsigma+2)(x-1)]^2 e^{(\Upsilon^2-1)} \\ -8(1+\varrho)^2 \left(\frac{(\Upsilon+1)(\Upsilon+2)}{2} + \frac{1}{2}(\Upsilon+2)(\Upsilon+\varsigma+3)(x-1) \right. \\ \left. + \frac{1}{8}(\Upsilon+\varsigma+3)(\Upsilon+\varsigma+4)(x-1)^2 \right) \end{array} \right].$$

Proof. Suppose $\Theta \in \mathfrak{G}_\Sigma(\varrho, \Upsilon, \varsigma, x)$. According to equations (11) and (12), for all $z, \varpi \in \mathbb{U}$ and analytic functions r and s with $r(0) = s(0) = 0$ and $|r(z)| < 1$, $|s(\varpi)| < 1$, we can express:

$$(1-\varrho) \frac{\mathbb{P}\Upsilon\Theta(z)}{z} + \varrho(\mathbb{P}\Upsilon\Theta(z))' = J_n(x, r(z)) \quad (13)$$

and

$$(1-\varrho) \frac{\mathbb{P}\Upsilon g(\varpi)}{\varpi} + \varrho(\mathbb{P}\Upsilon g(\varpi))' = J_n(x, s(\varpi)). \quad (14)$$

Thus we have

$$(1-\varrho) \frac{\mathbb{P}\Upsilon\Theta(z)}{z} + \varrho(\mathbb{P}\Upsilon\Theta(z))' = 1 + P_1^{(\vartheta, \varsigma)}(x)b_1z + [P_1^{(\vartheta, \varsigma)}(x)b_2 + P_2^{(\vartheta, \varsigma)}(x)b_1^2]z^2 + \dots \quad (15)$$

and

$$(1-\varrho) \frac{\mathbb{P}\Upsilon g(\varpi)}{\varpi} + \varrho(\mathbb{P}\Upsilon g(\varpi))' = 1 + P_1^{(\vartheta, \varsigma)}(x)d_1\varpi + [P_1^{(\vartheta, \varsigma)}(x)d_2 + P_2^{(\vartheta, \varsigma)}(x)d_1^2]\varpi^2 + \dots \quad (16)$$

It is well known that if

$$|r(z)| = |b_1 z + b_2 z^2 + b_3 z^3 + \dots| < 1, \quad (z \in \mathbb{U})$$

and

$$|s(\varpi)| = |d_1 \varpi + d_2^2 \varpi + d_3^3 \varpi + \dots| < 1, \quad (\varpi \in \mathbb{U}),$$

then

$$|b_j| \leq 1 \text{ and } |d_j| \leq 1 \text{ for all } j \in \mathbb{N}. \quad (17)$$

Comparing the coefficients in (15) and (16), we get

$$\frac{2(1+\varrho)\Upsilon}{e^{\Upsilon^2-1}} a_2 = P_1^{(\Upsilon, \varsigma)}(x) b_1, \quad (18)$$

$$\frac{5(1+2\varrho)\Upsilon^2}{2e^{\Upsilon^2-1}} a_3 = P_1^{(\Upsilon, \varsigma)}(x) b_2 + P_2^{(\Upsilon, \varsigma)}(x) b_1^2, \quad (19)$$

$$-\frac{2(1+\varrho)\Upsilon}{e^{\Upsilon^2-1}} a_2 = P_1^{(\Upsilon, \varsigma)}(x) d_1, \quad (20)$$

and

$$\frac{5(1+2\varrho)\Upsilon^2}{2e^{\Upsilon^2-1}} (2a_2^2 - a_3) = P_1^{(\Upsilon, \varsigma)}(x) d_2 + P_2^{(\Upsilon, \varsigma)}(x) d_1^2. \quad (21)$$

From (18) and (20) it follows that

$$b_1 = -d_1 \quad (22)$$

and

$$\frac{8(1+\varrho)^2\Upsilon^2}{e^{2(\Upsilon^2-1)}} a_2^2 = \left[P_1^{(\Upsilon, \varsigma)}(x) \right]^2 (b_1^2 + d_1^2). \quad (23)$$

If we add (19) and (21), we get

$$\frac{5(1+2\varrho)\Upsilon^2}{e^{\Upsilon^2-1}} a_2^2 = P_1^{(\Upsilon, \varsigma)}(x) (b_2 + d_2) + P_2^{(\Upsilon, \varsigma)}(x) (b_1^2 + d_1^2). \quad (24)$$

Substituting the value of $(b_1^2 + d_1^2)$ from (23) in the right hand side of (24), we get

$$\frac{\Upsilon^2}{e^{\Upsilon^2-1}} \left[5(1+2\varrho) - \frac{P_2^{(\Upsilon, \varsigma)}(x)}{\left[P_1^{(\Upsilon, \varsigma)}(x) \right]^2} \frac{8(1+\varrho)^2}{e^{(\Upsilon^2-1)}} \right] a_2^2 = P_1^{(\Upsilon, \varsigma)}(x) (b_2 + d_2). \quad (25)$$

Using (10), (17) and (25), we find that

$$|a_2| \leq \frac{e^{(\Upsilon^2-1)} \left| (\Upsilon+1) + \frac{1}{2}(\Upsilon+\varsigma+2)(x-1) \right| \sqrt{\left| (\Upsilon+1) + \frac{1}{2}(\Upsilon+\varsigma+2)(x-1) \right|}}{\sqrt{\Upsilon(x, \Upsilon, \varsigma)}},$$

where

$$\Upsilon(x, \Upsilon, \varsigma) = \Upsilon^2 \left[\begin{array}{c} 5(1+2\varrho) \left[(\Upsilon+1) + \frac{1}{2}(\Upsilon+\varsigma+2)(x-1) \right]^2 e^{(\Upsilon^2-1)} \\ -8(1+\varrho)^2 \left(\frac{(\Upsilon+1)(\Upsilon+2)}{2} + \frac{1}{2}(\Upsilon+2)(\Upsilon+\varsigma+3)(x-1) \right. \\ \left. + \frac{1}{8}(\Upsilon+\varsigma+3)(\Upsilon+\varsigma+4)(x-1)^2 \right) \end{array} \right]$$

Moreover, if we subtract (21) from (19), we have

$$\frac{5(1+2\varrho)\Upsilon^2}{e^{\Upsilon^2-1}} (a_3 - a_2^2) = P_1^{(\Upsilon, \varsigma)}(x) (b_2 - d_2) + P_2^{(\Upsilon, \varsigma)}(x) (b_1^2 - d_1^2). \quad (26)$$

Then, in view of (23) and (26) becomes

$$a_3 = \frac{e^{2(\tau^2-1)} \left[P_1^{(\tau, \varsigma)}(x) \right]^2}{8(1+\varrho)^2 \tau^2} (b_1^2 + d_1^2) + \frac{e^{\tau^2-1}}{5(1+2\varrho)\tau^2} P_1^{(\tau, \varsigma)}(x) (b_2 - d_2).$$

Thus applying (10) and (17), we get

$$|a_3| \leq \frac{e^{2(\tau^2-1)} \left[(\tau + 1) + \frac{1}{2}(\tau + \varsigma + 2)(x - 1) \right]^2}{4(1+\varrho)^2 \tau^2} + \frac{2e^{\tau^2-1} \left[(\tau + 1) + \frac{1}{2}(\tau + \varsigma + 2)(x - 1) \right]}{5(1+2\varrho)\tau^2}.$$

Which completes the proof. $\square$

5 Fekete–Szegő problem for the subclass $\mathfrak{G}_\Sigma(\varrho, \tau, \varsigma, x)$

A prominent issue concerning coefficients of univalent analytic functions is the Fekete–Szegő inequality. Initially proposed by,³⁹ it asserts that for $\Theta \in \Sigma$, where Σ denotes

$$|a_3 - \tau a_2^2| \leq 1 + 2e^{-2\tau/(1-\mu)}.$$

When τ is real, this bound is sharp.

In this section, for functions in the class $\mathfrak{G}_\Sigma(\varrho, \tau, \varsigma, x)$ we provide Fekete–Szegő inequalities.

Theorem 5.1. Let $\Theta \in \Sigma$ given by (2) belongs to the class $\mathfrak{G}_\Sigma(\varrho, \tau, \varsigma, x)$. Then

$$|a_3 - \tau a_2^2| \leq \begin{cases} \frac{2e^{(\tau^2-1)} \left| (\tau+1) + \frac{1}{2}(\tau+\varsigma+2)(x-1) \right|}{5(1+2\varrho)\tau^2}, & |\tau - 1| \leq k(x) \\ \frac{2(1-\tau)e^{2(\tau^2-1)} \left[P_1^{(\vartheta, \varsigma)}(x) \right]^3}{\tau^2 \left[5(1+2\varrho) \left[P_1^{(\tau, \varsigma)}(x) \right]^2 e^{(\tau^2-1)} - 8(1+\varrho)^2 P_2^{(\tau, \varsigma)}(x) \right]}, & |\tau - 1| \geq k(x) \end{cases}$$

where

$$k(x) = \left| 1 - \frac{8(1+\varrho)^2 P_2^{(\tau, \varsigma)}(x)}{5(1+2\varrho)e^{(\tau^2-1)} \left[P_1^{(\vartheta, \varsigma)}(x) \right]^2} \right|.$$

Proof. From (25) and (26)

$$\begin{aligned} a_3 - \tau a_2^2 &= \frac{(1-\tau)e^{2(\tau^2-1)} \left[P_1^{(\tau, \varsigma)}(x) \right]^3 (b_2 + d_2)}{\tau^2 \left[5(1+2\varrho) \left[P_1^{(\tau, \varsigma)}(x) \right]^2 e^{(\tau^2-1)} - 8(1+\varrho)^2 P_2^{(\tau, \varsigma)}(x) \right]} + \frac{e^{\tau^2-1}}{5(1+2\varrho)\tau^2} P_1^{(\tau, \varsigma)}(x) (b_2 - d_2) \\ &= P_1^{(\tau, \varsigma)}(x) e^{(\tau^2-1)} \left[\left(h(\tau) + \frac{1}{5(1+2\varrho)\tau^2} \right) b_2 + \left(h(\tau) - \frac{1}{5(1+2\varrho)\tau^2} \right) d_2 \right], \end{aligned}$$

where

$$h(\tau) = \frac{(1-\tau)e^{(\tau^2-1)} \left[P_1^{(\vartheta, \varsigma)}(x) \right]^2}{\tau^2 \left[5(1+2\varrho) \left[P_1^{(\tau, \varsigma)}(x) \right]^2 e^{(\tau^2-1)} - 8(1+\varrho)^2 P_2^{(\tau, \varsigma)}(x) \right]},$$

Then, in view of (10), we have

$$|a_3 - \tau a_2^2| \leq \begin{cases} \frac{2e^{(\tau^2-1)} |P_1^{(\tau, \varsigma)}(x)|}{5(1+2\varrho)\tau^2} & 0 \leq |h(\tau)| \leq \frac{1}{5(1+2\varrho)\tau^2}, \\ 2e^{(\tau^2-1)} |P_1^{(\tau, \varsigma)}(x)| |h(\tau)| & |h(\tau)| \geq \frac{1}{5(1+2\varrho)\tau^2}. \end{cases}$$

Which completes the proof. $\square$

6 Corollaries and Consequences

We use our main results in this section to derive each of the new corollaries and implications that follow.

Corollary 6.1. Let $\Theta \in \Sigma$ given by (2) belongs to the class $\mathfrak{G}_\Sigma(0, \tau, \varsigma, x)$. Then

$$|a_2| \leq \frac{e^{(\tau^2-1)} |(\tau+1) + \frac{1}{2}(\tau+\varsigma+2)(x-1)| \sqrt{|(\tau+1) + \frac{1}{2}(\tau+\varsigma+2)(x-1)|}}{\sqrt{\Upsilon(x, \tau, \varsigma)}},$$

$$|a_3| \leq \frac{e^{2(\tau^2-1)} [(\tau+1) + \frac{1}{2}(\tau+\varsigma+2)(x-1)]^2}{4\tau^2} + \frac{2e^{\tau^2-1} [(\tau+1) + \frac{1}{2}(\tau+\varsigma+2)(x-1)]}{5\tau^2},$$

and

$$|a_3 - \tau a_2^2| \leq \begin{cases} \frac{2e^{(\tau^2-1)} |(\tau+1) + \frac{1}{2}(\tau+\varsigma+2)(x-1)|}{5\tau^2}, & |\tau-1| \leq k(x) \\ \frac{2(1-\tau)e^{2(\tau^2-1)} [P_1^{(\tau, \varsigma)}(x)]^3}{\tau^2 [5[P_1^{(\tau, \varsigma)}(x)]^2 e^{(\tau^2-1)} - 8P_2^{(\tau, \varsigma)}(x)]}, & |\tau-1| \geq k(x) \end{cases}$$

where

$$\Upsilon(x, \tau, \varsigma) = \tau^2 \left[-8 \left(\frac{5 [(\tau+1) + \frac{1}{2}(\tau+\varsigma+2)(x-1)]^2 e^{(\tau^2-1)}}{(\tau+1)(\tau+2) + \frac{1}{2}(\tau+2)(\tau+\varsigma+3)(x-1)} + \frac{1}{8}(\tau+\varsigma+3)(\tau+\varsigma+4)(x-1)^2 \right) \right].$$

and

$$k(x) = \left| 1 - \frac{8P_2^{(\tau, \varsigma)}(x)}{5e^{(\tau^2-1)} [P_1^{(\tau, \varsigma)}(x)]^2} \right|$$

Corollary 6.2. Let $\Theta \in \Sigma$ given by (2) belongs to the class $\mathfrak{G}_\Sigma(1, \tau, \varsigma, x)$. Then

$$|a_2| \leq \frac{e^{(\tau^2-1)} |(\tau+1) + \frac{1}{2}(\tau+\varsigma+2)(x-1)| \sqrt{|(\tau+1) + \frac{1}{2}(\tau+\varsigma+2)(x-1)|}}{\sqrt{\Upsilon(x, \tau, \varsigma)}},$$

$$|a_3| \leq \frac{e^{2(\tau^2-1)} [(\tau+1) + \frac{1}{2}(\tau+\varsigma+2)(x-1)]^2}{16\tau^2} + \frac{2e^{\tau^2-1} [(\tau+1) + \frac{1}{2}(\tau+\varsigma+2)(x-1)]}{15\tau^2},$$

and

$$|a_3 - \tau a_2^2| \leq \begin{cases} \frac{2e^{(\tau^2-1)} |(\tau+1) + \frac{1}{2}(\tau+\varsigma+2)(x-1)|}{15\tau^2}, & |\tau-1| \leq k(x) \\ \frac{2(1-\tau)e^{2(\tau^2-1)} [P_1^{(\tau, \varsigma)}(x)]^3}{\tau^2 [15[P_1^{(\tau, \varsigma)}(x)]^2 e^{(\tau^2-1)} - 32P_2^{(\tau, \varsigma)}(x)]}, & |\tau-1| \geq k(x) \end{cases}$$

where

$$\Upsilon(x, \Upsilon, \varsigma) = \Upsilon^2 \left[\begin{array}{c} 15 \left[(\Upsilon + 1) + \frac{1}{2}(\Upsilon + \varsigma + 2)(x - 1) \right]^2 e^{(\Upsilon^2 - 1)} \\ -32 \left(\frac{(\Upsilon + 1)(\Upsilon + 2)}{2} + \frac{1}{2}(\Upsilon + 2)(\Upsilon + \varsigma + 3)(x - 1) \right. \\ \left. + \frac{1}{8}(\Upsilon + \varsigma + 3)(\Upsilon + \varsigma + 4)(x - 1)^2 \right) \end{array} \right].$$

and

$$k(x) = \left| 1 - \frac{32P_2^{(\Upsilon, \varsigma)}(x)}{15e^{(\Upsilon^2 - 1)} [P_1^{(\Upsilon, \varsigma)}(x)]^2} \right|$$

7 Concluding Remark

In this study, we examined the concerns related to coefficients associated with the newly introduced subclasses $\mathfrak{G}_\Sigma(\varrho, \Upsilon, \varsigma, x)$, $\mathfrak{G}_\Sigma(0, \Upsilon, \varsigma, x)$, and $\mathfrak{G}_\Sigma(1, \Upsilon, \varsigma, x)$ within the class of bi-univalent functions defined in the open unit disk $\mathbb{U}$. The definitions of these subclasses are provided in the text. Our investigation focuses on estimating the Fekete-Szegő functional problems and the Maclaurin coefficients $|a_2|$ and $|a_3|$ for functions within different subclasses of bi-univalent functions. Furthermore, by specializing the parameters in our main results, we have made several novel discoveries.

References

- [1] A. Legendre, *Recherches sur laattraction des sphéroides homogènes*, Mémoires présentes par divers savants a laAcadémie des Sciences de laInstitut de France, Paris, 10 (1785), 411-434.
- [2] H. Bateman, *Higher Transcendental Functions*. McGraw-Hill, 1953.
- [3] P.L. Duren, *Univalent Functions*. Grundlehren der Mathematischen Wissenschaften, Band 259, Springer-Verlag, New York, Berlin, Heidelberg and Tokyo, 1983.
- [4] S. S. Miller, P. T. Mocanu, *Second Order Differential Inequalities in the Complex Plane*. J. Math. Anal. Appl. 65 (1978), 289–305.
- [5] S. S. Miller, P. T. Mocanu, *Differential Subordinations and Univalent Functions*. Mich. Math. J. 28 (1981), 157–172.
- [6] S. S. Miller, P. T. Mocanu, *Differential Subordinations. Theory and Applications*. Marcel Dekker, Inc.: New York, NY, USA, 2000.
- [7] S. Bulut, *Coefficient estimates for a class of analytic and bi-univalent functions*, Novi Sad Journal of Mathematics 43 (2013), 59–65.
- [8] B.A. Frasin, *Coefficient bounds for certain classes of bi-univalent functions*, Hacettepe Journal of Mathematics and Statistics 43.3 (2014), 383-389.
- [9] B. A. Frasin, M. K. Aouf, *New subclasses of bi-univalent functions*, Applied Mathematics Letters 24 (2011), 1569–1573.
- [10] Tariq Al-Hawary, Ala Amourah, Abdullah Alsoboh, and Omar Alsalhi. *A new comprehensive subclass of analytic bi-univalent functions related to gegenbauer polynomials*. Symmetry 15.3 (2023): 576.
- [11] G. Murugusundaramoorthy, N. Magesh, V. Prameela, *Coefficient bounds for certain subclasses of bi-univalent function*, Abstract and Applied Analysis, Volume 2013, Article ID 573017, 3 pages.
- [12] Z. Peng, G. Murugusundaramoorthy, T. Janani, *Coefficient estimate of bi-univalent functions of complex order associated with the Hohlov operator*, Journal of Complex Analysis, Volume 2014, Article ID 693908, 6 pages.

- [13] A. A. Amourah and Feras Yousef, *Some properties of a class of analytic functions involving a new generalized differential operator*. Boletim da Sociedade Paranaense de Matematica 38.6 (2020), 33-42.
- [14] F. Yousef, B.A. Frasin, T. Al-Hawary, *Fekete-Szegő inequality for analytic and bi-univalent functions subordinate to Chebyshev polynomials*. Filomat 32.9 (2018), 3229-3236.
- [15] F. Yousef, T. Al-Hawary, G. Murugusundaramoorthy, *Fekete-Szegő functional problems for some subclasses of bi-univalent functions defined by Frasin differential operator*. Afrika Matematika 30.3-4 (2019), 495–503.
- [16] F. Yousef, S. Alroud, M. Illafe, *New subclasses of analytic and bi-univalent functions endowed with coefficient estimate problems*. Analysis and Mathematical Physics 11. 2 (2021), 1-12.
- [17] F. Yousef, A. Amourah, B.A. Frasin, T. Bulboac, *An avant-Garde construction for subclasses of analytic bi-univalent functions*. Axioms 11.6 (2022), 267.
- [18] Abdullah Alsoboh, Muratlar, and Mucahit Buyankara. *Fekete-Szegő Inequality for a Subclass of Bi-Univalent Functions Linked to q -Ultraspherical Polynomials*. Contemporary Mathematics (2024), 2366-2380.
- [19] Abdullah Alsoboh and Georgia Irina Oros. *A Class of Bi-Univalent Functions in a Leaf-Like Domain Defined through Subordination via q -Calculus*. Mathematics 12.10 (2024), 1594.
- [20] F. Castellares, S. L. Ferrari, A. J. Lemonte, *On the Bell distribution and its associated regression model for count data*, Applied Mathematical Modelling 56 (2018), 172-185.
- [21] E. T. Bell, *Exponential polynomials*, Annals of Mathematics 35 (1934),258-277 .
- [22] E. T. Bell, *Exponential numbers*, The American Mathematical Monthly 41.7 (1934), 411-419.
- [23] M. Marcokova, V. Guldán, *Jacobi Polynomials and Some Related Functions*. In Mathematical Methods in Engineering (pp. 219-227). Springer Netherlands.
- [24] A. A. Amourah and Mohamed Illafe. *A Comprehensive Subclass of Analytic and Bi-Univalent Functions Associated with Subordination*.Palestine Journal of Mathematics 9.1 (2020).
- [25] A. Amourah, T. Al-Hawary, B. A. Frasin, *Application of Chebyshev polynomials to certain class of bi-Bazilevič functions of order $\alpha + i\beta$* . Afrika Matematika 32 .5 (2021), 1059-1066.
- [26] A. Amourah, M. Alomari, F. Yousef, A. Alsoboh, *Consolidation of a Certain Discrete Probability Distribution with a Subclass of Bi-Univalent Functions Involving Gegenbauer Polynomials*. Mathematical Problems in Engineering 2022 (2022).
- [27] A. Amourah, B. A. Frasin, M. Ahmad, F. Yousef, *Exploiting the Pascal distribution series and Gegenbauer polynomials to construct and study a new subclass of analytic bi-univalent functions*. Symmetry 14.1 (2022), 147.
- [28] T. Al-Hawary, A. Amourah, B. A. Frasin, *Fekete-Szegő inequality for bi-univalent functions by means of Horadam polynomials*. Boletín de la Sociedad Matematica Mexicana 27. 3 (2021), 1-12.
- [29] M. Illafe, A. Amourah, M. Haji Mohd, *Coefficient estimates and Fekete-Szegő functional inequalities for a certain subclass of analytic and bi-univalent functions*. Axioms 11.4 (2022), 147.
- [30] A. Hussen, M. Illafe, *Coefficient Bounds for a Certain Subclass of Bi-Univalent Functions Associated with Lucas-Balancing Polynomials*. Mathematics 11.24 (2023), 4941.
- [31] A. Hussen, A. Zeyani, *Coefficients and Fekete-Szegő Functional Estimations of Bi-Univalent Subclasses Based on Gegenbauer Polynomials*. Mathematics 11.13 (2023), 2852.
- [32] A. Hussen, *An Application of the Mittag-Leffler-type Borel Distribution and Gegenbauer Polynomials on a Certain Subclass of Bi-Univalent Functions*. Heliyon 9 (2024), e31469.
- [33] A. Hussen, M. S. Madi, A. M. Abominjil, *Bounding coefficients for certain subclasses of bi-univalent functions related to Lucas-Balancing polynomials*. AIMS Mathematics 9.7 (2024), e31469.

- [34] M. Illafe, F. Yousef, M. Haji Mohd, S. Supramaniam, *Initial Coefficients Estimates and Fekete-Szegő Inequality Problem for a General Subclass of Bi-Univalent Functions Defined by Subordination*. Axioms 12.3 (2023), 235.
- [35] B. A. Frasin, T. Al-Hawary, F. Yousef, I. Aldawish, *On subclasses of analytic functions associated with Struve functions*. Nonlinear Functional Analysis and Applications 27.1 (2022), 99-110.
- [36] A. Amourah, N. Anakira, M. J. Mohammed and M. Jasim, *Jacobi polynomials and bi-univalent functions*. Computer Science, 19(4) (2024), 957-968.
- [37] A. Amourah, A. Alsoboh, D. Breaz and S. M. El-Deeb, *A Bi-Starlike Class in a Leaf-like Domain Defined through Subordination via q -Calculus*. Mathematics, 12(11) (2024), 1735.
- [38] A. Amourah, B. A. Frasin, J. Salah and T. Al-Hawary, *Fibonacci Numbers Related to Some Subclasses of Bi-Univalent Functions*. International Journal of Mathematics and Mathematical Sciences, 2024(1) (2024), 8169496.
- [39] M. Fekete, G. Szegő, *Eine Bemerkung über ungerade schlichte Funktionen*. Journal of the London Mathematical Society 1.2 (1933), 85-89.