

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/385356775>

# New comprehensive two subclasses related to Gregory numbers of analytic bi-univalent functions

**Article** in Journal of Mathematics and Computer Science · October 2024

DOI: 10.22436/jmcs.037.03.07

CITATIONS

0

READS

70

5 authors, including:



**Tariq Al-Hawary**

Al-Balqa Applied University

64 PUBLICATIONS 741 CITATIONS

SEE PROFILE



**Ala. A. Amourah**

Sohar University

88 PUBLICATIONS 863 CITATIONS

SEE PROFILE



**Jamal Salah**

A'Sharqiyah University

104 PUBLICATIONS 476 CITATIONS

SEE PROFILE




**Mohammad Al-khlyleh**

Al-Balqa Applied University

9 PUBLICATIONS 23 CITATIONS

SEE PROFILE



## New comprehensive two subclasses related to Gregory numbers of analytic bi-univalent functions



Tariq Al-Hawary<sup>a,e,\*</sup>, Ala Amourah<sup>b,f</sup>, Jamal Salah<sup>c,\*</sup>, Mohammad Al-khlyleh<sup>a</sup>, Basem Aref Frasin<sup>d</sup>

<sup>a</sup>Department of Applied Science, Ajloun College, Al Balqa Applied University, Ajloun 26816, Jordan.

<sup>b</sup>Mathematics Education Program, Faculty of Education and Arts, Sohar University, Sohar 3111, Oman.

<sup>c</sup>College of Applied and Health Sciences, A'Sharqiyah University, Post Box No. 42, Post Code No. 400 Ibra, Sultanate of Oman.

<sup>d</sup>Faculty of Science, Department of Mathematics, Al al-Bayt University, Mafrq, Jordan.

<sup>e</sup>Jadara Research Center, Jadara University, Irbid 21110, Jordan.

<sup>f</sup>Applied Science Research Center, Applied Science Private University, Amman, Jordan.

### Abstract

In this paper, using subordinations with the functions whose coefficients are Gregory numbers, we present two novel subclasses  $\wp_{\Pi}(\vartheta, \gamma, \beta)$ , and  $\mathcal{F}_{\Pi}(\phi)$  within the bi-univalent function family. We study the estimates  $|a_2|$  and  $|a_3|$  of the Maclaurin coefficients and the Fekete-Szegő inequality regarding functions in every one of these two subclasses. Following the originality of the characterizations and the proofs may encourage additional research on these kinds of similarly defined analytic bi-univalent function subclasses.

**Keywords:** Gregory numbers, analytic, univalent, bi-univalent, Fekete-Szegő.

**2020 MSC:** 30C45.

©2025 All rights reserved.

### 1. Introduction and preliminaries

Let  $\mathcal{H}$  be the class of all analytic functions  $h$  in the open unit disk  $\Omega = \{\tau \in \mathbb{C} : |\tau| < 1\}$  normalized by  $h(0) = h'(0) - 1 = 0$  and has the form:

$$h(\tau) = \tau + \sum_{k=2}^{\infty} a_k \tau^k, \quad (\tau \in \Omega). \quad (1.1)$$

Further, let  $\mathcal{D}$  be the class of univalent functions in the class  $\mathcal{H}$  (for details, see [9]). Now,  $I \prec J$  (the subordination of analytic functions  $I$  and  $J$ ) if for all  $\tau \in \Omega$  there exists a function  $\lambda$  with  $\lambda(0) = 0$  and  $|\lambda(\tau)| < 1$ , such that  $I(\tau) = J(\lambda(\tau))$ . Also, if  $I(0) = J(0)$  and  $I(\Omega) \subset J(\Omega)$  iff  $J$  is univalent in  $\Omega$ , then  $I(\tau) \prec J(\tau)$  (see [11, 15]).

\*Corresponding authors

Email address: [tariq\\_amh@bau.edu.jo](mailto:tariq_amh@bau.edu.jo) (Tariq Al-Hawary)

doi: [10.22436/jmcs.037.03.07](https://doi.org/10.22436/jmcs.037.03.07)

Received: 2024-05-05 Revised: 2024-08-27 Accepted: 2024-09-14

The inverse function is well-known  $J(\tau) = I^{-1}(\tau)$  for analytic and univalent function  $I(\tau) : \mathbb{D}_1 \rightarrow \mathbb{D}_2$  defined by

$$J(I(\tau)) = \tau, \quad (\tau \in \mathbb{D}_1),$$

is a univalent and analytic function. Furthermore (see [9]), there is an inverse map  $I^{-1}$  for any function  $I \in \mathcal{D}$ , such that

$$I^{-1}(I(\tau)) = \tau \quad (\tau \in \Omega),$$

and

$$I(I^{-1}(\lambda)) = \lambda \quad \left( |\lambda| < r_0(I); r_0(I) \geq \frac{1}{4} \right).$$

Actually, the inverse function is written as

$$I^{-1}(\lambda) = \lambda - a_2\lambda^2 + (2a_2^2 - a_3)\lambda^3 - (5a_2^3 - 5a_2a_3 + a_4)\lambda^4 + \dots.$$

A function  $I$  given by (1.1) will be in the class  $\Pi$  ( $\Pi$  the class of bi-univalent functions in  $\Omega$ ) if  $I \in \mathcal{H}$  and both  $I(\tau)$  and  $I^{-1}(\tau)$  are univalent in  $\Omega$ . For details of the class  $\Pi$ , see [1, 7, 16, 19, 21, 24, 25].

Examining the class  $\Pi$ , Brannan and Clunie [6] conjectured that  $a_2 < \sqrt{2}$ , Lewin [13] found that  $a_2 < 1.51$ , and Netanyahu [18] showed that  $\max a_2 = \frac{4}{3}$ . Estimating the coefficient  $a_k$  of  $k \geq 3, k \in \mathbb{N}$ , is still an open problem.

The estimate for functions in the class  $\Pi$  was done by Tan [22] in 1984, and it is  $|a_2| < 1.485$ .

It is widely known that the Fekete-Szegő problem relates to the coefficients of functions in  $\mathcal{D}$ . The first to address this was [10], if  $I \in \Pi$ ,

$$|a_3 - \sigma a_2^2| \leq 1 + 2e^{-2\sigma/(1-\mu)}, \quad \sigma \in \mathbb{R}.$$

Lately, several scholars have begun to examining bi-univalent functions connected to orthogonal polynomials, like [2–5, 12, 23]. Gregory coefficients  $\Theta_k$ , are the numbers  $\frac{1}{2}, \frac{-1}{12}, \frac{1}{24}, \frac{-19}{720}, \frac{3}{160}, \frac{-863}{60.480}, \dots$ . They appear in the expansion of the reciprocal logarithm Maclaurin series

$$\frac{\tau}{\log(\tau+1)} = 1 + \frac{1}{2}\tau - \frac{1}{12}\tau^2 + \frac{1}{24}\tau^3 - \frac{19}{720}\tau^4 + \frac{3}{160}\tau^5 - \frac{863}{60.480}\tau^6 + \dots.$$

James Gregory in 1670 introduced these numbers and subsequently brought back to life by numerous mathematicians and appear in the works of contemporary authors.

The generating function of the Gregory coefficients  $\Theta_k$  (see [20]), are given by

$$\Upsilon(\tau) = \frac{\tau}{\log(\tau+1)} = \sum_{k=0}^{\infty} \Theta_k \tau^k = 1 + \frac{1}{2}\tau - \frac{1}{12}\tau^2 + \frac{1}{24}\tau^3 - \frac{19}{720}\tau^4 + \frac{3}{160}\tau^5 - \frac{863}{60.480}\tau^6 + \dots, \quad \tau \in \Omega. \quad (1.2)$$

Clearly,  $\Theta_k$  for some values of  $k \in \mathbb{N}$  are

$$\Theta_0 = 1, \quad \Theta_1 = \frac{1}{2}, \quad \Theta_2 = \frac{-1}{12}, \quad \Theta_3 = \frac{1}{24}, \quad \Theta_4 = \frac{-19}{720}, \quad \Theta_5 = \frac{3}{160}, \quad \Theta_6 = \frac{-863}{60.480}.$$

The aim of this paper is to make an attempt to improve the initial coefficients  $|a_2|$ ,  $|a_3|$ , and  $|a_3 - \xi a_2^2|$  for certain subclasses of class  $\Pi$ .

## 2. The class $\wp_{\Pi}(\vartheta, \gamma, \beta)$ coefficient bounds

In this section, we examine a subclass  $\wp_{\Pi}(\vartheta, \gamma, \beta)$  using the Gregory coefficients generating functions of analytic bi-univalent functions and obtain initial coefficients  $|a_2|$ ,  $|a_3|$ , and Fekete-Szegő inequality. In our study, the next two lemmas are employed.

**Lemma 2.1** ([8]). If  $\varepsilon \in \mathcal{V}$ , then  $|s_k| \leq 2$ ,  $k \geq 1$  for each  $k$ , where  $\varepsilon$  analytic in  $\Omega$  and has the form  $\varepsilon(\tau) = 1 + s_1\tau + s_2\tau^2 + \dots$ , and sharp for all  $k \in \mathbb{N}$ .

**Lemma 2.2** ([26]). Let  $j, g \in \mathbb{R}$  and  $\tau_1, \tau_2 \in \mathbb{C}$ . If  $|\tau_1|, |\tau_2| < \hbar$ , then

$$|(j+g)\tau_1 + (j-g)\tau_2| \leq \begin{cases} 2|j|\hbar, & \text{for } |j| \geq |g|, \\ 2|g|\hbar, & \text{for } |j| \leq |g|. \end{cases}$$

**Definition 2.3.** Let  $\vartheta \geq 0$ ,  $\beta \geq 1$ ,  $\gamma \in \mathbb{C}$  and  $\operatorname{Re}(\gamma) \geq 0$ . A function  $h \in \Pi$  given by (1.1) is in the class  $\wp_{\Pi}(\vartheta, \gamma, \beta)$  if satisfies the subordinations

$$(1-\vartheta)h'(z) + \vartheta (h'(z))^{\beta} \left( \frac{h(z)}{z} \right)^{\gamma-1} \prec \Upsilon(z) \quad (2.1)$$

and

$$(1-\vartheta)p'(\omega) + \vartheta (p'(\omega))^{\beta} \left( \frac{p(\omega)}{\omega} \right)^{\gamma-1} \prec \Upsilon(\omega), \quad (2.2)$$

where  $\Upsilon(z)$  is given by (1.2) and  $p(\omega) = h^{-1}(\omega)$ .

**Remark 2.4.**

- 1) By utilizing particular values for the parameters  $\vartheta$ ,  $\beta$ , and  $\gamma$  in Definition 2.3, we receive several popular subclasses in  $\mathcal{H}$  studied by several authors.
- 2) If we consider  $h_*(\tau) = \frac{\tau}{1-c\tau}$ ,  $|c| \leq 1$ , then we can check that  $h_*(\tau) \in \wp_{\Pi}(\vartheta, \gamma, \beta)$ , so the class  $\wp_{\Pi}(\vartheta, \gamma, \beta)$  is not empty (see [26, Remark 1]).

**Theorem 2.5.** If  $h \in \wp_{\Pi}(\vartheta, \gamma, \beta)$ . Then

$$|a_2| \leq \min \left\{ \frac{1}{2\vartheta(2\beta + \gamma - 3) + 4}, \frac{1}{\sqrt{\vartheta((\gamma+2)(\gamma-3) + 4\beta(\gamma-1) + 2\beta(2\beta+1)) + \frac{14}{3}(\vartheta(2\beta + \gamma - 3) + 2)^2 + 6}} \right\},$$

$$|a_3| \leq \min \left\{ \frac{1}{4(\vartheta(2\beta + \gamma - 3) + 2)^2} + \frac{1}{2\vartheta(3\beta + \gamma - 4) + 6}, \frac{1}{\vartheta((\gamma+2)(\gamma-3) + 4\beta(\gamma-1) + 2\beta(2\beta+1)) + \frac{14}{3}(\vartheta(2\beta + \gamma - 3) + 2)^2 + 6} + \frac{1}{2\vartheta(3\beta + \gamma - 4) + 6} \right\},$$

and

$$|a_3 - \xi a_2^2| \leq \begin{cases} \frac{1}{2\vartheta(3\beta + \gamma - 4) + 6}, & 0 \leq |\Theta(\xi)| \leq \frac{1}{8\vartheta(3\beta + \gamma - 4) + 24}, \\ \frac{1}{4|\Theta(\xi)|}, & |\Theta(\xi)| \geq \frac{1}{8\vartheta(3\beta + \gamma - 4) + 24}, \end{cases}$$

where

$$\Theta(\xi) = \frac{1 - \xi}{4\vartheta((\gamma+2)(\gamma-3) + 4\beta(\gamma-1) + 2\beta(2\beta+1)) + \frac{56}{3}(\vartheta(2\beta + \gamma - 3) + 2)^2 + 24}.$$

*Proof.* Let  $h \in \wp_{\Pi}(\vartheta, \gamma, \beta)$ . From subordinations (2.1) and (2.2), there exists two analytic functions  $c$  and  $b$  such that  $c(0) = 0 = b(0)$  and  $|c(z)|, |b(\omega)| < 1$ , while

$$(1-\vartheta)h'(z) + \vartheta (h'(z))^{\beta} \left( \frac{h(z)}{z} \right)^{\gamma-1} = \Upsilon(c(z)), \quad z \in \Omega,$$

and

$$(1 - \vartheta)p'(\omega) + \vartheta (p'(\omega))^{\beta} \left( \frac{p(\omega)}{\omega} \right)^{\gamma-1} = \Upsilon(b(\omega)), \quad \omega \in \Omega.$$

So, the function

$$t(z) = \frac{1 + c(z)}{1 - c(z)} = 1 + s_1 z + s_2 z^2 + \cdots \in \mathcal{V},$$

hence

$$c(z) = \frac{s_1}{2} z + \frac{1}{2} \left( s_2 - \frac{s_1^2}{2} \right) z^2 + \frac{1}{2} \left( s_3 - s_1 s_2 + \frac{s_1^3}{4} \right) z^3 + \cdots$$

and

$$\Upsilon(c(z)) = 1 + \frac{s_1}{4} z + \frac{1}{48} (12s_2 - 7s_1^2) z^2 + \frac{1}{192} (17s_1^3 - 56s_1 s_2 + 48s_3) z^3 + \cdots, \quad z \in \Omega.$$

Also, the function

$$l(\omega) = \frac{1 + b(\omega)}{1 - b(\omega)} = 1 + e_1 \omega + e_2 \omega^2 + \cdots \in \mathcal{V},$$

$$b(\omega) = \frac{e_1}{2} \omega + \frac{1}{2} \left( e_2 - \frac{e_1^2}{2} \right) \omega^2 + \frac{1}{2} \left( e_3 - e_1 e_2 + \frac{e_1^3}{4} \right) \omega^3 + \cdots,$$

and

$$\Upsilon(b(\omega)) = 1 + \frac{e_1}{4} \omega + \frac{1}{48} (12e_2 - 7e_1^2) \omega^2 + \frac{1}{192} (17e_1^3 - 56e_1 e_2 + 48e_3) \omega^3 + \cdots, \quad \omega \in \Omega.$$

Consequently, we get

$$\begin{aligned} & (1 - \vartheta)h'(z) + \vartheta (h'(z))^{\beta} \left( \frac{h(z)}{z} \right)^{\gamma-1} \\ &= 1 + \frac{s_1}{4} z + \frac{1}{48} (12s_2 - 7s_1^2) z^2 + \frac{1}{192} (17s_1^3 - 56s_1 s_2 + 48s_3) z^3 + \cdots, \quad z \in \Omega. \end{aligned}$$

and

$$\begin{aligned} & (1 - \vartheta)p'(\omega) + \vartheta (p'(\omega))^{\beta} \left( \frac{p(\omega)}{\omega} \right)^{\gamma-1} \\ &= 1 + \frac{e_1}{4} \omega + \frac{1}{48} (12e_2 - 7e_1^2) \omega^2 + \frac{1}{192} (17e_1^3 - 56e_1 e_2 + 48e_3) \omega^3 + \cdots, \quad \omega \in \Omega. \end{aligned}$$

Comparing the coefficients in the last two equations, we have

$$(\vartheta (2\beta + \gamma - 3) + 2) a_2 = \frac{s_1}{4}, \quad (2.3)$$

$$\left[ \vartheta \left( \frac{(\gamma-1)(\gamma-2)}{2} + 2\beta(\gamma-1) + 2\beta(\beta-1) \right) \right] a_2^2 + [\vartheta (3\beta + \gamma - 4) + 3] a_3 = \frac{1}{48} (12s_2 - 7s_1^2), \quad (2.4)$$

$$-(\vartheta (2\beta + \gamma - 3) + 2) a_2 = \frac{e_1}{4}, \quad (2.5)$$

and

$$\left[ \vartheta \left( \frac{(\gamma-2)(\gamma+3)}{2} + 2\beta(\gamma-1) + 2\beta(\beta+2) - 4 \right) + 6 \right] a_2^2 - [\vartheta (3\beta + \gamma - 4) + 3] a_3 = \frac{1}{48} (12e_2 - 7e_1^2). \quad (2.6)$$

From (2.3) and (2.5) it follows that

$$s_1 = -e_1 \quad (2.7)$$

and

$$s_1^2 + e_1^2 = 32 (\vartheta (2\beta + \gamma - 3) + 2)^2 a_2^2. \quad (2.8)$$

If we add (2.4) to (2.6), we get

$$[\vartheta ((\gamma + 2)(\gamma - 3) + 4\beta(\gamma - 1) + 2\beta(2\beta + 1)) + 6] a_2^2 = \frac{1}{4} (s_2 + e_2) - \frac{7}{48} (s_1^2 + e_1^2). \quad (2.9)$$

Substituting the value of  $s_1^2 + e_1^2$  from (2.8) in (2.9), we have

$$\left\{ [\vartheta ((\gamma + 2)(\gamma - 3) + 4\beta(\gamma - 1) + 2\beta(2\beta + 1)) + 6] + \frac{14}{3} (\vartheta (2\beta + \gamma - 3) + 2)^2 \right\} a_2^2 = \frac{1}{4} (s_2 + e_2). \quad (2.10)$$

By applying Lemma 2.1 and the triangle inequality for the relations (2.3) and (2.10), we get, respectively,

$$|a_2| \leq \frac{1}{2\vartheta (2\beta + \gamma - 3) + 4}$$

and

$$|a_2| \leq \frac{1}{\sqrt{\vartheta ((\gamma + 2)(\gamma - 3) + 4\beta(\gamma - 1) + 2\beta(2\beta + 1)) + \frac{14}{3} (\vartheta (2\beta + \gamma - 3) + 2)^2 + 6}}.$$

Moreover, if we subtract (2.6) from (2.4), we have

$$2 [\vartheta (3\beta + \gamma - 4) + 3] (a_3 - a_2^2) = \frac{1}{4} (s_2 - e_2) - \frac{7}{48} (s_1^2 - e_1^2).$$

Then, in view of (2.7), last equation becomes

$$a_3 = a_2^2 + \frac{s_2 - e_2}{8\vartheta (3\beta + \gamma - 4) + 24}. \quad (2.11)$$

The above equation with (2.3) becomes

$$a_3 = \frac{s_1^2}{16 (\vartheta (2\beta + \gamma - 3) + 2)^2} + \frac{s_2 - e_2}{8\vartheta (3\beta + \gamma - 4) + 24}.$$

By applying Lemma 2.1 and the triangle inequality for the last relation, we get

$$|a_3| \leq \frac{1}{4 (\vartheta (2\beta + \gamma - 3) + 2)^2} + \frac{1}{2\vartheta (3\beta + \gamma - 4) + 6}$$

and using our first assertion with (2.11), it follows that

$$|a_3| \leq \frac{1}{\vartheta ((\gamma + 2)(\gamma - 3) + 4\beta(\gamma - 1) + 2\beta(2\beta + 1)) + \frac{14}{3} (\vartheta (2\beta + \gamma - 3) + 2)^2 + 6} + \frac{1}{2\vartheta (3\beta + \gamma - 4) + 6}.$$

Also, from (2.11) we have

$$\begin{aligned} a_3 - \xi a_2^2 &= \frac{s_2 - e_2}{8\vartheta (3\beta + \gamma - 4) + 24} + (1 - \xi) a_2^2 \\ &= \frac{s_2 - e_2}{8\vartheta (3\beta + \gamma - 4) + 24} \\ &\quad + \frac{(1 - \xi) (s_2 + e_2)}{4\vartheta ((\gamma + 2)(\gamma - 3) + 4\beta(\gamma - 1) + 2\beta(2\beta + 1)) + \frac{56}{3} (\vartheta (2\beta + \gamma - 3) + 2)^2 + 24} \end{aligned}$$

$$= \left( \Theta(\xi) + \frac{1}{8\vartheta(3\beta + \gamma - 4) + 24} \right) s_2 + \left( \Theta(\xi) - \frac{1}{8\vartheta(3\beta + \gamma - 4) + 24} \right) e_2,$$

where

$$\Theta(\xi) = \frac{1 - \xi}{4\vartheta((\gamma + 2)(\gamma - 3) + 4\beta(\gamma - 1) + 2\beta(2\beta + 1)) + \frac{56}{3}(\vartheta(2\beta + \gamma - 3) + 2)^2 + 24}.$$

Then, in view of Lemma 2.1 for  $s_2$  and  $e_2$ , and Lemma 2.2, we obtain

$$|a_3 - \xi a_2^2| \leq \begin{cases} \frac{1}{2\vartheta(3\beta + \gamma - 4) + 6}, & 0 \leq |\Theta(\xi)| \leq \frac{1}{8\vartheta(3\beta + \gamma - 4) + 24}, \\ \frac{1}{4|\Theta(\xi)|}, & |\Theta(\xi)| \geq \frac{1}{8\vartheta(3\beta + \gamma - 4) + 24}, \end{cases}$$

which completes the proof.  $\square$

Fixing  $\vartheta = 0$  or  $\vartheta = \beta = \gamma = 1$  in Theorem 2.5, we get the following coefficient estimates, which is due to [17] in Theorems 1 and 2.

**Corollary 2.6.** *If  $h \in \wp_{\Pi}(\vartheta, \gamma, \beta)$ , then  $|a_2| \leq \sqrt{\frac{3}{74}}$ ,  $|a_3| \leq \frac{23}{111}$ , and*

$$|a_3 - \xi a_2^2| \leq \begin{cases} \frac{1}{6}, & |K(\xi)| \leq \frac{1}{24}, \\ \frac{1}{4|K(\xi)|}, & |K(\xi)| \geq \frac{1}{24}, \end{cases}$$

where  $K(\xi) = \frac{3(1-\xi)}{296}$ .

### 3. The class $\mathcal{F}_{\Pi}(\phi)$ coefficient bounds

In this part, we examine a subclass  $\mathcal{F}_{\Pi}(\phi)$  of analytic bi-univalent functions and obtain initial coefficients  $|a_2|$ ,  $|a_3|$ , and Fekete-Szegő inequality. We utilize the subsequent lemma to establish our results for the class  $\mathcal{F}_{\Pi}(\phi)$ .

**Lemma 3.1** ([14]). *If  $\varepsilon(\tau) = 1 + s_1\tau + s_2\tau^2 + \dots \in \mathcal{V}$ ,  $\tau \in \Omega$ , then there exist some  $\alpha, \delta$  with  $|\alpha|, |\delta| \leq 1$ , such that*

$$2s_2 = s_1^2 + \alpha(4 - s_1^2) \text{ and } 4s_3 = s_1^3 + 2s_1\alpha(4 - s_1^2) - (4 - s_1^2)s_1\alpha^2 + 2(4 - s_1^2)(1 - |\alpha|^2)\delta.$$

**Definition 3.2.** A function  $h \in \Pi$  given by (1.1) is in the class  $\mathcal{F}_{\Pi}(\phi)$ , where  $\phi \in (-\pi, \pi]$ , iff

$$h'(z) + \left( \frac{e^{i\phi} + 1}{2} \right) zh''(z) \prec \Upsilon(z) \quad \text{and} \quad p'(\omega) + \left( \frac{e^{i\phi} + 1}{2} \right) \omega p''(\omega) \prec \Upsilon(\omega), \quad (3.1)$$

where  $\Upsilon(z)$  is given by (1.2) and  $p(\omega) = h^{-1}(\omega)$ .

**Remark 3.3.**

- 1) By utilizing particular values for  $\phi \in (-\pi, \pi]$  in Definition 3.2, we obtain several popular subclasses of  $\mathcal{H}$ .
- 2) If we consider  $h_*(\tau) = \frac{\tau}{1-c\tau}$ ,  $|c| \leq 1$ , then we can check that  $h_*(\tau) \in \mathcal{F}_{\Pi}(\phi)$ , so the class  $\mathcal{F}_{\Pi}(\phi)$  is not empty (see [26, Remark 5]).

**Theorem 3.4.** *If  $h \in \mathcal{F}_{\Pi}(\phi)$ . Then*

$$|a_2| \leq \min \left\{ \frac{1}{\sqrt{2}|e^{i\phi} + 3|}, \frac{1}{\sqrt{6|e^{i\phi} + 2| + \frac{7}{3}|e^{i\phi} + 3|^2}} \right\},$$

$$|a_3| \leq \left\{ \frac{1}{2|e^{i\Phi} + 3|^2} + \frac{1}{6|e^{i\Phi} + 2|} \frac{1}{|33 + 20e^{i\Phi} + \frac{7}{3}e^{i2\Phi}|} + \frac{1}{6|e^{i\Phi} + 2|} \right\},$$

and

$$|a_3 - \rho a_2^2| \leq \begin{cases} \frac{1}{6|e^{i\Phi} + 2|}, & |1 - \rho| \leq \frac{|e^{i\Phi} + 3|^2}{3|e^{i\Phi} + 2|}, \\ \frac{|1 - \rho|}{2|e^{i\Phi} + 3|^2}, & |1 - \rho| \geq \frac{|e^{i\Phi} + 3|^2}{3|e^{i\Phi} + 2|}. \end{cases} \quad (3.2)$$

*Proof.* Let  $h \in \mathcal{F}_\Pi(\Phi)$ . From subordinations (3.1), there exists two analytic functions  $c$  and  $b$  such that  $c(0) = 0 = b(0)$  and  $|c(z)|, |b(\omega)| < 1$ , such that

$$h'(z) + \left( \frac{e^{i\Phi} + 1}{2} \right) zh''(z) = \Upsilon(c(z)), \quad z \in \Omega,$$

and

$$p'(\omega) + \left( \frac{e^{i\Phi} + 1}{2} \right) \omega p''(\omega) = \Upsilon(b(\omega)), \quad \omega \in \Omega.$$

Thus we have

$$h'(z) + \left( \frac{e^{i\Phi} + 1}{2} \right) zh''(z) = 1 + \frac{s_1}{4}z + \frac{1}{48}(12s_2 - 7s_1^2)z^2 + \frac{1}{192}(17s_1^3 - 56s_1s_2 + 48s_3)z^3 + \dots, \quad z \in \Omega,$$

and

$$\begin{aligned} p'(\omega) + \left( \frac{e^{i\Phi} + 1}{2} \right) \omega p''(\omega) &= 1 + \frac{e_1}{4}\omega + \frac{1}{48}(12e_2 - 7e_1^2)\omega^2 \\ &+ \frac{1}{192}(17e_1^3 - 56e_1e_2 + 48e_3)\omega^3 + \dots, \quad \omega \in \Omega. \end{aligned}$$

Comparing the coefficients in last two equations, we have

$$\begin{aligned} (e^{i\Phi} + 3)a_2 &= \frac{s_1}{4}, \\ 3(e^{i\Phi} + 2)a_3 &= \frac{1}{48}(12s_2 - 7s_1^2), \end{aligned} \quad (3.3)$$

$$\begin{aligned} -(e^{i\Phi} + 3)a_2 &= \frac{e_1}{4}, \\ 3(e^{i\Phi} + 2)(2a_2^2 - a_3) &= \frac{1}{48}(12e_2 - 7e_1^2). \end{aligned} \quad (3.4)$$

From (2.3) and (2.5) it follows that

$$s_1 = -e_1 \quad (3.5)$$

and

$$16(e^{i\Phi} + 3)^2 a_2^2 = s_1^2 + e_1^2. \quad (3.6)$$

If we add (3.3) to (3.4), we get

$$6(e^{i\Phi} + 2)a_2^2 = \frac{1}{4}(s_2 + e_2) - \frac{7}{48}(s_1^2 + e_1^2). \quad (3.7)$$

Substituting the value of  $s_1^2 + e_1^2$  from (3.6) in (3.7), we have

$$\left\{ 6(e^{i\Phi} + 2) + \frac{7}{3}(e^{i\Phi} + 3)^2 \right\} a_2^2 = \frac{1}{4}(s_2 + e_2). \quad (3.8)$$

By applying Lemma 2.1 and the triangle inequality for the relations (3.6) and (3.8), we get, respectively,

$$|a_2| \leq \frac{1}{\sqrt{2}|e^{i\Phi} + 3|} \quad \text{and} \quad |a_2| \leq \frac{1}{\sqrt{6|e^{i\Phi} + 2| + \frac{7}{3}|e^{i\Phi} + 3|^2}}.$$

Moreover, if we subtract (3.4) from (3.3), we have

$$6(e^{i\Phi} + 2)(a_3 - a_2^2) = \frac{1}{4}(s_2 - e_2) - \frac{7}{48}(s_1^2 - e_1^2).$$

Then, in view of (3.5), last equation becomes

$$a_3 = a_2^2 + \frac{s_2 - e_2}{24(e^{i\Phi} + 2)}. \quad (3.9)$$

The above equation with (3.6) becomes

$$a_3 = \frac{s_1^2 + e_1^2}{16(e^{i\Phi} + 3)^2} + \frac{s_2 - e_2}{24(e^{i\Phi} + 2)}.$$

By applying Lemma 2.1 and the triangle inequality for the last relation, we get

$$|a_3| \leq \frac{1}{2|e^{i\Phi} + 3|^2} + \frac{1}{6|e^{i\Phi} + 2|}.$$

Similarly, using of (3.8) in relation (3.9) follows that

$$|a_3| \leq \frac{1}{|33 + 20e^{i\Phi} + \frac{7}{3}e^{i2\Phi}|} + \frac{1}{6|e^{i\Phi} + 2|}.$$

Also, using (3.5) and (3.6), we get  $a_2^2 = \frac{s_1^2}{8(e^{i\Phi} + 3)^2}$ . Thus, from (3.9), we have

$$a_3 - \rho a_2^2 = \frac{s_2 - e_2}{24(e^{i\Phi} + 2)} + (1 - \rho)a_2^2 = \frac{s_2 - e_2}{24(e^{i\Phi} + 2)} + (1 - \rho)\frac{s_1^2}{8(e^{i\Phi} + 3)^2}.$$

From Lemma 3.1, we have  $2s_2 = s_1^2 + \alpha(4 - s_1^2)$  and  $2e_2 = e_1^2 + \delta(4 - e_1^2)$ ,  $|\alpha|, |\delta| \leq 1$ , and using (3.5), we obtain

$$s_2 - e_2 = \frac{4 - s_1^2}{2}(\alpha - \delta),$$

and thus

$$a_3 - \rho a_2^2 = \frac{(4 - s_1^2)(\alpha - \delta)}{48(e^{i\Phi} + 2)} + \frac{(1 - \rho)s_1^2}{8(e^{i\Phi} + 3)^2}.$$

Using the triangle inequality, taking  $|\alpha| = \mu$ ,  $|\delta| = \eta$ ,  $\mu, \eta \in [0, 1]$ , and assuming that  $s_1 \in \mathbb{R}$ ,  $s_1 = d \in [0, 2]$ , thus, we get

$$|a_3 - \rho a_2^2| \leq \frac{(4 - d^2)(\mu + \eta)}{48|e^{i\Phi} + 2|} + \frac{|1 - \rho|d^2}{8|e^{i\Phi} + 3|^2}. \quad (3.10)$$

Assume that  $\Psi(d) = \frac{|1 - \rho|d^2}{8|e^{i\Phi} + 3|^2} \geq 0$  and  $\Phi(d) = \frac{4 - d^2}{48|e^{i\Phi} + 2|} \geq 0$ , the relation (3.10) can be rewritten as

$$|a_3 - \rho a_2^2| \leq \Psi(d) + \Phi(d)(\mu + \eta) =: \mathcal{L}(\mu, \eta), \quad \mu, \eta \in [0, 1].$$

Therefore

$$\max\{\mathcal{L}(\mu, \eta) : \mu, \eta \in [0, 1]\} = \mathcal{L}(1, 1) = \Psi(d) + 2\Phi(d) =: K(d), \quad d \in [0, 2],$$

where

$$K(d) = \frac{1}{8|e^{i\phi} + 3|^2} \left( |1 - \rho| - \frac{|e^{i\phi} + 3|^2}{3|e^{i\phi} + 2|} \right) d^2 + \frac{1}{6|e^{i\phi} + 2|}.$$

Since

$$K'(d) = \frac{1}{4|e^{i\phi} + 3|^2} \left( |1 - \rho| - \frac{|e^{i\phi} + 3|^2}{3|e^{i\phi} + 2|} \right) d,$$

it is clear that  $K'(d) \leq 0$  iff  $|1 - \rho| \leq \frac{|e^{i\phi} + 3|^2}{3|e^{i\phi} + 2|}$ , hence, the function  $K$  is decreasing on  $[0, 2]$ , therefore,

$$\max\{K(d) : d \in [0, 2]\} = K(0) = \frac{1}{6|e^{i\phi} + 2|}.$$

Also,  $K'(d) \geq 0$  iff  $|1 - \rho| \geq \frac{|e^{i\phi} + 3|^2}{3|e^{i\phi} + 2|}$ , so,  $K$  is an increasing function over  $[0, 2]$ , so

$$\max\{K(d) : d \in [0, 2]\} = K(2) = \frac{|1 - \rho|}{2|e^{i\phi} + 3|^2}$$

and the estimation (3.2) has been validated. □

**Remark 3.5.** By taking specific values for  $\vartheta$ ,  $\gamma$ , and  $\beta$  in Theorem 2.5, and for  $\phi$  in Theorem 3.4 we get various well-known subclasses of  $\Pi$ . Also, there are several functions  $\Upsilon(z)$  that may produce interesting subclasses of function class  $\Pi$ . As an example, if

$$\Upsilon(z) = \frac{1 + (1 - 2\hbar)z}{1 - z} = 1 + 2(1 - \hbar)z + 2(1 - \hbar)z^2 + \dots \quad (0 \leq \hbar < 1)$$

or

$$\Upsilon(z) = \left( \frac{1 + z}{1 - z} \right)^\hbar = 1 + 2\hbar z + 2\hbar^2 z^2 + \dots \quad (0 < \hbar \leq 1),$$

which gives special cases for our previous results.

## 4. Conclusions

In this study, utilizing the functions whose coefficients are Gregory numbers, for the class of bi-univalent functions we present two new subclasses  $\wp_\Pi(\vartheta, \gamma, \beta)$  and  $\mathcal{F}_\Pi(\phi)$ . We study the estimates  $|a_2|$  and  $|a_3|$  and the Fekete-Szegő inequality regarding the functions in these classes. The originality of the characterizations and the proofs in this research may encourage additional research on these kinds of similarly defined analytic bi-univalent function subclasses.

## 5. Acknowledgments

All authors are thankful to the reviewers of this article, who gave valuable remarks and comments in order to improve the quality of the paper.

## References

- [1] I. Aldawish, T. Al-Hawary, B. A. Frasin, *Subclasses of bi-univalent functions defined by Frasin differential operator*, *Mathematics*, 8 (2020), 11 pages. 1

- [2] T. Al-Hawary, A. Amourah, A. Alsoboh, O. Alsalhi, *A new comprehensive subclass of analytic bi-univalent functions related to gegenbauer polynomials*, Symmetry, **15**, (2023), 11 pages. 1
- [3] T. Al-Hawary, A. Amourah, B. A. Frasin, *Fekete-Szegő inequality for bi-univalent functions by means of Horadam polynomials*, Bol. Soc. Mat. Mex. (3), **27** (2021), 12 pages.
- [4] A. Amourah, T. Al-Hawary, B. A. Frasin, *Application of Chebyshev polynomials to certain class of bi-Bazilevič functions of order  $\alpha + i\beta$* , Afr. Mat., **32** (2021), 1059–1066.
- [5] A. Amourah, B. A. Frasin, T. M. Seoudy, *An Application of Miller–Ross-Type Poisson Distribution on Certain Subclasses of Bi-Univalent Functions Subordinate to Gegenbauer Polynomials  $\alpha + i\beta$* , Mathematics, **10** (2022), 10 pages. 1
- [6] D. A. Brannan, J. G. Clunie, *Aspects of contemporary complex analysis*, Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], London-New York, (1980). 1
- [7] S. Bulut, *Coefficient estimates for a class of analytic and bi-univalent functions*, Novi Sad J. Math., **43** (2013), 59–65. 1
- [8] C. Carathéodory, *Über den Variabilitätsbereich der Koeffizienten von Potenzreihen, die gegebene Werte nicht annehmen*, Math. Ann., **64** (1907), 95–115. 2.1
- [9] P. L. Duren, *Univalent functions*, Springer-Verlag, New York, (1983). 1
- [10] M. Fekete, G. Szegő, *Eine Bemerkung Über Ungerade Schlichte Funktionen*, J. London Math. Soc., **8** (1933), 85–89. 1
- [11] B. A. Frasin, *Subordination results for a class of analytic functions defined by a linear operator*, JIPAM. J. Inequal. Pure Appl. Math., **7** (2006), 7 pages. 1
- [12] B. A. Frasin, T. Al-Hawary, F. Yousef, I. Aldawish, *On subclasses of analytic functions associated with Struve functions*, Nonlinear Funct. Anal. Appl., **27** (2022), 99–110. 1
- [13] M. Lewin, *On a coefficient problem for bi-univalent functions*, Proc. Amer. Math. Soc., **18** (1967), 63–68. 1
- [14] R. J. Libera, E. J. Złotkiewicz, *Coefficient bounds for the inverse of a function with derivative in  $\mathcal{P}$* , Proc. Amer. Math. Soc., **87** (1983), 251–257. 3.1
- [15] S. Miller, P. Mocanu, *Differential Subordination: theory and applications*, CRC Press, New York, (2000). 1
- [16] G. Murugusundaramoorthy, N. Magesh, V. Prameela, *Coefficient bounds for certain subclasses of bi-univalent function*, Abstr. Appl. Anal., **2013** (2013), 3 pages. 1
- [17] G. Murugusundaramoorthy, K. Vijaya, T. Bulboacă, *Initial Coefficient Bounds for Bi-Univalent Functions Related to Gregory Coefficients*, Mathematics, **11** (2023), 1–16. 2
- [18] E. Netanyahu, *The minimal distance of the image boundary from the origin and the second coefficient of a univalent function in  $z < 1$* , Arch. Rational Mech. Anal., **32** (1969), 100–112. 1
- [19] Z. Peng, G. Murugusundaramoorthy, T. Janani, *Coefficient estimate of biunivalent functions of complex order associated with the Hohlov operator*, J. Complex Anal., **2014** (2014), 6 pages. 1
- [20] G. M. Phillips, *Gregory's method for numerical integration*, Amer. Math. Monthly, **79** (1972), 270–274. 1
- [21] H. M. Srivastava, A. K. Mishra, P. Gochhayat, *Certain subclasses of analytic and bi-univalent functions*, Appl. Math. Lett., **23** (2010), 1188–1192. 1
- [22] D. L. Tan, *Coefficient estimates for bi-univalent functions*, Chin. Ann. Math. Ser. A, **5** (1984), 559–568. 1
- [23] K. Vijaya, G. Murugusundaramoorthy, *Bi-Starlike Function of Complex Order Involving Mathieu-Type Series Associated with Telephone Numbers*, Symmetry, **15** (2023), 18 pages. 1
- [24] F. Yousef, T. Al-Hawary, G. Murugusundaramoorthy, *Fekete-Szegő functional problems for some subclasses of bi-univalent functions defined by Frasin differential operator*, Afr. Mat., **30** (2019), 495–503. 1
- [25] F. Yousef, B. A. Frasin, T. Al-Hawary, *Fekete-Szegő inequality for analytic and bi-univalent functions subordinate to Chebyshev polynomials*, Filomat, **32** (2018), 3229–3236. 1
- [26] P. Zaprawa, *Estimates of initial coefficients for bi-univalent functions*, Abstr. Appl. Anal., **2014** (2014), 6 pages. 2.2, 2, 2